



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2021 – 10:03 am BST

PDB ID : 7ORD
Title : The crystal structure of the domain-swapped dimer of onconase (2)
Authors : Merlino, A.; Loreto, D.
Deposited on : 2021-06-05
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	AAA	105	
1	BBB	105	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	AAA	201	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1778 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protein P-30.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	104	Total	C	N	O	S	0	1	0
			832	523	143	157	9			
1	BBB	104	Total	C	N	O	S	0	2	0
			840	527	145	159	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	1	PCA	-	insertion	UNP P22069
BBB	1	PCA	-	insertion	UNP P22069

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	51	Total	O	0	1
			51	51		
4	BBB	46	Total	O	0	1
			46	46		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: Protein P-30

Chain AAA:  76% 23%



- Molecule 1: Protein P-30

Chain BBB:  81% 17%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.75Å 28.54Å 61.48Å 90.00° 94.24° 90.00°	Depositor
Resolution (Å)	58.59 – 2.14	Depositor
% Data completeness (in resolution range)	94.1 (58.59-2.14)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.212 , 0.267	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1778	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, ACT, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	0.68	0/842	0.89	0/1137
1	BBB	0.68	0/850	0.86	0/1148
All	All	0.68	0/1692	0.88	0/2285

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	832	0	819	14	0
1	BBB	840	0	824	19	0
2	AAA	4	0	3	5	0
3	AAA	5	0	0	0	0
4	AAA	51	0	0	2	0
4	BBB	46	0	0	3	0
All	All	1778	0	1646	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:AAA:201:ACT:H3	4:BBB:204:HOH:O	1.81	0.80
2:AAA:201:ACT:H2	1:BBB:10:HIS:CD2	2.20	0.75
1:AAA:97:HIS:HA	2:AAA:201:ACT:H1	1.70	0.73
1:BBB:14:THR:O	1:BBB:16:ASP:N	2.24	0.70
1:BBB:14:THR:C	1:BBB:16:ASP:H	1.94	0.70
2:AAA:201:ACT:CH3	4:BBB:204:HOH:O	2.43	0.63
2:AAA:201:ACT:H2	1:BBB:10:HIS:NE2	2.14	0.62
1:BBB:64:TYR:OH	1:BBB:81:LYS:HD2	2.02	0.60
1:BBB:14:THR:C	1:BBB:16:ASP:N	2.52	0.60
1:BBB:15:ARG:HB3	1:BBB:38:TYR:CD1	2.38	0.59
1:AAA:2:ASP:OD2	4:AAA:301:HOH:O	2.17	0.58
1:AAA:48:CYS:O	1:AAA:93:GLN:HA	2.07	0.55
1:BBB:56:ASN:ND2	1:BBB:89:THR:OG1	2.43	0.52
1:BBB:19:CYS:HB3	1:BBB:77:TYR:HB2	1.92	0.52
1:BBB:50:GLY:H	1:BBB:93:GLN:HG2	1.78	0.48
1:BBB:59:THR:HA	4:BBB:226:HOH:O	2.16	0.46
1:AAA:41:PRO:HD3	1:BBB:14:THR:HG22	1.98	0.45
1:AAA:36:PHE:HB3	1:BBB:12:THR:OG1	2.16	0.45
1:BBB:15:ARG:HB3	1:BBB:38:TYR:CE1	2.51	0.45
1:AAA:56:ASN:HD22	1:AAA:89:THR:HA	1.83	0.44
1:AAA:50:GLY:H	1:AAA:93:GLN:HE21	1.65	0.44
1:AAA:13:ASN:OD1	1:AAA:38:TYR:CE1	2.71	0.44
1:BBB:56:ASN:HD22	1:BBB:89:THR:HA	1.82	0.43
1:AAA:64:TYR:OH	1:AAA:81:LYS:HD2	2.18	0.43
1:BBB:51:ILE:O	1:BBB:93:GLN:N	2.49	0.42
1:AAA:42:GLU:HG2	4:AAA:337:HOH:O	2.19	0.42
1:AAA:51:ILE:O	1:AAA:93:GLN:N	2.43	0.42
1:BBB:47:ILE:CD1	1:BBB:59:THR:HG22	2.50	0.41
1:AAA:54:SER:HA	1:AAA:90:CYS:O	2.21	0.41
1:BBB:38:TYR:HB3	1:BBB:64:TYR:HB3	2.01	0.41
1:BBB:50:GLY:H	1:BBB:93:GLN:HE21	1.67	0.41
1:AAA:56:ASN:ND2	1:AAA:89:THR:OG1	2.50	0.41
1:AAA:19:CYS:HB3	1:AAA:77:TYR:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	103/105 (98%)	97 (94%)	6 (6%)	0	100	100
1	BBB	104/105 (99%)	99 (95%)	4 (4%)	1 (1%)	15	8
All	All	207/210 (99%)	196 (95%)	10 (5%)	1 (0%)	29	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	BBB	15	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	98/98 (100%)	93 (95%)	5 (5%)	24	19
1	BBB	99/98 (101%)	97 (98%)	2 (2%)	55	57
All	All	197/196 (100%)	190 (96%)	7 (4%)	35	32

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	12	THR
1	AAA	61	SER
1	AAA	75	CYS
1	AAA	78	LYS
1	AAA	103	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	BBB	15	ARG
1	BBB	67	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	BBB	1	1	7,8,9	0.50	0	9,10,12	0.80	1 (11%)
1	PCA	AAA	1	1	7,8,9	0.50	0	9,10,12	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	BBB	1	1	-	0/0/11/13	0/1/1/1
1	PCA	AAA	1	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	1	PCA	O-C-CA	-2.17	119.09	124.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	AAA	201	-	1,3,3	2.67	1 (100%)	0,3,3	-	-
3	SO4	AAA	202	-	4,4,4	0.35	0	6,6,6	0.08	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	201	ACT	CH3-C	2.67	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	201	ACT	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.