



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 01:50 am BST

PDB ID : 6OUN
Title : Structure of HIV-1 Reverse Transcriptase (RT) in complex with dsDNA and (-)3TC-TP
Authors : Bertoletti, N.; Anderson, K.S.
Deposited on : 2019-05-04
Resolution : 2.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	558	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 87%, grey 11%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 87% 11% . </div> </div>
2	B	452	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 2%, yellow 9%, green 79%, grey 12%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 2% 79% 9% 12% </div> </div>
3	P	21	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 38%, yellow 48%, grey 14%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 38% 48% 14% </div> </div>
4	T	27	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 41%, yellow 41%, grey 19%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 41% 41% 19% </div> </div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 8228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reverse transcriptase/ribonuclease H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	546	Total	C	N	O	S	0	0	0
			4234	2736	707	783	8			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	172	ARG	LYS	conflict	UNP P03366
A	258	CYS	GLN	engineered mutation	UNP P03366
A	280	SER	CYS	engineered mutation	UNP P03366
A	461	ARG	LYS	conflict	UNP P03366
A	468	THR	PRO	conflict	UNP P03366
A	471	ASP	ASN	conflict	UNP P03366
A	512	GLN	LYS	conflict	UNP P03366

- Molecule 2 is a protein called P51 RT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	397	Total	C	N	O	S	0	0	0
			3116	2034	508	569	5			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-11	MET	-	expression tag	UNP P03366
B	-10	GLY	-	expression tag	UNP P03366
B	-9	SER	-	expression tag	UNP P03366
B	-8	SER	-	expression tag	UNP P03366
B	-7	HIS	-	expression tag	UNP P03366
B	-6	HIS	-	expression tag	UNP P03366
B	-5	HIS	-	expression tag	UNP P03366
B	-4	HIS	-	expression tag	UNP P03366
B	-3	HIS	-	expression tag	UNP P03366

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	HIS	-	expression tag	UNP P03366
B	-1	SER	-	expression tag	UNP P03366
B	0	SER	-	expression tag	UNP P03366
B	172	ARG	LYS	conflict	UNP P03366
B	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is a DNA chain called DNA primer 20-mer.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	P	18	Total	C	N	O	P	S	0	0	0
			366	175	64	109	17	1			


- Molecule 4 is a DNA chain called DNA template 27-mer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	T	22	Total	C	N	O	P	0	0	0
			457	213	93	129	22			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mg	0	0
			2	2		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O S 5 4 1	0	0
6	P	1	Total O S 5 4 1	0	0
6	T	1	Total O S 5 4 1	0	0

- Molecule 7 is Lamivudine Triphosphate (three-letter code: 1RZ) (formula: $C_8H_{14}N_3O_{12}P_3S$) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C N O P S 27 8 3 12 3 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	2	Total O 2 2	0	0
8	B	7	Total O 7 7	0	0
8	T	2	Total O 2 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Reverse transcriptase/ribonuclease H



- Molecule 2: P51 RT



- Molecule 3: DNA primer 20-mer



- Molecule 4: DNA template 27-mer



DA	DT	DG	G704	G705	G706	G707	G708	G709	G710	G711	G712	G713	G714	A715	A716	G717	A718	G719	G720	G721	G722	G723	G724	G725	DT	DG
----	----	----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	----	----

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	167.01Å 171.78Å 105.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.57 – 2.66 29.57 – 2.66	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.57-2.66) 99.9 (29.57-2.66)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 2.64Å)	Xtriage
Refinement program	PHENIX (1.14 _3260: ???)	Depositor
R, R_{free}	0.207 , 0.259 0.209 , 0.261	Depositor DCC
R_{free} test set	2208 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	76.1	Xtriage
Anisotropy	0.542	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.006 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8228	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 1RZ, DDG, SO4, G47

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/4346	0.51	0/5947
2	B	0.32	0/3205	0.48	0/4386
3	P	0.85	0/355	0.97	0/543
4	T	0.62	0/514	0.79	0/792
All	All	0.39	0/8420	0.55	0/11668

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	183	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4234	0	4041	36	0
2	B	3116	0	2993	21	0
3	P	366	0	207	10	0
4	T	457	0	244	9	0
5	A	2	0	0	0	0
6	A	5	0	0	0	0
6	P	5	0	0	0	0
6	T	5	0	0	0	0
7	A	27	0	13	1	0
8	A	2	0	0	0	0
8	B	7	0	0	0	0
8	T	2	0	0	0	0
All	All	8228	0	7498	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:P:816:DG:H2'	3:P:817:G47:H8	1.55	0.87
1:A:492:GLU:HG2	1:A:530:LYS:HB2	1.74	0.69
1:A:277:ARG:NH2	1:A:355:ALA:O	2.30	0.65
4:T:713:DC:H2'	4:T:714:DG:C8	2.32	0.65
3:P:820:DC:C2'	3:P:821:DC:H5'	2.29	0.63
1:A:60:VAL:HG12	1:A:75:VAL:HG22	1.81	0.62
2:B:73:LYS:NZ	2:B:146:TYR:OH	2.33	0.61
2:B:163:SER:O	2:B:167:ILE:HG12	2.01	0.61
2:B:84:THR:HG21	2:B:153:TRP:HE1	1.64	0.61
3:P:816:DG:H2'	3:P:817:G47:C8	2.29	0.60
1:A:339:TYR:CZ	1:A:352:GLY:HA3	2.39	0.58
1:A:474:ASN:O	1:A:477:THR:HG22	2.03	0.57
3:P:806:DT:H2'	3:P:807:DC:C6	2.38	0.57
2:B:54:ASN:HB3	2:B:143:ARG:HH21	1.70	0.57
2:B:38:CYS:SG	2:B:132:ILE:HD11	2.45	0.57
1:A:181:TYR:HB2	1:A:188:TYR:HB3	1.87	0.56
1:A:114:ALA:N	7:A:604:1RZ:O2A	2.38	0.55
3:P:820:DC:H2''	3:P:821:DC:H5'	1.87	0.55
1:A:475:GLN:HB3	1:A:501:TYR:CE2	2.41	0.55
2:B:115:TYR:HB3	2:B:149:LEU:HB2	1.88	0.54
2:B:249:LYS:HB2	2:B:252:TRP:CE2	2.42	0.54
1:A:444:GLY:HA3	1:A:477:THR:HG23	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:ARG:HD3	1:A:74:LEU:HD21	1.91	0.51
1:A:332:GLN:HB2	1:A:336:GLN:HB2	1.91	0.51
1:A:460:ASN:OD1	1:A:461:ARG:HG3	2.11	0.51
1:A:273:GLY:H	1:A:338:THR:HG21	1.76	0.51
1:A:458:VAL:HG22	1:A:464:GLN:HG2	1.93	0.51
4:T:710:DG:H2'	4:T:711:DC:C6	2.46	0.50
1:A:114:ALA:HB1	1:A:160:PHE:CZ	2.46	0.50
1:A:17:ASP:O	1:A:83:ARG:HD3	2.12	0.49
1:A:460:ASN:HD22	2:B:288:ALA:HA	1.77	0.49
2:B:157:PRO:HG3	2:B:184:MET:HA	1.95	0.48
4:T:711:DC:H2'	4:T:712:DC:C6	2.47	0.48
1:A:281:LYS:O	1:A:284:ARG:HG2	2.14	0.48
4:T:714:DG:H2''	4:T:715:DA:C8	2.50	0.47
2:B:61:PHE:CZ	2:B:74:LEU:HD23	2.50	0.47
1:A:23:GLN:HE22	1:A:60:VAL:H	1.62	0.46
1:A:390:LYS:HB3	1:A:417:VAL:HG21	1.98	0.46
3:P:819:DG:H2'	3:P:820:DC:C6	2.50	0.46
1:A:172:ARG:NH2	2:B:139:THR:HG22	2.31	0.46
1:A:260:LEU:HD21	1:A:303:LEU:HD13	1.97	0.46
2:B:260:LEU:HD21	2:B:303:LEU:HD13	1.98	0.45
1:A:399:GLU:HA	1:A:402:TRP:HD1	1.80	0.45
2:B:19:PRO:HB2	2:B:58:THR:HG22	1.98	0.45
2:B:121:ASP:O	2:B:125:ARG:HG3	2.17	0.45
4:T:717:DC:H2''	4:T:718:DA:C8	2.51	0.45
1:A:494:ASN:HB3	2:B:289:LEU:HD12	1.98	0.45
4:T:712:DC:H2''	4:T:713:DC:C6	2.52	0.44
1:A:434:ILE:HD13	1:A:530:LYS:HB3	1.99	0.44
3:P:805:DG:H3'	3:P:806:DT:H73	1.99	0.44
1:A:175:ASN:HB3	1:A:178:ILE:HD12	1.99	0.44
1:A:405:TYR:CE2	1:A:407:GLN:HB2	2.54	0.43
1:A:259:LYS:HA	3:P:819:DG:H5'	1.99	0.43
2:B:175:ASN:HB3	2:B:178:ILE:HG13	2.00	0.43
1:A:160:PHE:CE1	1:A:184:MET:O	2.72	0.42
4:T:712:DC:H2''	4:T:713:DC:H6	1.84	0.42
3:P:814:DC:H2''	3:P:815:DG:C8	2.55	0.42
1:A:89:GLU:HG3	4:T:708:DG:OP1	2.19	0.42
4:T:706:DC:H2'	4:T:707:DG:C8	2.54	0.42
2:B:105:SER:O	2:B:190:GLY:HA2	2.19	0.42
2:B:21:VAL:HB	2:B:59:PRO:HD3	2.02	0.42
2:B:342:TYR:HB3	2:B:348:ASN:HA	2.01	0.42
1:A:491:LEU:HD23	1:A:491:LEU:HA	1.87	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:HIS:CG	1:A:97:PRO:HD2	2.55	0.41
1:A:222:GLN:HE22	1:A:228:LEU:HB2	1.85	0.41
1:A:350:LYS:HB2	1:A:350:LYS:HE3	1.86	0.41
2:B:331:LYS:HB2	2:B:337:TRP:CZ3	2.56	0.41
1:A:370:GLU:HG2	2:B:394:GLN:HE22	1.86	0.40
3:P:819:DG:H2'	3:P:820:DC:H6	1.86	0.40
1:A:330:GLN:NE2	1:A:340:GLN:OE1	2.43	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	542/558 (97%)	526 (97%)	16 (3%)	0	100	100
2	B	391/452 (86%)	379 (97%)	12 (3%)	0	100	100
All	All	933/1010 (92%)	905 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	428/498 (86%)	420 (98%)	8 (2%)	57	74

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	320/411 (78%)	316 (99%)	4 (1%)	69	82
All	All	748/909 (82%)	736 (98%)	12 (2%)	62	78

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	THR
1	A	241	VAL
1	A	253	THR
1	A	314	VAL
1	A	373	GLN
1	A	402	TRP
1	A	477	THR
1	A	507	GLN
2	B	29	GLU
2	B	163	SER
2	B	234	LEU
2	B	396	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	221	HIS
2	B	207	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	G47	P	817	1,3,4	20,27,28	1.12	2 (10%)	21,38,41	2.18	6 (28%)
3	DDG	P	822	3,4	17,23,24	1.80	4 (23%)	15,33,36	2.69	10 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	G47	P	817	1,3,4	-	2/7/25/26	0/3/3/3
3	DDG	P	822	3,4	-	2/3/18/19	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	817	G47	C6-C5	3.96	1.48	1.41
3	P	822	DDG	C2-N1	-3.50	1.29	1.35
3	P	822	DDG	O4'-C4'	-3.06	1.38	1.44
3	P	822	DDG	C4-N3	-2.79	1.31	1.35
3	P	822	DDG	C6-N1	-2.45	1.28	1.33
3	P	817	G47	C5-C4	2.29	1.47	1.40

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	817	G47	C2-N3-C4	5.68	121.73	115.28
3	P	822	DDG	C2-N3-C4	4.73	120.76	115.36
3	P	822	DDG	C6-C5-C4	-4.51	116.50	120.80
3	P	817	G47	C6-C5-C4	-4.00	116.98	120.80
3	P	817	G47	C6-N1-C2	3.91	122.19	115.18
3	P	817	G47	C5-C6-N1	-3.57	118.55	123.43
3	P	822	DDG	O4'-C1'-C2'	-3.40	102.98	106.67
3	P	822	DDG	C6-N1-C2	3.17	120.97	115.93
3	P	822	DDG	C4'-O4'-C1'	3.01	112.65	109.81
3	P	822	DDG	C5-C6-N1	-2.88	119.49	123.43
3	P	822	DDG	N3-C2-N1	-2.60	123.75	127.22
3	P	822	DDG	C3'-C2'-C1'	2.54	105.72	102.78
3	P	817	G47	C4-C5-N7	-2.50	106.80	109.40
3	P	822	DDG	C4-C5-N7	-2.45	106.84	109.40
3	P	817	G47	N3-C2-N1	-2.39	122.45	126.23
3	P	822	DDG	O4'-C4'-C3'	-2.31	100.98	104.80

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	P	817	G47	C3'-C4'-C5'-O5'
3	P	822	DDG	O4'-C4'-C5'-O5'
3	P	817	G47	O4'-C4'-C5'-O5'
3	P	822	DDG	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	P	817	G47	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	P	901	-	4,4,4	0.13	0	6,6,6	0.08	0
6	SO4	T	801	-	4,4,4	0.14	0	6,6,6	0.07	0
7	1RZ	A	604	5	22,28,28	0.67	0	27,43,43	1.14	2 (7%)
6	SO4	A	603	-	4,4,4	0.15	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	1RZ	A	604	5	-	3/19/31/31	0/2/2/2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	604	1RZ	C5'-C4'-S3'	-2.63	106.78	113.67
7	A	604	1RZ	PB-O3B-PG	-2.06	125.77	132.83

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	A	604	1RZ	PB-O3A-PA-O5'
7	A	604	1RZ	PA-O3A-PB-O2B
7	A	604	1RZ	C4'-C5'-O5'-PA

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	604	1RZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	546/558 (97%)	-0.22	6 (1%) 80 79	48, 79, 125, 164	0
2	B	397/452 (87%)	-0.10	11 (2%) 53 49	54, 83, 142, 183	0
3	P	16/21 (76%)	-0.53	0 100 100	73, 89, 105, 111	0
4	T	22/27 (81%)	-0.37	0 100 100	67, 97, 126, 144	0
All	All	981/1058 (92%)	-0.18	17 (1%) 70 67	48, 81, 135, 183	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	87	PHE	6.3
2	B	210	LEU	4.2
1	A	68	SER	3.3
2	B	360	ALA	3.2
2	B	334	GLN	2.6
2	B	359	GLY	2.5
1	A	156	SER	2.4
2	B	357	MET	2.4
1	A	69	THR	2.4
2	B	245	VAL	2.4
2	B	301	LEU	2.3
2	B	212	TRP	2.2
2	B	69	THR	2.2
1	A	290	THR	2.2
2	B	67	ASP	2.1
1	A	25	PRO	2.1
1	A	284	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	G47	P	817	25/26	0.94	0.13	79,95,113,118	0
3	DDG	P	822	21/22	0.98	0.11	46,56,66,73	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	A	602	1/1	0.81	0.05	78,78,78,78	0
7	1RZ	A	604	27/27	0.91	0.18	75,105,151,152	0
6	SO4	P	901	5/5	0.91	0.16	158,158,159,160	0
6	SO4	A	603	5/5	0.95	0.16	119,120,125,126	0
6	SO4	T	801	5/5	0.95	0.10	119,120,123,125	0
5	MG	A	601	1/1	0.99	0.16	57,57,57,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 1RZ A 604:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.