



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 02:58 pm BST

PDB ID : 6OUX
Title : Structure of SMUL_1544, a decarboxylase from *Sulfurospirillum multivorans*
Authors : Wetterhorn, K.M.; Rayment, I.; Vecellio, A.; Seeger, M.; Keller, S.; Schubert, T.
Deposited on : 2019-05-05
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>16%</div> <div>• •</div> </div> </div>
1	B	393	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>11%</div> <div>• 10%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6528 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Threonine phosphate decarboxylase-like enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	393	Total	C	N	O	S	0	0	0
			3165	2036	522	585	22			
1	B	354	Total	C	N	O	S	0	0	0
			2864	1853	466	525	20			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ALA	-	expression tag	UNP W6EE14
A	1	SER	-	expression tag	UNP W6EE14
B	0	ALA	-	expression tag	UNP W6EE14
B	1	SER	-	expression tag	UNP W6EE14


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	245	Total	O	0	0
			245	245		
2	B	254	Total	O	0	0
			254	254		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Threonine phosphate decarboxylase-like enzyme



- Molecule 1: Threonine phosphate decarboxylase-like enzyme



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	141.53Å 90.63Å 70.87Å 90.00° 109.96° 90.00°	Depositor
Resolution (Å)	74.90 – 1.94 33.31 – 1.94	Depositor EDS
% Data completeness (in resolution range)	94.3 (74.90-1.94) 94.3 (33.31-1.94)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.190 , 0.250 0.198 , 0.255	Depositor DCC
R_{free} test set	2932 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6528	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	2/3228 (0.1%)	1.04	14/4357 (0.3%)
1	B	0.89	0/2920	0.98	9/3941 (0.2%)
All	All	0.91	2/6148 (0.0%)	1.02	23/8298 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	180	ASN	C-O	8.54	1.39	1.23
1	A	242	SER	CB-OG	5.30	1.49	1.42

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	ARG	NE-CZ-NH1	-16.82	111.89	120.30
1	A	64	ARG	NE-CZ-NH2	14.17	127.38	120.30
1	A	324	MET	CG-SD-CE	-11.71	81.46	100.20
1	B	257	MET	CG-SD-CE	-10.41	83.55	100.20
1	B	372	ARG	NE-CZ-NH1	8.88	124.74	120.30
1	A	180	ASN	C-N-CD	7.84	144.86	128.40
1	A	230	ASP	CB-CG-OD2	-7.18	111.84	118.30
1	B	372	ARG	NE-CZ-NH2	-6.69	116.96	120.30
1	B	293	ARG	NE-CZ-NH1	6.59	123.59	120.30
1	A	307	ARG	NE-CZ-NH2	6.31	123.45	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	180	ASN	N-CA-C	6.19	127.71	111.00
1	B	190	LYS	CD-CE-NZ	5.86	125.17	111.70
1	A	64	ARG	CD-NE-CZ	5.85	131.79	123.60
1	A	307	ARG	NE-CZ-NH1	-5.79	117.41	120.30
1	B	126	MET	CG-SD-CE	-5.70	91.08	100.20
1	A	45	ASP	CB-CG-OD1	5.60	123.34	118.30
1	B	257	MET	CA-CB-CG	5.51	122.67	113.30
1	A	180	ASN	N-CA-CB	-5.35	100.97	110.60
1	A	180	ASN	CB-CA-C	-5.23	99.95	110.40
1	A	329	MET	CG-SD-CE	5.19	108.50	100.20
1	B	160	LYS	CD-CE-NZ	5.17	123.59	111.70
1	B	305	MET	CG-SD-CE	-5.09	92.05	100.20
1	A	230	ASP	CB-CG-OD1	5.02	122.82	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	180	ASN	Peptide
1	A	390	ASP	Peptide
1	B	272	TRP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3165	0	3193	71	0
1	B	2864	0	2883	44	0
2	A	245	0	0	22	0
2	B	254	0	0	24	0
All	All	6528	0	6076	112	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (112) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:83:GLN:NE2	1:B:273:ASN:HB3	1.42	1.33
1:A:329:MET:HE3	1:A:356:MET:SD	1.78	1.22
1:A:329:MET:CE	1:A:356:MET:SD	2.34	1.16
1:A:329:MET:SD	2:A:638:HOH:O	2.04	1.15
1:A:30:MET:O	2:A:401:HOH:O	1.66	1.14
1:B:16:PHE:N	2:B:402:HOH:O	1.78	1.13
1:B:83:GLN:NE2	1:B:273:ASN:CB	2.12	1.10
1:A:58:VAL:HG11	1:A:247:ILE:HD11	1.37	1.07
1:A:246:GLY:O	1:A:247:ILE:HG13	1.61	1.00
1:B:68:GLN:NE2	2:B:403:HOH:O	2.01	0.91
1:A:0:ALA:CB	2:A:626:HOH:O	2.18	0.89
1:A:56:GLU:HG2	2:A:598:HOH:O	1.72	0.89
1:B:50:SER:HB3	2:B:417:HOH:O	1.72	0.88
1:A:2:VAL:HB	2:A:404:HOH:O	1.74	0.88
1:B:83:GLN:CD	1:B:273:ASN:HB3	1.93	0.87
1:A:129:THR:CG2	1:A:180:ASN:HD21	1.89	0.85
1:A:0:ALA:HB1	2:A:626:HOH:O	1.76	0.85
1:A:247:ILE:HG22	1:A:250:VAL:HG13	1.60	0.84
1:A:58:VAL:HG11	1:A:247:ILE:CD1	2.07	0.84
1:A:83:GLN:HE22	1:A:103:ILE:H	1.26	0.82
1:A:180:ASN:OD1	1:A:181:PRO:HD3	1.80	0.80
1:B:48:TYR:O	1:B:372:ARG:NH2	2.14	0.80
1:A:247:ILE:CG2	1:A:250:VAL:HG13	2.12	0.79
1:A:247:ILE:HD12	1:A:285:PHE:HZ	1.47	0.79
1:B:14:LYS:NZ	2:B:404:HOH:O	2.14	0.79
1:B:13:THR:C	2:B:402:HOH:O	2.21	0.79
1:A:180:ASN:OD1	1:A:181:PRO:CD	2.31	0.79
1:A:329:MET:HE2	1:A:356:MET:SD	2.26	0.76
1:B:83:GLN:HE21	1:B:273:ASN:CB	1.95	0.75
1:B:242:SER:HB3	2:B:434:HOH:O	1.85	0.75
1:B:13:THR:O	2:B:402:HOH:O	2.06	0.73
1:A:317:ASN:O	1:A:333:PRO:HG3	1.89	0.73
1:A:0:ALA:HB2	2:A:626:HOH:O	1.87	0.72
1:A:390:ASP:OD1	2:A:403:HOH:O	2.09	0.71
1:B:119:GLU:HG3	2:B:416:HOH:O	1.90	0.70
1:A:129:THR:HG22	1:A:180:ASN:ND2	2.08	0.69
1:A:129:THR:HG23	1:A:180:ASN:HD21	1.58	0.67
1:B:14:LYS:C	2:B:402:HOH:O	2.32	0.66
1:A:33:LEU:N	2:A:401:HOH:O	2.28	0.66
1:A:186:LEU:HD23	1:A:191:MET:SD	2.35	0.66
1:A:105:ASN:OD1	1:A:272:TRP:CZ3	2.50	0.65
1:B:50:SER:CB	2:B:417:HOH:O	2.35	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:ASN:ND2	2:A:406:HOH:O	2.29	0.64
1:A:22:ASN:HD22	1:A:24:GLY:H	1.44	0.64
1:B:212:HIS:HD2	1:B:294:TYR:OH	1.82	0.63
1:A:129:THR:HG22	1:A:180:ASN:HD21	1.59	0.63
1:B:15:ALA:N	2:B:402:HOH:O	2.32	0.62
1:B:309:LYS:HE3	2:B:542:HOH:O	2.00	0.62
1:B:309:LYS:CE	2:B:542:HOH:O	2.47	0.62
1:A:73:LYS:O	1:A:77:THR:HB	2.01	0.60
1:A:56:GLU:CG	2:A:598:HOH:O	2.41	0.59
1:A:212:HIS:HD2	1:A:294:TYR:OH	1.85	0.59
1:A:329:MET:HE2	1:A:356:MET:CG	2.32	0.59
1:B:127:ILE:HD13	1:B:145:TYR:CD2	2.38	0.59
1:A:246:GLY:C	1:A:247:ILE:HG13	2.23	0.59
1:A:3:ASP:OD1	2:A:404:HOH:O	2.17	0.58
1:A:79:TYR:CD2	1:B:48:TYR:HB2	2.43	0.54
1:B:17:HIS:HB2	2:B:607:HOH:O	2.08	0.54
1:B:83:GLN:HE21	1:B:273:ASN:HB2	1.73	0.53
1:B:83:GLN:N	2:B:414:HOH:O	2.41	0.52
1:B:127:ILE:HD12	2:B:529:HOH:O	2.09	0.52
1:A:126:MET:SD	1:A:156:PHE:CE1	3.03	0.52
1:A:247:ILE:CD1	1:A:285:PHE:HZ	2.19	0.51
1:A:58:VAL:CG1	1:A:247:ILE:HD11	2.25	0.51
1:B:90:MET:HG2	1:B:286:VAL:HG11	1.92	0.51
1:B:83:GLN:HE22	1:B:273:ASN:ND2	2.08	0.51
1:B:103:ILE:CD1	2:B:633:HOH:O	2.58	0.51
1:A:247:ILE:HD12	1:A:285:PHE:CZ	2.36	0.51
1:B:103:ILE:HD11	2:B:633:HOH:O	2.11	0.51
1:B:83:GLN:NE2	1:B:273:ASN:ND2	2.60	0.50
1:A:52:PRO:O	1:A:56:GLU:HG3	2.13	0.49
1:A:188:LEU:HD21	2:A:538:HOH:O	2.13	0.48
1:A:227:VAL:HG13	2:A:563:HOH:O	2.12	0.48
1:B:83:GLN:NE2	1:B:273:ASN:CG	2.65	0.48
1:A:129:THR:CG2	1:A:180:ASN:ND2	2.63	0.48
1:A:8:ARG:NH1	1:A:348:GLU:OE2	2.47	0.47
1:A:31:GLU:C	2:A:401:HOH:O	2.52	0.47
1:A:378:ASN:ND2	2:A:420:HOH:O	2.47	0.47
1:B:64:ARG:NH1	2:B:421:HOH:O	2.47	0.47
1:A:329:MET:CE	1:A:356:MET:CG	2.93	0.47
1:A:67:ILE:HG13	1:B:72:TYR:CZ	2.51	0.46
1:A:34:LYS:N	2:A:401:HOH:O	1.99	0.45
1:A:30:MET:N	2:A:421:HOH:O	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16:PHE:HB3	2:B:402:HOH:O	2.17	0.45
1:B:343:SER:HB3	2:B:569:HOH:O	2.16	0.44
1:A:36:MET:HA	2:A:625:HOH:O	2.16	0.44
1:A:391:SER:OG	1:A:392:LYS:N	2.48	0.44
1:A:212:HIS:HB2	2:A:464:HOH:O	2.17	0.44
1:A:271:LEU:O	1:A:271:LEU:HD12	2.18	0.44
1:A:79:TYR:CE2	1:B:48:TYR:HB2	2.53	0.44
1:B:125:LEU:CD2	1:B:174:VAL:HB	2.48	0.44
1:A:161:TYR:OH	1:A:173:VAL:HG13	2.18	0.43
1:B:127:ILE:HG22	1:B:147:PRO:HA	1.99	0.43
1:A:127:ILE:CG2	1:A:147:PRO:HA	2.48	0.43
1:A:308:PHE:CD1	1:A:377:ASN:HB3	2.53	0.43
1:A:26:HIS:HB3	2:A:421:HOH:O	2.18	0.43
1:B:62:ILE:HG22	1:B:67:ILE:HD13	2.01	0.42
1:A:268:HIS:H	1:A:269:GLY:HA3	1.85	0.42
1:A:105:ASN:OD1	1:A:272:TRP:HZ3	2.01	0.41
1:A:237:ILE:O	1:A:255:ALA:HA	2.19	0.41
1:A:58:VAL:HG21	1:A:247:ILE:HD11	2.02	0.41
1:A:247:ILE:O	1:A:247:ILE:HG22	2.21	0.41
1:B:292:LYS:HE3	2:B:611:HOH:O	2.21	0.41
1:A:305:MET:SD	1:A:371:GLY:HA3	2.61	0.41
1:B:161:TYR:O	1:B:164:PHE:HB3	2.20	0.41
1:A:215:TYR:H	1:A:324:MET:HE1	1.86	0.41
1:B:83:GLN:NE2	1:B:273:ASN:HD22	2.19	0.40
1:A:180:ASN:OD1	1:A:181:PRO:N	2.54	0.40
1:A:181:PRO:HB2	1:A:329:MET:HE3	2.03	0.40
1:B:9:ASN:ND2	2:B:410:HOH:O	2.37	0.40
1:A:28:PRO:HB2	2:A:456:HOH:O	2.22	0.40
1:B:16:PHE:CB	2:B:402:HOH:O	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	391/393 (100%)	377 (96%)	8 (2%)	6 (2%)	10	2
1	B	348/393 (88%)	340 (98%)	7 (2%)	1 (0%)	41	32
All	All	739/786 (94%)	717 (97%)	15 (2%)	7 (1%)	17	7

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	247	ILE
1	A	391	SER
1	A	246	GLY
1	A	268	HIS
1	A	181	PRO
1	B	70	ASN
1	A	269	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/349 (100%)	334 (96%)	15 (4%)	29	14
1	B	314/349 (90%)	305 (97%)	9 (3%)	42	28
All	All	663/698 (95%)	639 (96%)	24 (4%)	35	20

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	30	MET
1	A	77	THR
1	A	85	SER
1	A	132	SER
1	A	148	LEU
1	A	181	PRO
1	A	190	LYS
1	A	212	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	260	ARG
1	A	268	HIS
1	A	334	SER
1	A	352	TYR
1	A	375	GLU
1	A	390	ASP
1	B	37	PHE
1	B	39	THR
1	B	69	THR
1	B	84	ARG
1	B	88	LYS
1	B	119	GLU
1	B	129	THR
1	B	194	LEU
1	B	212	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	ASN
1	A	70	ASN
1	A	78	HIS
1	A	83	GLN
1	A	117	GLN
1	A	121	GLN
1	A	212	HIS
1	B	83	GLN
1	B	167	ASN
1	B	169	GLN
1	B	212	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	393/393 (100%)	-0.10	16 (4%) 37 44	9, 21, 48, 95	0
1	B	354/393 (90%)	-0.15	13 (3%) 41 49	9, 21, 46, 63	0
All	All	747/786 (95%)	-0.12	29 (3%) 39 47	9, 21, 47, 95	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	270	PHE	7.5
1	A	247	ILE	4.9
1	B	37	PHE	4.5
1	B	40	LEU	4.2
1	A	271	LEU	3.9
1	B	83	GLN	3.7
1	A	35	LYS	3.6
1	B	85	SER	3.5
1	B	18	ALA	3.4
1	A	248	ALA	3.3
1	A	272	TRP	3.2
1	A	246	GLY	3.2
1	B	41	GLU	3.1
1	A	180	ASN	3.1
1	B	38	PRO	2.9
1	A	245	PHE	2.7
1	A	392	LYS	2.6
1	B	314	GLY	2.6
1	A	390	ASP	2.5
1	B	69	THR	2.5
1	A	268	HIS	2.4
1	B	39	THR	2.4
1	A	267	GLU	2.3
1	A	39	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	36	MET	2.3
1	B	139	LYS	2.3
1	B	68	GLN	2.2
1	A	118	GLU	2.2
1	A	391	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.