



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 01:44 am BST

PDB ID : 2OVY  
Title : Crystal structure of the catalytic domain of rat phosphodiesterase 10A  
Authors : Pandit, J.; Marr, E.S.  
Deposited on : 2007-02-15  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	<div> <div>2%</div> <div>67%</div> <div>14%</div> <div>•</div> <div>16%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	997[B]	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphodiesterase-10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	305	2488	1586	433	447	22	0	1	0

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	423	MET	-	EXPRESSION TAG	UNP Q9QYJ6
A	424	HIS	-	EXPRESSION TAG	UNP Q9QYJ6
A	425	HIS	-	EXPRESSION TAG	UNP Q9QYJ6
A	426	HIS	-	EXPRESSION TAG	UNP Q9QYJ6
A	427	HIS	-	EXPRESSION TAG	UNP Q9QYJ6
A	428	HIS	-	EXPRESSION TAG	UNP Q9QYJ6
A	429	HIS	-	EXPRESSION TAG	UNP Q9QYJ6
A	430	SER	-	EXPRESSION TAG	UNP Q9QYJ6
A	431	SER	-	EXPRESSION TAG	UNP Q9QYJ6
A	432	GLY	-	EXPRESSION TAG	UNP Q9QYJ6
A	433	LEU	-	EXPRESSION TAG	UNP Q9QYJ6
A	434	VAL	-	EXPRESSION TAG	UNP Q9QYJ6
A	435	PRO	-	EXPRESSION TAG	UNP Q9QYJ6
A	436	ARG	-	EXPRESSION TAG	UNP Q9QYJ6
A	437	GLY	-	EXPRESSION TAG	UNP Q9QYJ6
A	438	SER	-	EXPRESSION TAG	UNP Q9QYJ6
A	439	ALA	-	EXPRESSION TAG	UNP Q9QYJ6
A	440	MET	-	EXPRESSION TAG	UNP Q9QYJ6
A	441	GLY	-	EXPRESSION TAG	UNP Q9QYJ6

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	1
			10	8	2		


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is 6,7-DIMETHOXY-4-[(3R)-3-(QUINOXALIN-2-YLOXY)PYRROLIDIN-1-YL]QUINAZOLINE (three-letter code: PFJ) (formula: C<sub>22</sub>H<sub>21</sub>N<sub>5</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			30	22	5	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	334	Total	O	0	0
			334	334		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Phosphodiesterase-10A



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.80 Å   120.80 Å   83.50 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	65.94 – 1.80 35.74 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (65.94-1.80) 99.7 (35.74-1.80)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.79 (at 1.81 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.157 , 0.198 0.156 , 0.196	Depositor DCC
$R_{free}$ test set	3228 reflections (7.69%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.8	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 50.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2864	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PFJ, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.70	27/2549 (1.1%)	1.30	21/3451 (0.6%)

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	720	TYR	CD1-CE1	9.69	1.53	1.39
1	A	610	GLU	CG-CD	9.45	1.66	1.51
1	A	629	PHE	CG-CD2	8.33	1.51	1.38
1	A	509	TYR	CD1-CE1	8.19	1.51	1.39
1	A	608	GLU	CD-OE2	7.81	1.34	1.25
1	A	582	GLU	CG-CD	7.56	1.63	1.51
1	A	685	GLU	CG-CD	6.75	1.62	1.51
1	A	618	LYS	CE-NZ	6.72	1.65	1.49
1	A	668	VAL	CB-CG2	6.62	1.66	1.52
1	A	528	TYR	CD2-CE2	6.57	1.49	1.39
1	A	463	GLU	CD-OE1	6.20	1.32	1.25
1	A	720	TYR	CD2-CE2	6.14	1.48	1.39
1	A	686	PHE	CD1-CE1	6.13	1.51	1.39
1	A	590	VAL	CB-CG1	5.99	1.65	1.52
1	A	551	LEU	CG-CD2	5.74	1.73	1.51
1	A	657	GLY	N-CA	5.74	1.54	1.46
1	A	692	GLU	CG-CD	5.74	1.60	1.51
1	A	683	TYR	CG-CD1	5.68	1.46	1.39
1	A	494	GLU	CB-CG	5.49	1.62	1.52
1	A	610	GLU	CD-OE1	5.44	1.31	1.25
1	A	596	GLU	CD-OE1	5.35	1.31	1.25
1	A	685	GLU	CB-CG	-5.33	1.42	1.52
1	A	482	TYR	CD2-CE2	5.24	1.47	1.39
1	A	629	PHE	CE2-CZ	5.18	1.47	1.37
1	A	655	VAL	CB-CG2	5.11	1.63	1.52
1	A	517	TRP	CE3-CZ3	5.10	1.47	1.38

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	586	PHE	CD1-CE1	5.03	1.49	1.39

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	746	ARG	NE-CZ-NH2	-9.51	115.54	120.30
1	A	652	ARG	NE-CZ-NH2	-9.38	115.61	120.30
1	A	719	PHE	CB-CG-CD1	-8.88	114.59	120.80
1	A	681	ASP	CB-CG-OD2	-8.30	110.83	118.30
1	A	653	ASP	CB-CG-OD2	-8.25	110.88	118.30
1	A	629	PHE	CB-CG-CD1	-7.32	115.67	120.80
1	A	565	LEU	CB-CG-CD2	-6.98	99.13	111.00
1	A	510	ARG	NE-CZ-NH1	-6.92	116.84	120.30
1	A	632	ARG	NE-CZ-NH2	-6.67	116.97	120.30
1	A	691	ASP	CB-CG-OD2	-6.61	112.36	118.30
1	A	661	THR	CA-CB-CG2	-6.28	103.60	112.40
1	A	567	LYS	CD-CE-NZ	-6.13	97.60	111.70
1	A	613	LEU	CB-CG-CD1	-6.06	100.70	111.00
1	A	691	ASP	CB-CG-OD1	5.69	123.42	118.30
1	A	632	ARG	NE-CZ-NH1	-5.61	117.49	120.30
1	A	632	ARG	NH1-CZ-NH2	5.51	125.46	119.40
1	A	528	TYR	CZ-CE2-CD2	-5.46	114.89	119.80
1	A	704	MET	CG-SD-CE	-5.44	91.49	100.20
1	A	653	ASP	OD1-CG-OD2	5.35	133.47	123.30
1	A	658	LEU	CB-CG-CD2	-5.28	102.03	111.00
1	A	609	TYR	CB-CG-CD1	-5.22	117.87	121.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2488	0	2458	39	0
2	A	10	0	0	3	0
3	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
5	A	30	0	21	3	0
6	A	334	0	0	14	1
All	All	2864	0	2479	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:492:CYS:SG	1:A:544:LYS:HE2	1.53	1.46
1:A:492:CYS:SG	1:A:544:LYS:CE	2.23	1.25
1:A:710:ASP:HB2	6:A:317:HOH:O	1.39	1.20
1:A:459:CYS:HB3	6:A:322:HOH:O	1.01	1.18
1:A:558[A]:ARG:HH11	1:A:558[A]:ARG:HG2	1.13	1.05
2:A:997[B]:SO4:S	6:A:250:HOH:O	2.21	0.97
1:A:492:CYS:SG	1:A:544:LYS:CD	2.54	0.95
2:A:997[B]:SO4:O3	6:A:250:HOH:O	1.98	0.79
5:A:999:PFJ:H3	5:A:999:PFJ:C16	2.13	0.79
1:A:485:HIS:CD2	6:A:266:HOH:O	2.36	0.77
1:A:639:TYR:HB2	6:A:211:HOH:O	1.83	0.77
1:A:558[A]:ARG:CG	1:A:558[A]:ARG:HH11	1.97	0.76
5:A:999:PFJ:H3	5:A:999:PFJ:H162	1.67	0.76
1:A:497:LYS:NZ	6:A:280:HOH:O	2.20	0.73
1:A:558[A]:ARG:NH1	1:A:558[A]:ARG:HG2	1.92	0.73
2:A:997[B]:SO4:O2	6:A:250:HOH:O	2.08	0.70
1:A:492:CYS:SG	1:A:544:LYS:HD3	2.35	0.67
1:A:488:CYS:SG	1:A:544:LYS:HG3	2.36	0.65
1:A:491:SER:O	1:A:492:CYS:C	2.35	0.64
1:A:492:CYS:SG	1:A:544:LYS:CG	2.93	0.56
1:A:488:CYS:SG	1:A:544:LYS:CG	2.95	0.55
1:A:709:ARG:NH1	1:A:709:ARG:HG2	2.21	0.54
1:A:540:ASP:OD1	1:A:543:ARG:NH2	2.42	0.53
1:A:492:CYS:SG	1:A:544:LYS:HG2	2.47	0.53
1:A:535:ASN:HB2	6:A:254:HOH:O	2.08	0.53
1:A:558[A]:ARG:CG	1:A:558[A]:ARG:NH1	2.62	0.53
5:A:999:PFJ:C3	5:A:999:PFJ:C16	2.86	0.52
1:A:499:CYS:HB3	6:A:220:HOH:O	2.09	0.52
1:A:511:ARG:HG2	1:A:511:ARG:HH11	1.75	0.51
1:A:482:TYR:CZ	1:A:486:ARG:HD2	2.46	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:488:CYS:HA	6:A:252:HOH:O	2.09	0.51
1:A:488:CYS:HB3	1:A:544:LYS:HE2	1.95	0.49
1:A:488:CYS:HB3	1:A:544:LYS:CE	2.43	0.48
1:A:534:ASN:HB3	1:A:537:LEU:HD12	1.95	0.47
1:A:511:ARG:CG	1:A:511:ARG:HH11	2.27	0.47
1:A:510:ARG:NE	1:A:558[B]:ARG:HD2	2.30	0.46
1:A:709:ARG:HH11	1:A:709:ARG:HG2	1.81	0.45
1:A:459:CYS:CB	6:A:322:HOH:O	1.91	0.44
1:A:614:GLU:HG3	1:A:618:LYS:HE3	1.99	0.44
1:A:690:GLY:HA3	1:A:704:MET:O	2.18	0.43
1:A:488:CYS:CA	6:A:252:HOH:O	2.68	0.41
1:A:511:ARG:NH1	1:A:511:ARG:CG	2.83	0.41
1:A:454:LEU:HB2	1:A:459:CYS:SG	2.61	0.41
1:A:482:TYR:OH	1:A:486:ARG:HD2	2.20	0.41
1:A:558[A]:ARG:NH2	6:A:277:HOH:O	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:318:HOH:O	6:A:321:HOH:O 3_555	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	304/362 (84%)	299 (98%)	5 (2%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	275/322 (85%)	267 (97%)	8 (3%)	42 29

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	474	ASN
1	A	494	GLU
1	A	497	LYS
1	A	535	ASN
1	A	558[A]	ARG
1	A	558[B]	ARG
1	A	632	ARG
1	A	709	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	474	ASN
1	A	748	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	997[B]	-	4,4,4	1.01	0	6,6,6	2.05	2 (33%)
2	SO4	A	997[A]	-	4,4,4	0.32	0	6,6,6	0.66	0
5	PFJ	A	999	-	34,34,34	2.08	9 (26%)	46,48,48	3.69	22 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PFJ	A	999	-	-	2/12/21/21	0/5/5/5

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	999	PFJ	C21-N24	6.28	1.38	1.30
5	A	999	PFJ	C3-C2	4.42	1.44	1.36
5	A	999	PFJ	C6-C5	-4.28	1.35	1.42
5	A	999	PFJ	O13-C2	3.80	1.43	1.37
5	A	999	PFJ	C7-N8	3.22	1.37	1.33
5	A	999	PFJ	C4-C1	2.47	1.41	1.36
5	A	999	PFJ	C22-N23	2.30	1.35	1.31
5	A	999	PFJ	C7-C6	-2.14	1.39	1.43
5	A	999	PFJ	C9-N10	2.09	1.35	1.32

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	999	PFJ	C21-N24-C25	11.20	123.98	115.79
5	A	999	PFJ	C22-C21-N24	-9.76	119.16	124.09

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	999	PFJ	C21-C22-N23	-9.72	115.98	121.86
5	A	999	PFJ	C22-N23-C26	6.96	125.29	116.91
5	A	999	PFJ	C21-O20-C17	6.53	124.44	117.75
5	A	999	PFJ	C14-O13-C2	6.01	126.60	117.53
5	A	999	PFJ	N10-C9-N8	-4.89	121.04	128.68
5	A	999	PFJ	O20-C17-C16	4.09	118.13	108.19
5	A	999	PFJ	C3-C6-C7	-4.00	120.74	124.94
2	A	997[B]	SO4	O4-S-O2	3.47	127.42	109.31
5	A	999	PFJ	C25-C26-N23	-3.22	117.92	121.00
2	A	997[B]	SO4	O3-S-O2	-3.11	93.08	109.31
5	A	999	PFJ	C30-C25-C26	3.05	123.06	119.25
5	A	999	PFJ	C26-C25-N24	-3.03	117.24	121.18
5	A	999	PFJ	O13-C2-C3	3.02	129.14	125.24
5	A	999	PFJ	C12-O11-C1	2.81	121.78	117.53
5	A	999	PFJ	O11-C1-C4	-2.73	121.70	125.24
5	A	999	PFJ	C27-C26-N23	2.67	122.90	118.52
5	A	999	PFJ	C19-N15-C7	-2.64	115.73	123.56
5	A	999	PFJ	C16-N15-C7	2.56	131.02	123.61
5	A	999	PFJ	O20-C21-C22	2.37	122.12	117.54
5	A	999	PFJ	C9-N10-C5	2.35	118.64	115.40
5	A	999	PFJ	O13-C2-C1	-2.16	112.40	115.41
5	A	999	PFJ	O11-C1-C2	2.06	118.28	115.41

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	999	PFJ	C2-C1-O11-C12
5	A	999	PFJ	C4-C1-O11-C12


There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	997[B]	SO4	3	0
5	A	999	PFJ	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	305/362 (84%)	-0.33	7 (2%) 60 56	15, 23, 42, 71	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	492	CYS	6.3
1	A	488	CYS	5.8
1	A	453	ASN	5.6
1	A	490	THR	4.4
1	A	491	SER	4.1
1	A	489	GLY	3.4
1	A	496	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	997[B]	5/5	0.89	0.14	18,22,29,31	5

*Continued on next page...*






*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	997[A]	5/5	0.89	0.14	26,26,28,31	5
5	PFJ	A	999	30/30	0.92	0.09	24,30,37,38	0
4	MG	A	9002	1/1	0.99	0.09	20,20,20,20	0
3	ZN	A	9001	1/1	1.00	0.08	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

#### Electron density around PFJ A 999:

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.