



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 03:24 pm BST

PDB ID : 2OWP
Title : Crystal structure of a cystatin-like fold protein (bx_e_b1374) from burkholderia xenovorans lb400 at 2.00 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2007-02-16
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	129	
1	B	129	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2398 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein Bxe_B1374.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	129	Total	C	N	O	S	Se	0	1	0
			1022	646	181	190	1	4			
1	B	129	Total	C	N	O	S	Se	0	0	0
			1013	641	180	187	1	4			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP Q13MU9
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q13MU9
A	31	MSE	MET	MODIFIED RESIDUE	UNP Q13MU9
A	103	MSE	MET	MODIFIED RESIDUE	UNP Q13MU9
A	127	MSE	MET	MODIFIED RESIDUE	UNP Q13MU9
B	0	GLY	-	LEADER SEQUENCE	UNP Q13MU9
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q13MU9
B	31	MSE	MET	MODIFIED RESIDUE	UNP Q13MU9
B	103	MSE	MET	MODIFIED RESIDUE	UNP Q13MU9
B	127	MSE	MET	MODIFIED RESIDUE	UNP Q13MU9

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	148	Total O 148 148	0	0
4	B	149	Total O 149 149	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hypothetical protein Bxe_B1374

Chain A:  91% 9%



- Molecule 1: Hypothetical protein Bxe_B1374

Chain B:  91% 9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	80.52Å 80.52Å 116.26Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.89 – 2.00 29.90 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.5 (29.89-2.00) 98.5 (29.90-2.00)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.96 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.192 , 0.236 0.201 , 0.239	Depositor DCC
R_{free} test set	1504 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.775	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.039 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2398	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/1046	0.71	0/1418
1	B	0.72	0/1034	0.73	0/1402
All	All	0.72	0/2080	0.72	0/2820

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1022	0	1002	13	0
1	B	1013	0	992	14	0
2	A	10	0	0	0	0
3	A	32	0	48	2	0
3	B	24	0	36	1	0
4	A	148	0	0	0	0
4	B	149	0	0	0	0
All	All	2398	0	2078	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MSE:HE1	1:B:84:PHE:CE1	1.87	1.09
1:B:127:MSE:HE3	1:B:128:PRO:HD2	1.49	0.95
1:A:1:MSE:HE1	1:B:84:PHE:HE1	1.44	0.80
1:A:1:MSE:CE	1:B:84:PHE:CE1	2.65	0.80
1:A:1:MSE:SE	1:B:84:PHE:CE1	3.01	0.64
1:A:1:MSE:SE	1:B:84:PHE:CZ	3.01	0.64
1:A:61:GLU:OE1	3:A:135:EDO:O1	2.11	0.57
1:A:1:MSE:CE	1:B:84:PHE:HE1	2.11	0.56
1:A:1:MSE:HE1	1:B:84:PHE:CD1	2.39	0.54
1:B:127:MSE:HE3	1:B:128:PRO:CD	2.34	0.51
1:A:37:HIS:ND1	3:A:137:EDO:H12	2.28	0.48
1:B:45:GLY:HA3	1:B:48:GLU:OE2	2.16	0.46
1:B:2:GLU:OE2	1:B:5:GLN:HG2	2.16	0.46
1:A:100:GLY:HA2	1:A:124:LEU:HD23	2.00	0.44
1:B:28:ILE:HA	1:B:31:MSE:HE3	2.00	0.43
1:A:28:ILE:HA	1:A:31:MSE:HE3	2.00	0.43
1:A:45:GLY:HA3	1:A:48:GLU:OE2	2.19	0.42
1:A:37:HIS:CE1	1:A:57:ARG:HD2	2.55	0.42
1:B:122:VAL:HG11	3:B:134:EDO:H21	2.01	0.41
1:B:22:ALA:HB1	1:B:31:MSE:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/129 (99%)	125 (98%)	3 (2%)	0	100	100
1	B	127/129 (98%)	126 (99%)	1 (1%)	0	100	100
All	All	255/258 (99%)	251 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/102 (105%)	106 (99%)	1 (1%)	78	83
1	B	105/102 (103%)	104 (99%)	1 (1%)	76	81
All	All	212/204 (104%)	210 (99%)	2 (1%)	78	83

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	ARG
1	B	101	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	B	5	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	132	-	3,3,3	0.47	0	2,2,2	0.37	0
3	EDO	A	137	-	3,3,3	0.93	0	2,2,2	0.73	0
3	EDO	B	131	-	3,3,3	0.38	0	2,2,2	0.40	0
3	EDO	B	129	-	3,3,3	0.77	0	2,2,2	0.36	0
2	SO4	A	130	-	4,4,4	0.20	0	6,6,6	0.19	0
3	EDO	A	133	-	3,3,3	0.52	0	2,2,2	0.24	0
3	EDO	A	134	-	3,3,3	0.51	0	2,2,2	0.14	0
3	EDO	B	134	-	3,3,3	0.77	0	2,2,2	0.05	0
2	SO4	A	129	-	4,4,4	0.13	0	6,6,6	0.24	0
3	EDO	A	135	-	3,3,3	0.60	0	2,2,2	0.17	0
3	EDO	B	130	-	3,3,3	0.48	0	2,2,2	0.39	0
3	EDO	A	131	-	3,3,3	0.81	0	2,2,2	0.35	0
3	EDO	B	133	-	3,3,3	0.53	0	2,2,2	0.14	0
3	EDO	A	138	-	3,3,3	0.57	0	2,2,2	0.27	0
3	EDO	A	136	-	3,3,3	0.65	0	2,2,2	0.39	0
3	EDO	B	132	-	3,3,3	0.46	0	2,2,2	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	132	-	-	1/1/1/1	-
3	EDO	A	137	-	-	1/1/1/1	-
3	EDO	B	131	-	-	0/1/1/1	-
3	EDO	B	129	-	-	1/1/1/1	-
3	EDO	A	133	-	-	1/1/1/1	-
3	EDO	A	134	-	-	0/1/1/1	-
3	EDO	B	134	-	-	1/1/1/1	-
3	EDO	A	135	-	-	1/1/1/1	-
3	EDO	B	130	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	131	-	-	1/1/1/1	-
3	EDO	B	133	-	-	1/1/1/1	-
3	EDO	A	138	-	-	1/1/1/1	-
3	EDO	A	136	-	-	1/1/1/1	-
3	EDO	B	132	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	129	EDO	O1-C1-C2-O2
3	A	133	EDO	O1-C1-C2-O2
3	A	135	EDO	O1-C1-C2-O2
3	A	131	EDO	O1-C1-C2-O2
3	A	138	EDO	O1-C1-C2-O2
3	B	133	EDO	O1-C1-C2-O2
3	A	132	EDO	O1-C1-C2-O2
3	A	136	EDO	O1-C1-C2-O2
3	B	132	EDO	O1-C1-C2-O2
3	A	137	EDO	O1-C1-C2-O2
3	B	134	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	137	EDO	1	0
3	B	134	EDO	1	0
3	A	135	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.