



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 01:37 pm BST

PDB ID : 5OX8
Title : Structure of Enhanced Cyan Fluorescent Protein at pH 5.0
Authors : Gotthard, G.; von Stetten, D.; Clavel, D.; Noirclerc-Savoye, M.; Royant, A.
Deposited on : 2017-09-06
Resolution : 1.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2377 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	43	0
			2072	1333	335	399	5			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P42212
A	1	VAL	-	expression tag	UNP P42212
A	64	LEU	PHE	engineered mutation	UNP P42212
A	66	B2H	SER	chromophore	UNP P42212
A	66	B2H	TYR	chromophore	UNP P42212
A	66	B2H	GLY	chromophore	UNP P42212
A	146	ILE	ASN	engineered mutation	UNP P42212
A	153	THR	MET	engineered mutation	UNP P42212
A	163	ALA	VAL	conflict	UNP P42212
A	231	LEU	HIS	conflict	UNP P42212

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	296	Total	O	0	50
			304	304		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.25Å 62.62Å 69.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.66 – 1.29 39.66 – 1.29	Depositor EDS
% Data completeness (in resolution range)	98.6 (39.66-1.29) 98.6 (39.66-1.29)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 1.29Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.132 , 0.166 0.140 , 0.167	Depositor DCC
R_{free} test set	2841 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	15.7	Xtriage
Anisotropy	0.784	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2377	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, B2H

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	0/2172	0.84	2/2938 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	19[A]	ASP	CB-CG-OD1	5.82	123.53	118.30
1	A	19[B]	ASP	CB-CG-OD1	5.82	123.53	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2072	0	2037	13	0
2	A	1	0	0	0	0
3	A	304	0	0	6	0
All	All	2377	0	2037	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145[B]:TYR:CD1	1:A:205[B]:SER:O	1.75	1.39
1:A:145[B]:TYR:OH	1:A:204[B]:GLN:OE1	1.72	1.06
1:A:145[B]:TYR:CE1	1:A:205[B]:SER:O	2.12	1.03
1:A:145[B]:TYR:HD1	1:A:205[B]:SER:O	1.38	1.00
1:A:19[B]:ASP:OD2	3:A:401[B]:HOH:O	1.96	0.84
1:A:190[B]:ASP:HB2	3:A:647[B]:HOH:O	1.85	0.76
1:A:148[B]:HIS:NE2	3:A:403[B]:HOH:O	2.24	0.70
1:A:190[B]:ASP:CB	3:A:647[B]:HOH:O	2.49	0.59
1:A:45:LYS:HE2	1:A:47[A]:ILE:HD11	1.89	0.54
1:A:124[B]:GLU:CD	3:A:459[B]:HOH:O	2.48	0.52
1:A:203[A]:THR:HG23	1:A:224[A]:VAL:HG22	1.93	0.50
1:A:145[B]:TYR:CE2	1:A:147[B]:SER:HB2	2.51	0.46
1:A:204[A]:GLN:HA	3:A:563[A]:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/237 (111%)	261 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/206 (114%)	234 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	B2H	A	66[A]	-	22,26,27	2.14	7 (31%)	21,37,39	2.96	11 (52%)
1	B2H	A	66[B]	-	22,26,27	2.16	7 (31%)	21,37,39	3.01	12 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	B2H	A	66[A]	-	-	3/8/15/16	0/3/3/3
1	B2H	A	66[B]	-	-	3/8/15/16	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66[A]	B2H	O2-C2	-4.35	1.19	1.32
1	A	66[B]	B2H	O2-C2	-4.35	1.19	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66[A]	B2H	CB2-CA2	-4.00	1.48	1.51
1	A	66[B]	B2H	CB2-CA2	-4.00	1.48	1.51
1	A	66[A]	B2H	CZ3-CE3	3.79	1.45	1.36
1	A	66[B]	B2H	CZ3-CE3	3.79	1.45	1.36
1	A	66[A]	B2H	C1-N2	3.12	1.39	1.34
1	A	66[B]	B2H	C1-N2	3.12	1.39	1.34
1	A	66[A]	B2H	CA3-C3	-2.85	1.42	1.49
1	A	66[B]	B2H	CA3-C3	-2.85	1.42	1.49
1	A	66[A]	B2H	CZ2-CE2	2.72	1.46	1.41
1	A	66[B]	B2H	CZ2-CE2	2.72	1.46	1.41
1	A	66[A]	B2H	CE3-CD2	-2.61	1.36	1.42
1	A	66[B]	B2H	CE3-CD2	-2.61	1.36	1.42

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66[A]	B2H	C3-CA3-N3	6.53	121.82	111.92
1	A	66[B]	B2H	C3-CA3-N3	6.53	121.82	111.92
1	A	66[A]	B2H	CE3-CD2-CE2	6.48	126.76	118.17
1	A	66[B]	B2H	CE3-CD2-CE2	6.48	126.76	118.17
1	A	66[A]	B2H	CZ3-CH2-CZ2	5.13	127.63	120.44
1	A	66[B]	B2H	CZ3-CH2-CZ2	5.13	127.63	120.44
1	A	66[A]	B2H	CH2-CZ2-CE2	-4.09	114.20	120.08
1	A	66[B]	B2H	CH2-CZ2-CE2	-4.09	114.20	120.08
1	A	66[A]	B2H	CZ3-CE3-CD2	-3.27	116.36	120.89
1	A	66[B]	B2H	CZ3-CE3-CD2	-3.27	116.36	120.89
1	A	66[A]	B2H	CH2-CZ3-CE3	-2.64	116.75	120.44
1	A	66[B]	B2H	CH2-CZ3-CE3	-2.64	116.75	120.44
1	A	66[A]	B2H	CB2-CG2-CD2	-2.63	121.34	126.50
1	A	66[B]	B2H	CB2-CG2-CD2	-2.63	121.34	126.50
1	A	66[A]	B2H	CE3-CD2-CG2	-2.49	129.85	134.42
1	A	66[B]	B2H	CE3-CD2-CG2	-2.49	129.85	134.42
1	A	66[B]	B2H	CG1-CB1-CA1	2.40	117.83	112.16
1	A	66[A]	B2H	CD2-CE2-NE1	2.39	113.23	107.92
1	A	66[B]	B2H	CD2-CE2-NE1	2.39	113.23	107.92
1	A	66[A]	B2H	O3-C3-CA3	-2.26	119.88	126.32
1	A	66[B]	B2H	O3-C3-CA3	-2.26	119.88	126.32
1	A	66[A]	B2H	CG2-CB2-CA2	2.14	118.22	114.50
1	A	66[B]	B2H	CG2-CB2-CA2	2.14	118.22	114.50

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	66[A]	B2H	C2-CA2-CB2-CG2
1	A	66[B]	B2H	C2-CA2-CB2-CG2
1	A	66[A]	B2H	CA2-CB2-CG2-CD2
1	A	66[B]	B2H	CA2-CB2-CG2-CD2
1	A	66[A]	B2H	N2-CA2-CB2-CG2
1	A	66[B]	B2H	N2-CA2-CB2-CG2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/237 (94%)	-0.14	5 (2%) 62 61	14, 19, 31, 57	1 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	229	ILE	3.8
1	A	230	THR	3.7
1	A	151[A]	TYR	2.5
1	A	145[A]	TYR	2.2
1	A	190[A]	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	B2H	A	66[A]	24/25	0.96	0.07	13,16,18,18	3
1	B2H	A	66[B]	24/25	0.96	0.07	13,15,17,18	3

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	A	301	1/1	1.00	0.04	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.