



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 05:32 am BST

PDB ID : 6OX2  
Title : SETD3in Complex with an Actin Peptide with the Target Histidine Fully Methylated  
Authors : Horton, J.R.; Dai, S.; Cheng, X.  
Deposited on : 2019-05-13  
Resolution : 2.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Y	15	<div> <div>20%</div> <div> <div></div> <div>93%</div> <div>7%</div> </div> </div>
1	Z	15	<div> <div>33%</div> <div> <div></div> <div>100%</div> </div> </div>
2	A	599	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>•</div> <div>19%</div> </div> </div>
2	B	599	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>•</div> <div>19%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	B	608	-	-	-	X
5	GOL	B	610	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8766 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin Peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	Y	15	Total	C	N	O	0	0	0
			125	83	20	22			
1	Z	15	Total	C	N	O	0	0	0
			121	80	19	22			

- Molecule 2 is a protein called Histone-lysine N-methyltransferase setd3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	483	Total	C	N	O	S	0	10	0
			3920	2507	654	741	18			
2	B	483	Total	C	N	O	S	0	9	0
			3936	2522	656	739	19			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP Q86TU7
A	-4	PRO	-	expression tag	UNP Q86TU7
A	-3	LEU	-	expression tag	UNP Q86TU7
A	-2	GLY	-	expression tag	UNP Q86TU7
A	-1	SER	-	expression tag	UNP Q86TU7
B	-5	GLY	-	expression tag	UNP Q86TU7
B	-4	PRO	-	expression tag	UNP Q86TU7
B	-3	LEU	-	expression tag	UNP Q86TU7
B	-2	GLY	-	expression tag	UNP Q86TU7
B	-1	SER	-	expression tag	UNP Q86TU7

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula:  $C_{14}H_{20}N_6O_5S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	A	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	Y	7	Total	O	0	0
			7	7		
6	A	266	Total	O	0	0
			266	266		
6	Z	4	Total	O	0	0
			4	4		
6	B	237	Total	O	0	0
			237	237		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



#### • Molecule 1: Actin Peptide





#### • Molecule 1: Actin Peptide



#### • Molecule 2: Histone-lysine N-methyltransferase setd3



#### • Molecule 2: Histone-lysine N-methyltransferase setd3





N211		GLY
R214		VAL
Y218		GLN
D236		ASP
P258		ALA
D261		LEU
G262		ASN
S263		ILE
R284		ARG
W273		GLU
N280		ALA
F408		LYS
P416		THR
L456		GLU
I464		ASN
R495		GLY
L498		ASN
Y501		GLY
		THR
		ARG
		SER
		SER
		GLU
		ASN
		LEU
		GLY
		LEU
		SER
		LEU
		GLN
		ASN
		GLU
		SER
		VAL
		SER
		LYS
		ARG
		ALA
		SER
		VAL
		GLU
		ASP
		PRO
		LEU
		VAL
		LYS
		GLY
		ARG
		ASN
		SER
		LEU
		ASP
		GLU
		GLU
		THR
		ALA

GLY
VAL
LYS
GLU

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.10Å 175.94Å 66.26Å 90.00° 92.45° 90.00°	Depositor
Resolution (Å)	43.56 – 2.09 43.56 – 2.09	Depositor EDS
% Data completeness (in resolution range)	96.7 (43.56-2.09) 96.7 (43.56-2.09)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.32 (at 2.08Å)	Xtriage
Refinement program	PHENIX (1.14_3260)	Depositor
R, $R_{free}$	0.199 , 0.239 0.199 , 0.239	Depositor DCC
$R_{free}$ test set	1997 reflections (2.54%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.8	Xtriage
Anisotropy	0.762	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 52.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.064 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8766	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, HIC, EDO, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	Y	0.22	0/116	0.36	0/157
1	Z	0.22	0/112	0.39	0/153
2	A	0.24	0/4032	0.40	0/5465
2	B	0.24	0/4044	0.39	0/5480
All	All	0.24	0/8304	0.39	0/11255

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Y	125	0	116	2	0
1	Z	121	0	105	0	0
2	A	3920	0	3821	12	0
2	B	3936	0	3851	10	0
3	A	26	0	19	0	0
3	B	26	0	19	0	0
4	A	60	0	90	0	0
4	B	32	0	48	0	0
5	B	6	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	266	0	0	2	0
6	B	237	0	0	1	0
6	Y	7	0	0	0	0
6	Z	4	0	0	0	0
All	All	8766	0	8077	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:491:GLN:NE2	6:A:710:HOH:O	2.36	0.59
2:B:211:ASN:OD1	2:B:214:ARG:NH2	2.38	0.56
2:A:68:VAL:HG21	2:A:165:ALA:HA	1.87	0.56
2:A:83:LYS:NZ	2:A:87:GLU:OE2	2.38	0.56
2:A:339:LYS:HD2	2:A:366:VAL:HG22	1.89	0.53

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Y	12/15 (80%)	11 (92%)	1 (8%)	0	100	100
1	Z	12/15 (80%)	12 (100%)	0	0	100	100
2	A	491/599 (82%)	482 (98%)	9 (2%)	0	100	100
2	B	490/599 (82%)	484 (99%)	6 (1%)	0	100	100
All	All	1005/1228 (82%)	989 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Y	12/13 (92%)	12 (100%)	0	100	100
1	Z	11/13 (85%)	11 (100%)	0	100	100
2	A	423/521 (81%)	422 (100%)	1 (0%)	93	95
2	B	426/521 (82%)	426 (100%)	0	100	100
All	All	872/1068 (82%)	871 (100%)	1 (0%)	93	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	228	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	255	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	HIC	Y	73	1	8,11,12	0.75	0	6,14,16	0.52	0
1	HIC	Z	73	1	8,11,12	0.77	0	6,14,16	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HIC	Y	73	1	-	0/5/6/8	0/1/1/1
1	HIC	Z	73	1	-	0/5/6/8	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	Y	73	HIC	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	614	-	3,3,3	0.45	0	2,2,2	0.26	0
4	EDO	A	603	-	3,3,3	0.47	0	2,2,2	0.25	0
4	EDO	A	615	-	3,3,3	0.46	0	2,2,2	0.23	0
4	EDO	B	609	-	3,3,3	0.44	0	2,2,2	0.30	0
4	EDO	B	603	-	3,3,3	0.43	0	2,2,2	0.39	0
4	EDO	A	604	-	3,3,3	0.45	0	2,2,2	0.38	0
3	SAH	A	601	-	21,28,28	0.72	1 (4%)	20,40,40	0.52	0
4	EDO	B	605	-	3,3,3	0.44	0	2,2,2	0.31	0
4	EDO	B	608	-	3,3,3	0.44	0	2,2,2	0.31	0
4	EDO	B	607	-	3,3,3	0.46	0	2,2,2	0.32	0
4	EDO	A	609	-	3,3,3	0.44	0	2,2,2	0.44	0
4	EDO	B	604	-	3,3,3	0.45	0	2,2,2	0.33	0
4	EDO	A	607	-	3,3,3	0.45	0	2,2,2	0.42	0
4	EDO	A	611	-	3,3,3	0.45	0	2,2,2	0.31	0
4	EDO	B	606	-	3,3,3	0.46	0	2,2,2	0.32	0
4	EDO	A	606	-	3,3,3	0.45	0	2,2,2	0.34	0
4	EDO	A	613	-	3,3,3	0.42	0	2,2,2	0.49	0
4	EDO	B	602	-	3,3,3	0.45	0	2,2,2	0.33	0
4	EDO	A	605	-	3,3,3	0.45	0	2,2,2	0.32	0
4	EDO	A	610	-	3,3,3	0.45	0	2,2,2	0.32	0
5	GOL	B	610	-	5,5,5	0.90	0	5,5,5	0.98	0
4	EDO	A	612	-	3,3,3	0.45	0	2,2,2	0.32	0
4	EDO	A	616	-	3,3,3	0.46	0	2,2,2	0.31	0
4	EDO	A	602	-	3,3,3	0.45	0	2,2,2	0.27	0
4	EDO	A	608	-	3,3,3	0.46	0	2,2,2	0.30	0
3	SAH	B	601	-	21,28,28	0.80	1 (4%)	20,40,40	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	614	-	-	0/1/1/1	-
4	EDO	A	603	-	-	0/1/1/1	-
4	EDO	A	615	-	-	0/1/1/1	-
4	EDO	B	609	-	-	1/1/1/1	-
4	EDO	B	603	-	-	0/1/1/1	-
4	EDO	A	604	-	-	0/1/1/1	-
3	SAH	A	601	-	-	2/7/31/31	0/3/3/3
4	EDO	B	605	-	-	0/1/1/1	-
4	EDO	B	608	-	-	1/1/1/1	-

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	607	-	-	0/1/1/1	-
4	EDO	A	609	-	-	0/1/1/1	-
4	EDO	B	604	-	-	0/1/1/1	-
4	EDO	A	607	-	-	0/1/1/1	-
4	EDO	A	611	-	-	0/1/1/1	-
4	EDO	B	606	-	-	0/1/1/1	-
4	EDO	A	606	-	-	0/1/1/1	-
4	EDO	A	613	-	-	0/1/1/1	-
4	EDO	B	602	-	-	0/1/1/1	-
4	EDO	A	605	-	-	0/1/1/1	-
4	EDO	A	610	-	-	0/1/1/1	-
5	GOL	B	610	-	-	0/4/4/4	-
4	EDO	A	612	-	-	0/1/1/1	-
4	EDO	A	616	-	-	0/1/1/1	-
4	EDO	A	602	-	-	0/1/1/1	-
4	EDO	A	608	-	-	0/1/1/1	-
3	SAH	B	601	-	-	2/7/31/31	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	SAH	C2'-C1'	2.61	1.57	1.53
3	A	601	SAH	O4'-C1'	2.06	1.44	1.41

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	SAH	C-CA-CB-CG
3	B	601	SAH	C-CA-CB-CG
4	B	609	EDO	O1-C1-C2-O2
3	B	601	SAH	CB-CG-SD-C5'
3	A	601	SAH	CB-CG-SD-C5'


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will



also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	Y	14/15 (93%)	1.88	3 (21%) 0 0	46, 58, 119, 123	0
1	Z	14/15 (93%)	1.68	5 (35%) 0 0	64, 76, 118, 122	0
2	A	483/599 (80%)	0.19	17 (3%) 44 49	23, 40, 70, 97	0
2	B	483/599 (80%)	0.29	18 (3%) 41 46	24, 44, 77, 102	0
All	All	994/1228 (80%)	0.28	43 (4%) 35 40	23, 43, 79, 123	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Y	66	THR	7.8
2	A	20	SER	5.2
1	Z	66	THR	5.0
1	Y	67	LEU	4.7
1	Z	67	LEU	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	HIC	Z	73	11/12	0.94	0.13	65,67,70,72	0
1	HIC	Y	73	11/12	0.97	0.10	40,41,47,53	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	A	603	4/4	0.63	0.40	100,101,102,102	0
4	EDO	A	615	4/4	0.66	0.22	59,60,60,61	0
4	EDO	A	611	4/4	0.71	0.33	73,80,84,86	0
4	EDO	B	609	4/4	0.73	0.25	64,64,64,64	0
4	EDO	A	606	4/4	0.76	0.36	63,68,71,72	0
4	EDO	A	613	4/4	0.76	0.39	77,79,83,85	0
4	EDO	B	608	4/4	0.77	0.52	56,61,63,66	0
5	GOL	B	610	6/6	0.77	0.41	62,66,69,70	0
4	EDO	A	602	4/4	0.78	0.18	66,67,69,71	0
4	EDO	A	616	4/4	0.80	0.37	64,66,67,68	0
4	EDO	A	614	4/4	0.81	0.27	69,71,75,80	0
4	EDO	B	606	4/4	0.81	0.22	73,75,79,82	0
4	EDO	B	602	4/4	0.81	0.33	72,72,73,73	0
4	EDO	B	604	4/4	0.82	0.47	84,87,89,94	0
4	EDO	B	605	4/4	0.83	0.20	67,69,70,71	0
4	EDO	A	608	4/4	0.83	0.22	58,58,58,58	0
4	EDO	A	610	4/4	0.85	0.30	67,69,70,70	0
4	EDO	B	603	4/4	0.86	0.17	44,45,51,55	0
4	EDO	A	612	4/4	0.88	0.17	73,73,73,74	0
4	EDO	A	604	4/4	0.89	0.19	59,60,61,64	0
4	EDO	B	607	4/4	0.89	0.15	58,58,58,59	0
4	EDO	A	607	4/4	0.92	0.33	63,67,70,74	0
4	EDO	A	609	4/4	0.92	0.15	44,51,55,55	0
4	EDO	A	605	4/4	0.92	0.20	54,57,60,63	0
3	SAH	A	601	26/26	0.96	0.16	15,25,38,43	0
3	SAH	B	601	26/26	0.96	0.17	20,27,32,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around SAH A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers i

There are no such residues in this entry.