



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 11:52 am BST

PDB ID : 1OYF
Title : Crystal Structure of Russelles viper (*Daboia russellii pulchella*) phospholipase A2 in a complex with venom 6-methyl heptanol
Authors : Singh, N.; Jabeen, T.; Sharma, S.; Singh, T.P.
Deposited on : 2003-04-04
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	121	
2	B	121	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phospholipase A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	121	Total	C	N	O	S	0	1	0
			960	601	163	180	16			

- Molecule 2 is a protein called Phospholipase A2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	121	Total	C	N	O	S	0	0	0
			946	591	161	178	16			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 6-METHYLHEPTAN-1-OL (three-letter code: MHN) (formula: C₈H₁₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			9	8	1		
4	B	1	Total	C	O	0	0
			9	8	1		

- Molecule 5 is ACETIC ACID (three-letter code: ACY) (formula: $C_2H_4O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	76	Total 76	O 76	0	0
6	B	72	Total 72	O 72	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Phospholipase A2



• Molecule 2: Phospholipase A2



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.87Å 69.07Å 75.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.86 – 2.45	Depositor
% Data completeness (in resolution range)	79.1 (19.86-2.45)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.192 , 0.235	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2081	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MHN, ACY, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	5/982 (0.5%)	1.46	22/1319 (1.7%)
2	B	0.47	0/966	1.08	6/1295 (0.5%)
All	All	0.77	5/1948 (0.3%)	1.29	28/2614 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	32	GLY	N-CA	-16.32	1.21	1.46
1	A	36	THR	N-CA	14.09	1.74	1.46
1	A	29	CYS	C-N	-6.23	1.21	1.33
1	A	78	VAL	CA-CB	6.02	1.67	1.54
1	A	78	VAL	N-CA	5.07	1.56	1.46

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	ILE	C-N-CD	-21.80	72.63	120.60
2	B	19	ILE	C-N-CD	-21.23	73.91	120.60
2	B	19	ILE	C-N-CA	11.99	172.37	122.00
1	A	30	GLY	O-C-N	-11.80	103.81	122.70
1	A	31[A]	TRP	O-C-N	-11.64	103.42	123.20
1	A	31[B]	TRP	O-C-N	-11.64	103.42	123.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	ILE	C-N-CA	10.18	164.77	122.00
1	A	131	LYS	O-C-N	-10.13	106.50	122.70
2	B	127	LYS	N-CA-C	8.63	134.31	111.00
1	A	36	THR	CA-CB-CG2	-8.47	100.55	112.40
1	A	130	LEU	CA-CB-CG	7.95	133.59	115.30
1	A	30	GLY	CA-C-N	7.84	134.46	117.20
1	A	36	THR	N-CA-CB	-7.62	95.82	110.30
1	A	31[A]	TRP	CA-C-N	7.25	130.70	116.20
1	A	31[B]	TRP	CA-C-N	7.25	130.70	116.20
1	A	20	PRO	CA-N-CD	-7.11	101.55	111.50
1	A	105	CYS	CA-CB-SG	-6.62	102.08	114.00
1	A	36	THR	N-CA-C	-6.53	93.38	111.00
2	B	20	PRO	CA-N-CD	-6.37	102.58	111.50
1	A	30	GLY	N-CA-C	-6.28	97.39	113.10
2	B	129	GLU	N-CA-C	-6.09	94.55	111.00
1	A	78	VAL	O-C-N	5.59	131.65	122.70
1	A	31[A]	TRP	C-N-CA	5.55	133.95	122.30
1	A	31[B]	TRP	C-N-CA	5.55	133.95	122.30
1	A	36	THR	O-C-N	5.42	131.41	121.10
2	B	70	SER	N-CA-C	5.42	125.63	111.00
1	A	78	VAL	N-CA-CB	-5.18	100.10	111.50
1	A	131	LYS	N-CA-C	-5.10	97.22	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	31[A]	TRP	Mainchain,Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	960	0	910	48	0
2	B	946	0	907	32	0
3	A	5	0	0	0	0
4	A	9	0	18	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	9	0	18	0	0
5	A	4	0	3	0	0
6	A	76	0	0	1	0
6	B	72	0	0	3	0
All	All	2081	0	1856	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:THR:N	1:A:36:THR:CA	1.74	1.50
1:A:31[B]:TRP:CD1	2:B:43:ARG:HD2	2.01	0.95
1:A:19:ILE:HG22	1:A:20:PRO:CD	2.06	0.86
1:A:19:ILE:HG22	1:A:20:PRO:HD3	1.58	0.84
1:A:7:ARG:O	1:A:11:GLU:HG3	1.82	0.79
2:B:19:ILE:HG22	2:B:20:PRO:HD3	1.65	0.79
1:A:36:THR:N	1:A:36:THR:CB	2.46	0.79
1:A:31[B]:TRP:CG	2:B:43:ARG:HD2	2.20	0.77
1:A:50:CYS:SG	1:A:133:CYS:CB	2.72	0.77
1:A:36:THR:N	1:A:36:THR:C	2.38	0.75
2:B:19:ILE:HG22	2:B:20:PRO:CD	2.17	0.74
1:A:31[B]:TRP:CG	2:B:43:ARG:CD	2.72	0.73
1:A:50:CYS:CB	1:A:133:CYS:SG	2.75	0.69
2:B:19:ILE:CG2	2:B:20:PRO:HD3	2.24	0.67
1:A:50:CYS:CB	1:A:133:CYS:HG	2.01	0.65
2:B:54:ASN:C	2:B:56:PRO:HD3	2.15	0.65
1:A:23:SER:O	1:A:30:GLY:HA3	1.98	0.64
1:A:122:ASP:HB3	6:B:1072:HOH:O	1.98	0.64
1:A:35:GLY:C	1:A:36:THR:CA	2.64	0.63
1:A:19:ILE:HG22	1:A:20:PRO:HD2	1.78	0.63
1:A:19:ILE:CG2	1:A:20:PRO:HD3	2.29	0.62
1:A:114:SER:OG	1:A:116:LYS:HB3	2.00	0.61
1:A:61:CYS:O	1:A:68:PRO:HD3	2.01	0.61
1:A:31[B]:TRP:CB	2:B:43:ARG:CD	2.78	0.61
1:A:31[B]:TRP:CG	2:B:43:ARG:HD3	2.36	0.60
1:A:31[B]:TRP:CD1	2:B:43:ARG:CD	2.80	0.60
2:B:89:THR:OG1	2:B:92:GLU:HG3	2.01	0.59
1:A:131:LYS:O	1:A:133:CYS:C	2.37	0.59
1:A:30:GLY:C	1:A:31[B]:TRP:N	2.55	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:TYR:O	1:A:29:CYS:HB2	2.04	0.58
1:A:46:PHE:HZ	1:A:133:CYS:HA	1.69	0.57
1:A:19:ILE:CG2	1:A:20:PRO:CD	2.80	0.57
2:B:77:ARG:HA	2:B:81:ALA:O	2.07	0.55
2:B:121:PRO:HD2	2:B:125:LEU:HD12	1.88	0.55
1:A:104:ILE:O	1:A:108:GLN:HG3	2.07	0.53
1:A:86:LYS:HD2	1:A:86:LYS:N	2.22	0.53
1:A:89:THR:OG1	1:A:92:GLU:HG3	2.09	0.53
2:B:30:GLY:HA3	6:B:1015:HOH:O	2.10	0.52
2:B:19:ILE:HG22	2:B:20:PRO:HD2	1.92	0.52
2:B:104:ILE:O	2:B:108:GLN:HG3	2.09	0.51
2:B:19:ILE:CG2	2:B:20:PRO:CD	2.85	0.51
1:A:107:ARG:HG3	1:A:108:GLN:N	2.26	0.51
1:A:51:CYS:O	1:A:55:LEU:HD13	2.10	0.50
2:B:7:LYS:HE2	2:B:11:GLU:OE2	2.12	0.50
2:B:54:ASN:O	2:B:56:PRO:HD3	2.12	0.49
1:A:46:PHE:CZ	1:A:133:CYS:HA	2.47	0.49
2:B:114:SER:HB2	2:B:116:LYS:HG2	1.94	0.49
2:B:27:CYS:HB3	2:B:34:LYS:O	2.12	0.49
1:A:61:CYS:C	1:A:68:PRO:HD3	2.34	0.48
2:B:17:LEU:O	2:B:21:SER:HB2	2.14	0.47
1:A:31[B]:TRP:CB	2:B:43:ARG:HD3	2.44	0.47
1:A:31[B]:TRP:CB	2:B:43:ARG:HD2	2.42	0.47
2:B:5:PHE:O	2:B:9:ILE:HG13	2.14	0.47
1:A:88:GLY:H	1:A:93:ASN:HD21	1.63	0.47
1:A:1:SER:OG	1:A:72:ARG:HD2	2.17	0.45
1:A:31[B]:TRP:HB2	2:B:43:ARG:CD	2.46	0.45
2:B:61:CYS:HB3	6:B:1060:HOH:O	2.16	0.45
2:B:3:LEU:HD12	2:B:3:LEU:HA	1.77	0.45
1:A:91:CYS:O	1:A:95:ILE:HG13	2.17	0.44
2:B:12:GLU:O	2:B:107:ARG:HD3	2.18	0.44
2:B:38:LYS:HD2	2:B:120:TYR:CE2	2.52	0.43
1:A:35:GLY:HA3	1:A:126:CYS:HB3	2.00	0.43
1:A:77:ARG:HA	1:A:81:ALA:O	2.19	0.43
1:A:33:GLY:HA3	6:A:3010:HOH:O	2.18	0.42
1:A:55:LEU:N	1:A:56:PRO:HD3	2.34	0.42
1:A:46:PHE:CZ	1:A:133:CYS:SG	3.13	0.42
2:B:9:ILE:O	2:B:13:THR:HG23	2.21	0.41
1:A:68:PRO:HA	1:A:95:ILE:HD13	2.01	0.41
1:A:54:ASN:C	1:A:56:PRO:HD3	2.40	0.41
1:A:121:PRO:C	1:A:124:PHE:N	2.73	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:124:PHE:CE2	2:B:125:LEU:HG	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/121 (99%)	114 (95%)	3 (2%)	3 (2%)	5	3
2	B	119/121 (98%)	112 (94%)	5 (4%)	2 (2%)	9	7
All	All	239/242 (99%)	226 (95%)	8 (3%)	5 (2%)	7	5

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	20	PRO
2	B	20	PRO
1	A	37	PRO
1	A	33	GLY
2	B	33	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/104 (101%)	92 (88%)	13 (12%)	4	4
2	B	104/104 (100%)	94 (90%)	10 (10%)	8	8
All	All	209/208 (100%)	186 (89%)	23 (11%)	6	5

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LEU
1	A	3	LEU
1	A	7	ARG
1	A	20	PRO
1	A	49	ASP
1	A	55	LEU
1	A	86	LYS
1	A	105	CYS
1	A	122	ASP
1	A	127	LYS
1	A	129	GLU
1	A	130	LEU
1	A	133	CYS
2	B	2	LEU
2	B	3	LEU
2	B	13	THR
2	B	20	PRO
2	B	69	LYS
2	B	72	ARG
2	B	76	LYS
2	B	114	SER
2	B	127	LYS
2	B	129	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
2	B	54	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MHN	A	1001	-	8,8,8	0.86	0	8,8,8	0.47	0
5	ACY	A	3001	-	1,3,3	2.44	1 (100%)	0,3,3	0.00	-
4	MHN	B	1002	-	8,8,8	0.47	0	8,8,8	1.80	2 (25%)
3	SO4	A	2001	-	4,4,4	1.74	2 (50%)	6,6,6	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MHN	A	1001	-	-	2/6/6/6	-
4	MHN	B	1002	-	-	1/6/6/6	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2001	SO4	O1-S	2.81	1.61	1.46
5	A	3001	ACY	CH3-C	2.44	1.51	1.48
3	A	2001	SO4	O3-S	-2.00	1.31	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1002	MHN	C5-C6-C7	-3.32	97.58	114.42
4	B	1002	MHN	C6-C5-C4	3.00	124.24	113.62

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1001	MHN	C1-C2-C4-C5
4	A	1001	MHN	C3-C2-C4-C5
4	B	1002	MHN	C1-C2-C4-C5

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	30:GLY	C	31[B]:TRP	N	2.55

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.