



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2020 – 11:13 pm BST

PDB ID : 3OYP
Title : HCV NS3/4A in complex with ligand 3
Authors : Hagel, M.; Niu, D.; St.Martin, T.; Sheets, M.P.; Qiao, L.; Bernard, H.; Karp, R.M.; Zhu, Z.; Labenski, M.T.; Chaturvedi, P.C.; Nacht, M.; Westlin, W.F.; Petter, R.C.; Singh, J.
Deposited on : 2010-09-23
Resolution : 2.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	187	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>7%</div> </div> </div>
1	B	187	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>6%</div> </div> </div>
2	C	16	<div> <div></div> <div> <div>69%</div> <div>6%</div> <div>25%</div> </div> </div>
2	D	16	<div> <div></div> <div> <div>63%</div> <div>6%</div> <div>6%</div> <div>25%</div> </div> </div>
3	E	5	<div> <div></div> <div> <div>80%</div> <div>20%</div> </div> </div>
3	F	5	<div> <div></div> <div> <div>80%</div> <div>20%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2827 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine protease NS3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	S	0	0	0
			1278	798	230	240	10			
1	B	175	Total	C	N	O	S	0	0	0
			1283	801	231	241	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	114	ILE	VAL	VARIANT	UNP P26662
A	132	VAL	ILE	VARIANT	UNP P26662
B	114	ILE	VAL	VARIANT	UNP P26662
B	132	VAL	ILE	VARIANT	UNP P26662

- Molecule 2 is a protein called Non-structural protein 4A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	12	Total	C	N	O	0	0	0
			84	55	15	14			
2	D	12	Total	C	N	O	0	0	0
			84	55	15	14			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	220	LYS	-	EXPRESSION TAG	UNP P26662
C	235	LYS	-	EXPRESSION TAG	UNP P26662
D	220	LYS	-	EXPRESSION TAG	UNP P26662
D	235	LYS	-	EXPRESSION TAG	UNP P26662

- Molecule 3 is a protein called Peptidomimetic inhibitor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	E	5	Total 43	Br 1	C 29	N 5	O 7	S 1	0	0	0
3	F	5	Total 43	Br 1	C 29	N 5	O 7	S 1	0	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Zn 1	0	0
4	A	1	Total 1	Zn 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total 5	O 5	0	0
5	B	2	Total 2	O 2	0	0
5	C	2	Total 2	O 2	0	0
5	D	1	Total 1	O 1	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine protease NS3



- Molecule 1: Serine protease NS3




- Molecule 2: Non-structural protein 4A




- Molecule 2: Non-structural protein 4A



- Molecule 3: Peptidomimetic inhibitor



- Molecule 3: Peptidomimetic inhibitor



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	94.40 Å 94.40 Å 83.13 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	81.65 – 2.76 47.20 – 2.76	Depositor EDS
% Data completeness (in resolution range)	100.0 (81.65-2.76) 100.0 (47.20-2.76)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 2.77 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.215 , 0.274 0.215 , 0.272	Depositor DCC
R_{free} test set	762 reflections (6.98%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 17.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.148 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2827	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0Y8, 0Y9, ZN, DAL, PPI, 0YA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1305	0.51	0/1778
1	B	0.33	0/1310	0.53	0/1786
2	C	0.40	0/83	0.61	0/111
2	D	0.40	0/83	0.60	0/111
All	All	0.33	0/2781	0.52	0/3786

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1278	0	1286	10	0
1	B	1283	0	1293	10	0
2	C	84	0	99	0	0
2	D	84	0	99	1	0
3	E	43	0	30	2	0
3	F	43	0	30	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	2	0	0	0	0
5	C	2	0	0	0	0
5	D	1	0	0	0	0
All	All	2827	0	2837	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:ARG:HD2	3:F:3:0Y8:BR1	2.49	0.68
1:A:158:VAL:CG1	1:A:166:ALA:HB3	2.29	0.62
1:B:75:TYR:OH	1:B:80:GLN:NE2	2.32	0.61
1:B:158:VAL:CG1	1:B:166:ALA:HB3	2.34	0.56
1:A:154:PHE:CE2	3:E:4:0Y9:H14	2.46	0.51
1:A:139:SER:OG	3:E:4:0Y9:C26	2.60	0.50
1:A:75:TYR:OH	1:A:80:GLN:NE2	2.45	0.49
1:A:36:LEU:HD11	1:A:45:ALA:HB2	1.95	0.48
1:B:8:GLN:HE21	2:D:228:ARG:HE	1.62	0.48
1:B:97:CYS:SG	1:B:145:CYS:HB2	2.54	0.47
1:A:104:LEU:CD2	1:A:151:VAL:HG21	2.45	0.47
1:A:77:ASN:HD22	1:A:77:ASN:C	2.19	0.45
1:A:104:LEU:HD22	1:A:151:VAL:HG21	1.99	0.45
1:B:118:ARG:HD2	1:B:120:GLY:O	2.17	0.44
1:B:77:ASN:CB	1:B:80:GLN:HE21	2.31	0.44
1:A:14:LEU:HD12	1:A:18:ILE:HD11	2.00	0.44
1:B:104:LEU:HD22	1:B:151:VAL:HG11	2.01	0.42
1:B:77:ASN:HD22	1:B:77:ASN:C	2.21	0.42
1:A:130:ARG:HG2	1:A:134:TYR:CD2	2.54	0.42
1:B:114:ILE:HA	1:B:115:PRO:HD3	1.92	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	172/187 (92%)	165 (96%)	7 (4%)	0	100	100
1	B	173/187 (92%)	167 (96%)	6 (4%)	0	100	100
2	C	10/16 (62%)	10 (100%)	0	0	100	100
2	D	10/16 (62%)	10 (100%)	0	0	100	100
All	All	365/406 (90%)	352 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/154 (92%)	135 (95%)	7 (5%)	25	43
1	B	142/154 (92%)	133 (94%)	9 (6%)	18	31
2	C	10/13 (77%)	9 (90%)	1 (10%)	7	13
2	D	10/13 (77%)	8 (80%)	2 (20%)	1	1
All	All	304/334 (91%)	285 (94%)	19 (6%)	18	31

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	74	MET
1	A	77	ASN
1	A	95	THR
1	A	98	THR
1	A	99	CYS
1	A	130	ARG
1	B	9	GLN
1	B	14	LEU
1	B	28	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	40	THR
1	B	74	MET
1	B	77	ASN
1	B	130	ARG
1	B	155	ARG
1	B	163	VAL
2	C	228	ARG
2	D	222	SER
2	D	228	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	77	ASN
1	A	80	GLN
1	B	8	GLN
1	B	27	ASN
1	B	34	GLN
1	B	77	ASN
1	B	80	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	0Y9	F	4	3	5,8,9	0.88	0	7,12,14	3.00	3 (42%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	0Y9	E	4	3	5,8,9	0.90	0	7,12,14	3.07	3 (42%)
3	0Y8	F	3	3	16,21,22	1.43	2 (12%)	21,29,31	2.09	3 (14%)
3	0Y8	E	3	3	16,21,22	1.45	2 (12%)	21,29,31	2.32	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	0Y9	F	4	3	-	2/2/13/16	0/1/1/1
3	0Y9	E	4	3	-	0/2/13/16	0/1/1/1
3	0Y8	F	3	3	-	0/4/15/17	0/3/3/3
3	0Y8	E	3	3	-	0/4/15/17	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	3	0Y8	C4-C6	-3.25	1.39	1.43
3	F	3	0Y8	C4-C6	-2.81	1.40	1.43
3	F	3	0Y8	C1-N2	2.24	1.39	1.34
3	E	3	0Y8	C1-N2	2.21	1.39	1.34

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	4	0Y9	C26-C21-N20	-6.79	106.83	116.39
3	E	4	0Y9	C26-C21-N20	-6.75	106.88	116.39
3	E	3	0Y8	C4-O12-C13	6.45	127.83	118.56
3	F	3	0Y8	C6-C4-N2	-5.84	119.46	124.37
3	E	3	0Y8	C6-C4-N2	-5.47	119.77	124.37
3	F	3	0Y8	C4-O12-C13	5.00	125.74	118.56
3	E	3	0Y8	C3-C1-N2	-4.09	119.78	123.81
3	F	3	0Y8	C3-C1-N2	-3.83	120.04	123.81
3	E	4	0Y9	O27-C26-C21	-3.17	117.49	125.31
3	F	4	0Y9	O27-C26-C21	-2.81	118.39	125.31
3	E	4	0Y9	C22-C24-C25	-2.79	116.93	125.42
3	E	3	0Y8	C13-C14-C15	-2.59	99.73	102.92
3	F	4	0Y9	C22-C24-C25	-2.48	117.88	125.42

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	4	0Y9	C23-C22-C24-C25
3	F	4	0Y9	C21-C22-C24-C25

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	4	0Y9	2	0
3	F	3	0Y8	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	174/187 (93%)	0.24	1 (0%) 89 92	25, 37, 49, 57	1 (0%)
1	B	175/187 (93%)	0.34	5 (2%) 51 61	27, 39, 50, 58	0
2	C	12/16 (75%)	0.12	0 100 100	28, 32, 40, 42	0
2	D	12/16 (75%)	0.05	0 100 100	26, 28, 42, 43	0
3	E	0/5	-	-	-	-
3	F	0/5	-	-	-	-
All	All	373/416 (89%)	0.27	6 (1%) 72 79	25, 37, 49, 58	1 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	40	THR	2.6
1	B	1	ALA	2.5
1	B	121	ASP	2.4
1	A	18	ILE	2.2
1	B	39	ALA	2.2
1	B	14	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	0Y8	F	3	19/20	0.91	0.22	29,42,45,48	1
3	0Y9	E	4	8/9	0.94	0.16	28,35,37,39	0
3	DAL	F	2	5/6	0.95	0.18	34,35,37,37	0
3	0Y8	E	3	19/20	0.95	0.20	34,40,42,45	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	0Y9	F	4	8/9	0.97	0.22	33,38,41,41	0
3	DAL	E	2	5/6	0.97	0.21	34,35,38,40	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	ZN	A	201	1/1	0.96	0.14	41,41,41,41	0
4	ZN	B	201	1/1	0.96	0.11	65,65,65,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.