



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:55 am BST

PDB ID : 4OY7
Title : Structure of cellulose active LPMO CelS2 (ScLPMO10C) in complex with Copper.
Authors : Forsberg, Z.; Mackenzie, A.K.; Sorlie, M.; Rohr, A.K.; Helland, R.; Arvai, A.S.; Vaaje-Kolstad, G.; Eijsink, V.G.H.
Deposited on : 2014-02-11
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	196	<div><div></div><div>96% .</div></div>
1	B	196	<div><div>%</div><div>95% 5%</div></div>
1	C	196	<div><div>2%</div><div>93% 7%</div></div>
1	D	196	<div><div>3%</div><div>98% ..</div></div>
1	E	196	<div><div>%</div><div>95% . .</div></div>
1	F	196	<div><div>%</div><div>96% . .</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	196	<div><div></div><div>2%</div><div>98%</div><div></div><div>..</div></div>
1	H	196	<div><div></div><div>2%</div><div>97%</div><div></div><div>..</div></div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative secreted cellulose binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	2	0
			1496	935	255	299	7			
1	B	196	Total	C	N	O	S	0	4	0
			1507	941	256	302	8			
1	C	196	Total	C	N	O	S	0	2	0
			1492	934	253	297	8			
1	D	196	Total	C	N	O	S	0	2	0
			1496	935	255	299	7			
1	E	195	Total	C	N	O	S	0	3	0
			1496	936	254	298	8			
1	F	195	Total	C	N	O	S	0	2	0
			1490	933	253	296	8			
1	G	196	Total	C	N	O	S	0	2	0
			1492	933	253	299	7			
1	H	195	Total	C	N	O	S	0	3	0
			1493	935	253	297	8			

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Cu	0	0
			1	1		
2	D	1	Total	Cu	0	0
			1	1		
2	E	1	Total	Cu	0	0
			1	1		
2	H	1	Total	Cu	0	0
			1	1		
2	B	1	Total	Cu	0	0
			1	1		
2	C	1	Total	Cu	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Cu 1	0	0
2	F	1	Total 1	Cu 1	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	1	Total 1	Ca 1	0	0
3	G	1	Total 1	Ca 1	0	0
3	B	1	Total 1	Ca 1	0	0
3	E	1	Total 1	Ca 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	248	Total 248	O 248	0	0
4	B	250	Total 250	O 250	0	0
4	C	225	Total 225	O 225	0	0
4	D	221	Total 221	O 221	0	0
4	E	268	Total 269	O 269	0	1
4	F	271	Total 271	O 271	0	0
4	G	225	Total 225	O 225	0	0
4	H	214	Total 214	O 214	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative secreted cellulose binding protein

Chain A:  96%





- Molecule 1: Putative secreted cellulose binding protein

Chain B:  95% 5%




- Molecule 1: Putative secreted cellulose binding protein

Chain C:  93% 7%



- Molecule 1: Putative secreted cellulose binding protein

Chain D:  98%



- Molecule 1: Putative secreted cellulose binding protein

Chain E:  95%




- Molecule 1: Putative secreted cellulose binding protein



- Molecule 1: Putative secreted cellulose binding protein



- Molecule 1: Putative secreted cellulose binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.86Å 122.86Å 156.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.90 – 1.50 47.89 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.0 (47.90-1.50) 99.1 (47.89-1.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.194 , 0.222 0.200 , 0.227	Depositor DCC
R_{free} test set	12750 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	10.2	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13897	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.1728e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/1543	0.73	2/2111 (0.1%)
1	B	0.53	0/1554	0.76	1/2127 (0.0%)
1	C	0.52	0/1542	0.76	4/2110 (0.2%)
1	D	0.51	0/1543	0.75	1/2111 (0.0%)
1	E	0.56	0/1543	0.79	3/2114 (0.1%)
1	F	0.55	0/1537	0.80	5/2105 (0.2%)
1	G	0.51	0/1542	0.74	3/2110 (0.1%)
1	H	0.53	0/1543	0.75	2/2113 (0.1%)
All	All	0.53	0/12347	0.76	21/16901 (0.1%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	125	ARG	NE-CZ-NH2	9.74	125.17	120.30
1	F	125	ARG	NE-CZ-NH1	-8.77	115.91	120.30
1	H	108	ARG	NE-CZ-NH2	-8.61	116.00	120.30
1	G	108	ARG	NE-CZ-NH1	8.53	124.56	120.30
1	E	108	ARG	NE-CZ-NH1	8.38	124.49	120.30
1	D	108	ARG	NE-CZ-NH1	8.09	124.34	120.30
1	C	108	ARG	NE-CZ-NH1	7.99	124.30	120.30
1	H	108	ARG	NE-CZ-NH1	7.98	124.29	120.30
1	G	108	ARG	NE-CZ-NH2	-7.86	116.37	120.30
1	F	108	ARG	NE-CZ-NH1	7.81	124.20	120.30
1	E	108	ARG	NE-CZ-NH2	-7.37	116.62	120.30
1	A	108	ARG	NE-CZ-NH1	7.35	123.98	120.30
1	B	125	ARG	NE-CZ-NH1	6.86	123.73	120.30
1	C	108	ARG	NE-CZ-NH2	-6.77	116.91	120.30
1	F	108	ARG	NE-CZ-NH2	-6.68	116.96	120.30
1	G	125	ARG	NE-CZ-NH2	6.50	123.55	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	99	ASP	CB-CG-OD1	5.96	123.66	118.30
1	A	108	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	E	125	ARG	NE-CZ-NH2	5.63	123.11	120.30
1	C	166	ASP	CB-CG-OD1	5.34	123.10	118.30
1	C	125	ARG	NE-CZ-NH2	-5.13	117.74	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1496	0	1357	3	0
1	B	1507	0	1368	5	0
1	C	1492	0	1359	6	0
1	D	1496	0	1357	2	0
1	E	1496	0	1361	4	0
1	F	1490	0	1356	4	0
1	G	1492	0	1354	2	0
1	H	1493	0	1360	2	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	B	1	0	0	0	0
3	E	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
4	A	248	0	0	0	0
4	B	250	0	0	2	0
4	C	225	0	0	1	0
4	D	221	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	269	0	0	2	0
4	F	271	0	0	1	0
4	G	225	0	0	1	0
4	H	214	0	0	1	0
All	All	13897	0	10872	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:53:LYS:NZ	4:H:614:HOH:O	1.98	0.97
1:E:40[B]:MET:HE3	4:E:487:HOH:O	1.77	0.83
1:B:40[B]:MET:HG3	4:B:1247:HOH:O	1.81	0.80
1:B:158[B]:SER:OG	4:B:1233:HOH:O	2.07	0.72
1:F:40[B]:MET:HE2	4:F:588:HOH:O	1.96	0.66
1:D:108:ARG:NH2	4:D:571:HOH:O	2.30	0.60
1:C:62:THR:HG23	4:C:521:HOH:O	2.01	0.59
1:F:40[B]:MET:HA	1:F:40[B]:MET:HE3	1.84	0.59
1:B:59:LEU:HD12	1:B:75:ALA:HB1	1.87	0.57
1:E:40[B]:MET:HE2	4:E:544:HOH:O	2.09	0.53
1:D:59:LEU:HD12	1:D:75:ALA:HB1	1.93	0.50
1:E:46:TYR:O	1:E:49:GLN:HG3	2.11	0.49
1:A:59:LEU:HD12	1:A:75:ALA:HB1	1.95	0.48
1:A:35:HIS:HA	1:A:219:PHE:CE2	2.49	0.48
1:F:35:HIS:HA	1:F:219:PHE:CE2	2.49	0.48
1:C:59:LEU:HD12	1:C:75:ALA:HB1	1.97	0.47
1:C:46:TYR:O	1:C:49:GLN:HG3	2.15	0.47
1:C:35:HIS:HA	1:C:219:PHE:CE2	2.49	0.47
1:A:46:TYR:O	1:A:49:GLN:HG3	2.16	0.46
1:C:101:THR:OG1	1:C:106:GLY:HA2	2.17	0.44
1:G:108:ARG:NH2	4:G:583:HOH:O	2.37	0.42
1:H:46:TYR:O	1:H:49:GLN:HG3	2.19	0.42
1:G:101:THR:OG1	1:G:106:GLY:HA2	2.20	0.41
1:E:101:THR:OG1	1:E:106:GLY:HA2	2.20	0.41
1:B:35:HIS:HA	1:B:219:PHE:CE2	2.56	0.41
1:C:210:TRP:HB3	1:C:219:PHE:HB2	2.03	0.40
1:F:85:LEU:C	1:F:85:LEU:HD12	2.42	0.40
1:B:150:VAL:HB	1:B:172:GLN:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/196 (100%)	193 (98%)	3 (2%)	0	100	100
1	B	198/196 (101%)	194 (98%)	4 (2%)	0	100	100
1	C	196/196 (100%)	193 (98%)	3 (2%)	0	100	100
1	D	196/196 (100%)	195 (100%)	1 (0%)	0	100	100
1	E	196/196 (100%)	191 (97%)	5 (3%)	0	100	100
1	F	195/196 (100%)	189 (97%)	6 (3%)	0	100	100
1	G	196/196 (100%)	189 (96%)	7 (4%)	0	100	100
1	H	196/196 (100%)	191 (97%)	5 (3%)	0	100	100
All	All	1569/1568 (100%)	1535 (98%)	34 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	155/153 (101%)	155 (100%)	0	100	100
1	B	157/153 (103%)	157 (100%)	0	100	100
1	C	155/153 (101%)	155 (100%)	0	100	100
1	D	155/153 (101%)	155 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	156/153 (102%)	154 (99%)	2 (1%)	69	44
1	F	155/153 (101%)	155 (100%)	0	100	100
1	G	155/153 (101%)	155 (100%)	0	100	100
1	H	156/153 (102%)	156 (100%)	0	100	100
All	All	1244/1224 (102%)	1242 (100%)	2 (0%)	93	86

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	158[A]	SER
1	E	158[B]	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	196/196 (100%)	0.11	0 100 100	5, 10, 19, 23	0
1	B	196/196 (100%)	0.07	2 (1%) 82 85	5, 9, 16, 22	0
1	C	196/196 (100%)	0.17	4 (2%) 65 70	6, 10, 18, 24	0
1	D	196/196 (100%)	0.22	5 (2%) 56 61	6, 10, 20, 27	0
1	E	195/196 (99%)	0.06	1 (0%) 91 93	4, 8, 14, 20	0
1	F	195/196 (99%)	0.07	1 (0%) 91 93	5, 8, 15, 20	0
1	G	196/196 (100%)	0.23	3 (1%) 73 78	6, 10, 19, 30	0
1	H	195/196 (99%)	0.20	3 (1%) 73 78	6, 11, 20, 28	0
All	All	1565/1568 (99%)	0.14	19 (1%) 79 82	4, 10, 18, 30	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	230	GLY	5.9
1	F	229	GLY	5.5
1	E	229	GLY	4.7
1	D	229	GLY	4.1
1	G	229	GLY	3.5
1	D	230	GLY	3.3
1	D	156	GLY	2.8
1	H	229	GLY	2.7
1	H	166	ASP	2.4
1	C	230	GLY	2.4
1	D	60	ASP	2.3
1	B	230	GLY	2.3
1	C	156	GLY	2.3
1	C	60	ASP	2.3
1	B	72	GLN	2.2
1	H	185	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	182	SER	2.1
1	C	229	GLY	2.1
1	D	185	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	B	302	1/1	0.98	0.18	25,25,25,25	0
2	CU	B	301	1/1	0.99	0.07	19,19,19,19	1
2	CU	G	301	1/1	0.99	0.09	8,8,8,8	1
2	CU	D	301	1/1	0.99	0.05	17,17,17,17	1
2	CU	H	301	1/1	0.99	0.06	9,9,9,9	1
2	CU	A	301	1/1	0.99	0.06	25,25,25,25	1
3	CA	H	302	1/1	0.99	0.24	32,32,32,32	0
2	CU	E	301	1/1	1.00	0.05	13,13,13,13	1
3	CA	E	302	1/1	1.00	0.10	15,15,15,15	0
2	CU	C	301	1/1	1.00	0.04	15,15,15,15	1
3	CA	G	302	1/1	1.00	0.06	11,11,11,11	0
2	CU	F	301	1/1	1.00	0.04	11,11,11,11	1

6.5 Other polymers [i](#)

There are no such residues in this entry.