



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 01:57 am BST

PDB ID : 3P56
Title : The structure of the human RNase H2 complex defines key interaction interfaces relevant to enzyme function and human disease
Authors : Bubeck, D.; Graham, S.C.; Jones, E.Y.
Deposited on : 2010-10-08
Resolution : 4.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1127 (4.42-3.70)
Clashscore	141614	1033 (4.40-3.72)
Ramachandran outliers	138981	1145 (4.42-3.70)
Sidechain outliers	138945	1133 (4.42-3.70)
RSRZ outliers	127900	1005 (4.44-3.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	<div> <div>2%</div> <div> <div></div> <div>70%</div> <div>13%</div> <div>17%</div> </div> </div>
1	D	299	<div> <div>6%</div> <div> <div></div> <div>64%</div> <div>11%</div> <div>25%</div> </div> </div>
2	B	237	<div> <div>%</div> <div> <div></div> <div>59%</div> <div>6%</div> <div>34%</div> </div> </div>
2	E	237	<div> <div>5%</div> <div> <div></div> <div>60%</div> <div>6%</div> <div>33%</div> </div> </div>
3	C	164	<div> <div>4%</div> <div> <div></div> <div>62%</div> <div>9%</div> <div>29%</div> </div> </div>
3	F	164	<div> <div>7%</div> <div> <div></div> <div>61%</div> <div>10%</div> <div>29%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonuclease H2 subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	0	0
			1875	1203	325	337	10			
1	D	225	Total	C	N	O	S	0	0	0
			1690	1085	294	302	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	ALA	ASP	ENGINEERED MUTATION	UNP O75792
A	169	ALA	ASP	ENGINEERED MUTATION	UNP O75792
D	34	ALA	ASP	ENGINEERED MUTATION	UNP O75792
D	169	ALA	ASP	ENGINEERED MUTATION	UNP O75792

- Molecule 2 is a protein called Ribonuclease H2 subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	156	Total	C	N	O	S	0	0	0
			1072	691	189	189	3			
2	E	158	Total	C	N	O	S	0	0	0
			1088	702	191	192	3			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	GLY	-	EXPRESSION TAG	UNP Q5TBB1
B	-9	PRO	-	EXPRESSION TAG	UNP Q5TBB1
B	-8	LEU	-	EXPRESSION TAG	UNP Q5TBB1
B	-7	GLY	-	EXPRESSION TAG	UNP Q5TBB1
B	-6	SER	-	EXPRESSION TAG	UNP Q5TBB1
B	-5	PRO	-	EXPRESSION TAG	UNP Q5TBB1
B	-4	GLU	-	EXPRESSION TAG	UNP Q5TBB1
B	-3	PHE	-	EXPRESSION TAG	UNP Q5TBB1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	PRO	-	EXPRESSION TAG	UNP Q5TBB1
B	-1	GLY	-	EXPRESSION TAG	UNP Q5TBB1
B	0	ARG	-	EXPRESSION TAG	UNP Q5TBB1
B	1	LEU	-	EXPRESSION TAG	UNP Q5TBB1
E	-10	GLY	-	EXPRESSION TAG	UNP Q5TBB1
E	-9	PRO	-	EXPRESSION TAG	UNP Q5TBB1
E	-8	LEU	-	EXPRESSION TAG	UNP Q5TBB1
E	-7	GLY	-	EXPRESSION TAG	UNP Q5TBB1
E	-6	SER	-	EXPRESSION TAG	UNP Q5TBB1
E	-5	PRO	-	EXPRESSION TAG	UNP Q5TBB1
E	-4	GLU	-	EXPRESSION TAG	UNP Q5TBB1
E	-3	PHE	-	EXPRESSION TAG	UNP Q5TBB1
E	-2	PRO	-	EXPRESSION TAG	UNP Q5TBB1
E	-1	GLY	-	EXPRESSION TAG	UNP Q5TBB1
E	0	ARG	-	EXPRESSION TAG	UNP Q5TBB1
E	1	LEU	-	EXPRESSION TAG	UNP Q5TBB1


- Molecule 3 is a protein called Ribonuclease H2 subunit C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	116	Total	C	N	O	S	0	0	0
			817	523	149	143	2			
3	F	116	Total	C	N	O	S	0	0	0
			817	523	149	143	2			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Ribonuclease H2 subunit A



• Molecule 1: Ribonuclease H2 subunit A




• Molecule 2: Ribonuclease H2 subunit B







• Molecule 2: Ribonuclease H2 subunit B



• Molecule 3: Ribonuclease H2 subunit C



• Molecule 3: Ribonuclease H2 subunit C



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	212.24Å 42.30Å 186.95Å 90.00° 98.11° 90.00°	Depositor
Resolution (Å)	30.00 – 4.06 29.81 – 4.06	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-4.06) 95.6 (29.81-4.06)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 4.11Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.9.2	Depositor
R, R_{free}	0.374 , 0.379 0.401 , 0.414	Depositor DCC
R_{free} test set	659 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	125.4	Xtriage
Anisotropy	0.667	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.20 , 98.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.35$, $\langle L^2 \rangle = 0.18$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.80	EDS
Total number of atoms	7359	wwPDB-VP
Average B, all atoms (Å ²)	188.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.12 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8675e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.41	0/1916	0.60	0/2607
1	D	0.40	0/1726	0.60	0/2347
2	B	0.38	0/1096	0.56	0/1498
2	E	0.38	0/1113	0.56	0/1521
3	C	0.38	0/838	0.59	0/1147
3	F	0.38	0/838	0.58	0/1147
All	All	0.39	0/7527	0.59	0/10267

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1875	0	1815	16	0
1	D	1690	0	1627	13	0
2	B	1072	0	854	5	0
2	E	1088	0	869	7	0
3	C	817	0	772	6	0
3	F	817	0	772	7	0
All	All	7359	0	6709	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:GLU:HB3	1:A:229:GLY:H	1.51	0.75
1:D:225:GLU:HB3	1:D:229:GLY:H	1.51	0.74
1:D:96:LEU:HD11	1:D:120:THR:HG21	1.79	0.65
1:A:96:LEU:HD11	1:A:120:THR:HG21	1.79	0.64
1:A:98:PRO:HG2	1:A:233:PHE:HA	1.86	0.56
1:D:98:PRO:HG2	1:D:233:PHE:HA	1.86	0.56
3:C:76:PRO:HB2	3:C:79:LEU:HD12	1.87	0.56
1:D:12:GLY:HA3	1:D:100:LEU:HD13	1.88	0.56
2:E:212:ALA:HA	2:E:215:LEU:HD12	1.88	0.56
2:B:212:ALA:HA	2:B:215:LEU:HD12	1.88	0.55
1:A:12:GLY:HA3	1:A:100:LEU:HD13	1.89	0.55
3:F:76:PRO:HB2	3:F:79:LEU:HD12	1.89	0.55
1:D:24:CYS:HB3	1:D:133:VAL:HG13	1.90	0.54
1:A:24:CYS:HB3	1:A:133:VAL:HG13	1.91	0.52
1:D:33:VAL:HB	1:D:140:VAL:HG13	1.92	0.52
1:D:20:VAL:HB	1:D:25:ARG:HH21	1.75	0.51
1:A:33:VAL:HB	1:A:140:VAL:HG13	1.93	0.51
1:A:20:VAL:HB	1:A:25:ARG:HH21	1.76	0.50
2:B:74:PHE:HB2	3:C:30:HIS:HB2	1.93	0.49
1:D:41:VAL:HG22	1:D:238:TRP:CD1	2.47	0.49
2:B:68:GLU:HG3	3:C:143:LYS:HD2	1.95	0.49
1:A:153:LEU:HB3	1:A:162:VAL:HG11	1.94	0.49
1:D:299:LEU:HB2	2:E:50:ILE:HG21	1.94	0.48
1:A:41:VAL:HG22	1:A:238:TRP:CD1	2.48	0.48
1:A:110:LYS:HE3	1:A:110:LYS:H	1.79	0.48
3:C:60:GLU:HG3	3:C:69:ARG:HD2	1.96	0.47
3:F:60:GLU:HG3	3:F:69:ARG:HD2	1.96	0.47
1:A:14:CYS:HB3	1:A:95:VAL:HG13	1.97	0.47
2:E:74:PHE:HB2	3:F:30:HIS:HB2	1.97	0.46
2:E:174:THR:HG22	3:F:157:ILE:HG12	1.97	0.46
1:A:94:ASP:HB2	1:A:124:LEU:HD21	1.98	0.46
1:A:64:LYS:HD3	1:A:173:PRO:HA	1.98	0.46
3:C:17:ARG:HH11	3:C:19:ALA:HB3	1.80	0.45
1:D:94:ASP:HB2	1:D:124:LEU:HD21	1.98	0.45
1:D:64:LYS:HD3	1:D:173:PRO:HA	1.99	0.45
2:E:66:PHE:HB3	2:E:85:LEU:HB3	1.99	0.45
3:F:17:ARG:HH11	3:F:19:ALA:HB3	1.81	0.44
2:B:66:PHE:HB3	2:B:85:LEU:HB3	1.99	0.44
1:D:14:CYS:HB3	1:D:95:VAL:HG13	1.99	0.44
1:A:113:LEU:HD12	1:A:116:LEU:HD23	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:20:VAL:HA	2:E:88:ALA:HB3	2.00	0.44
1:D:113:LEU:HD12	1:D:116:LEU:HD23	2.00	0.43
2:B:20:VAL:HA	2:B:88:ALA:HB3	2.01	0.43
3:F:45:GLY:HA2	3:F:49:THR:HB	2.01	0.42
3:C:45:GLY:HA2	3:C:49:THR:HB	2.01	0.41
1:A:149:TYR:CD2	1:A:164:VAL:HG11	2.55	0.41
1:A:48:ALA:HB2	1:A:93:LEU:HD23	2.03	0.40
2:E:15:GLN:HG2	3:F:85:VAL:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/299 (80%)	224 (94%)	13 (5%)	2 (1%)	19	58
1	D	213/299 (71%)	199 (93%)	12 (6%)	2 (1%)	17	55
2	B	144/237 (61%)	133 (92%)	10 (7%)	1 (1%)	22	61
2	E	146/237 (62%)	135 (92%)	10 (7%)	1 (1%)	22	61
3	C	112/164 (68%)	106 (95%)	5 (4%)	1 (1%)	17	55
3	F	112/164 (68%)	105 (94%)	6 (5%)	1 (1%)	17	55
All	All	966/1400 (69%)	902 (93%)	56 (6%)	8 (1%)	19	58

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	37	ALA
3	F	37	ALA
1	A	22	ALA
2	B	76	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	76	ASN
1	D	22	ALA
1	A	108	ARG
1	D	108	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/255 (73%)	177 (95%)	10 (5%)	22	51
1	D	167/255 (66%)	158 (95%)	9 (5%)	22	50
2	B	81/208 (39%)	74 (91%)	7 (9%)	10	36
2	E	82/208 (39%)	76 (93%)	6 (7%)	14	41
3	C	72/132 (54%)	68 (94%)	4 (6%)	21	49
3	F	72/132 (54%)	68 (94%)	4 (6%)	21	49
All	All	661/1190 (56%)	621 (94%)	40 (6%)	18	47

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LEU
1	A	35	GLU
1	A	47	TYR
1	A	94	ASP
1	A	110	LYS
1	A	148	THR
1	A	167	LYS
1	A	175	VAL
1	A	179	SER
1	A	285	HIS
2	B	26	ASP
2	B	74	PHE
2	B	91	VAL
2	B	94	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	98	LEU
2	B	136	LYS
2	B	218	ASP
3	C	16	LEU
3	C	21	LEU
3	C	57	GLU
3	C	130	THR
1	D	31	LEU
1	D	35	GLU
1	D	47	TYR
1	D	94	ASP
1	D	110	LYS
1	D	167	LYS
1	D	175	VAL
1	D	179	SER
1	D	285	HIS
2	E	26	ASP
2	E	74	PHE
2	E	94	LEU
2	E	98	LEU
2	E	136	LYS
2	E	218	ASP
3	F	16	LEU
3	F	21	LEU
3	F	57	GLU
3	F	130	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	GLN
2	B	86	HIS
1	D	118	HIS
2	E	86	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/299 (83%)	-0.06	7 (2%) 53 42	58, 146, 212, 243	0
1	D	225/299 (75%)	0.38	18 (8%) 12 11	165, 213, 267, 278	0
2	B	156/237 (65%)	-0.08	3 (1%) 66 58	132, 190, 228, 249	0
2	E	158/237 (66%)	0.40	13 (8%) 11 10	179, 224, 248, 270	0
3	C	116/164 (70%)	0.06	6 (5%) 27 24	115, 156, 223, 251	0
3	F	116/164 (70%)	0.42	12 (10%) 6 6	176, 208, 244, 256	0
All	All	1020/1400 (72%)	0.18	59 (5%) 23 19	58, 196, 243, 278	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	ASN	7.6
3	F	72	GLU	6.7
1	A	11	THR	6.1
3	F	146	GLY	5.7
2	E	23	TYR	5.3
3	F	17	ARG	4.9
1	D	221	LYS	4.8
1	A	194	GLN	4.4
1	D	37	GLY	4.2
3	F	23	ASP	3.9
2	E	174	THR	3.8
2	E	100	TYR	3.7
1	A	16	LEU	3.5
2	E	217	SER	3.3
1	A	193	TRP	3.3
1	D	17	SER	3.2
3	C	146	GLY	3.1
3	F	22	ARG	3.1
1	D	284	SER	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	16	LEU	3.0
2	E	105	ASP	3.0
1	D	16	LEU	3.0
1	D	194	GLN	2.9
3	F	71	GLU	2.9
1	D	229	GLY	2.9
2	E	218	ASP	2.9
2	E	25	LYS	2.8
2	E	157	LYS	2.8
1	D	92	ALA	2.8
2	E	125	CYS	2.7
1	D	256	ASP	2.7
1	D	226	PRO	2.6
3	F	143	LYS	2.6
3	F	147	ALA	2.4
2	E	132	PRO	2.4
3	C	17	ARG	2.4
3	C	121	GLY	2.3
2	B	218	ASP	2.3
1	D	76	ARG	2.3
1	D	193	TRP	2.3
1	D	87	ASP	2.3
3	C	120	ILE	2.3
3	F	18	SER	2.3
3	C	122	ALA	2.3
3	F	141	ASP	2.2
2	B	23	TYR	2.2
1	D	213	ASP	2.2
1	A	87	ASP	2.2
2	B	217	SER	2.2
1	D	89	VAL	2.1
1	A	221	LYS	2.1
2	E	104	ALA	2.1
1	D	10	ASN	2.1
2	E	207	ASP	2.0
3	F	53	ARG	2.0
2	E	126	ILE	2.0
3	F	145	ARG	2.0
1	D	90	GLY	2.0
1	D	225	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.