



# Full wwPDB NMR Structure Validation Report ⓘ

Sep 27, 2022 – 11:10 am BST

PDB ID : 7P5Q  
BMRB ID : 34652  
Title : NMR structure of a peptide deriving from SARS-CoV-2 Lineage B.1.1.7 S  
RBD 482-506 fragment in HFIP/H<sub>2</sub>O  
Authors : Santoro, A.; Buonocore, M.; Grimaldi, M.; D'Ursi, A.M.  
Deposited on : 2021-07-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.31.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 62%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	25	<div> <div>32%</div> <div>8%</div> <div>60%</div> </div>

## 2 Ensemble composition and analysis ⓘ

This entry contains 40 models. Model 29 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:490-A:499 (10)	0.06	29

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: Inconsistent models

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 374 atoms, of which 174 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms						Trace
1	A	25	Total	C	H	N	O	S	0
			374	133	174	29	37	1	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	501	TYR	ASN	engineered mutation	UNP P0DTC2

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Spike glycoprotein




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: Spike glycoprotein




#### 4.2.2 Score per residue for model 2

- Molecule 1: Spike glycoprotein




### 4.2.3 Score per residue for model 3

- Molecule 1: Spike glycoprotein




### 4.2.4 Score per residue for model 4

- Molecule 1: Spike glycoprotein




### 4.2.5 Score per residue for model 5

- Molecule 1: Spike glycoprotein




### 4.2.6 Score per residue for model 6

- Molecule 1: Spike glycoprotein



### 4.2.7 Score per residue for model 7

- Molecule 1: Spike glycoprotein




### 4.2.8 Score per residue for model 8

- Molecule 1: Spike glycoprotein




### 4.2.9 Score per residue for model 9

- Molecule 1: Spike glycoprotein




### 4.2.10 Score per residue for model 10

- Molecule 1: Spike glycoprotein




### 4.2.11 Score per residue for model 11

- Molecule 1: Spike glycoprotein




### 4.2.12 Score per residue for model 12

- Molecule 1: Spike glycoprotein




### 4.2.13 Score per residue for model 13

- Molecule 1: Spike glycoprotein




### 4.2.14 Score per residue for model 14

- Molecule 1: Spike glycoprotein




### 4.2.15 Score per residue for model 15

- Molecule 1: Spike glycoprotein




### 4.2.16 Score per residue for model 16

- Molecule 1: Spike glycoprotein



### 4.2.17 Score per residue for model 17


- Molecule 1: Spike glycoprotein






#### 4.2.18 Score per residue for model 18

- Molecule 1: Spike glycoprotein




#### 4.2.19 Score per residue for model 19

- Molecule 1: Spike glycoprotein



#### 4.2.20 Score per residue for model 20

- Molecule 1: Spike glycoprotein




#### 4.2.21 Score per residue for model 21

- Molecule 1: Spike glycoprotein




#### 4.2.22 Score per residue for model 22

- Molecule 1: Spike glycoprotein




### 4.2.23 Score per residue for model 23

- Molecule 1: Spike glycoprotein




### 4.2.24 Score per residue for model 24

- Molecule 1: Spike glycoprotein




### 4.2.25 Score per residue for model 25

- Molecule 1: Spike glycoprotein




### 4.2.26 Score per residue for model 26

- Molecule 1: Spike glycoprotein




### 4.2.27 Score per residue for model 27

- Molecule 1: Spike glycoprotein




#### 4.2.28 Score per residue for model 28

- Molecule 1: Spike glycoprotein




#### 4.2.29 Score per residue for model 29 (medoid)

- Molecule 1: Spike glycoprotein




#### 4.2.30 Score per residue for model 30

- Molecule 1: Spike glycoprotein




#### 4.2.31 Score per residue for model 31

- Molecule 1: Spike glycoprotein




#### 4.2.32 Score per residue for model 32

- Molecule 1: Spike glycoprotein




#### 4.2.33 Score per residue for model 33

- Molecule 1: Spike glycoprotein




#### 4.2.34 Score per residue for model 34

- Molecule 1: Spike glycoprotein



#### 4.2.35 Score per residue for model 35

- Molecule 1: Spike glycoprotein




#### 4.2.36 Score per residue for model 36

- Molecule 1: Spike glycoprotein




#### 4.2.37 Score per residue for model 37

- Molecule 1: Spike glycoprotein




#### 4.2.38 Score per residue for model 38

- Molecule 1: Spike glycoprotein




#### 4.2.39 Score per residue for model 39

- Molecule 1: Spike glycoprotein



#### 4.2.40 Score per residue for model 40

- Molecule 1: Spike glycoprotein



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 40 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	
TALOS-N	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	196
Number of shifts mapped to atoms	196
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	62%

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	84	76	76	0±0
All	All	3360	3040	3040	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:490:PHE:O	1:A:492:LEU:HD23	0.50	2.06	3	1

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	10/25 (40%)	9±0 (88±4%)	1±0 (12±4%)	0±0 (0±0%)	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	400/1000 (40%)	353 (88%)	47 (12%)	0 (0%)	100	100

There are no Ramachandran outliers.

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	9/20 (45%)	6±1 (64±13%)	3±1 (36±13%)	1	8
All	All	360/800 (45%)	231 (64%)	129 (36%)	1	8

All 7 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	494	SER	40
1	A	498	GLN	25
1	A	492	LEU	18
1	A	495	TYR	17
1	A	493	GLN	16
1	A	497	PHE	12
1	A	490	PHE	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.6 Ligand geometry

There are no ligands in this entry.

## 6.7 Other polymers

There are no such molecules in this entry.

## 6.8 Polymer linkage issues

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 62% for the well-defined parts and 61% for the entire structure.

### 7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch\_output*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	196
Number of shifts mapped to atoms	196
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 62%, i.e. 81 atoms were assigned a chemical shift out of a possible 131. 1 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	24/46 (52%)	18/18 (100%)	6/20 (30%)	0/8 (0%)
Sidechain	49/59 (83%)	32/37 (86%)	17/20 (85%)	0/2 (0%)
Aromatic	8/26 (31%)	8/14 (57%)	0/12 (0%)	0/0 (—%)
Overall	81/131 (62%)	58/69 (84%)	23/52 (44%)	0/10 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 61%, i.e. 179 atoms were assigned a chemical shift out of a possible 294. 1 out of 3 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	66/121 (55%)	47/48 (98%)	19/50 (38%)	0/23 (0%)
Sidechain	89/114 (78%)	57/69 (83%)	32/41 (78%)	0/4 (0%)
Aromatic	24/59 (41%)	24/31 (77%)	0/28 (0%)	0/0 (—%)
Overall	179/294 (61%)	128/148 (86%)	51/119 (43%)	0/27 (0%)

#### 7.1.4 Statistically unusual chemical shifts ⓘ

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

