



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 01:04 am BST

PDB ID : 2P7I  
Title : CRYSTAL STRUCTURE OF a SAM dependent methyl-transferase type 12 family protein (ECA1738) FROM PECTOBACTERIUM ATROSEPTICUM SCRI1043 AT 1.74 Å RESOLUTION  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-03-20  
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	<div> <div>6%</div> <div> <div></div> <div>82%</div> <div>8%</div> <div>10%</div> </div> </div>
1	B	250	<div> <div>8%</div> <div> <div></div> <div>86%</div> <div>5%</div> <div>8%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TRS	A	251	-	X	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4131 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	Se	0	13	0
			1858	1189	322	337	7	3			
1	B	229	Total	C	N	O	S	Se	0	8	0
			1872	1200	322	340	7	3			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP Q6D6E7
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q6D6E7
A	28	MSE	MET	MODIFIED RESIDUE	UNP Q6D6E7
A	32	MSE	MET	MODIFIED RESIDUE	UNP Q6D6E7
A	154	MSE	MET	MODIFIED RESIDUE	UNP Q6D6E7
B	0	GLY	-	LEADER SEQUENCE	UNP Q6D6E7
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q6D6E7
B	28	MSE	MET	MODIFIED RESIDUE	UNP Q6D6E7
B	32	MSE	MET	MODIFIED RESIDUE	UNP Q6D6E7
B	154	MSE	MET	MODIFIED RESIDUE	UNP Q6D6E7

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		
3	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Na	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	186	Total	O	0	5
			191	191		
6	B	180	Total	O	0	4
			184	184		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hypothetical protein



- Molecule 1: Hypothetical protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.83 Å   120.83 Å   149.97 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	29.11 – 1.74 29.11 – 1.74	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.11-1.74) 100.0 (29.11-1.74)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.90 (at 1.74 Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, $R_{free}$	0.155   ,   0.180 0.165   ,   0.192	Depositor DCC
$R_{free}$ test set	2883 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.5	Xtriage
Anisotropy	0.477	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 57.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4131	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, TRS, MPD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.69	0/1936	0.77	0/2614
1	B	0.65	0/1938	0.71	0/2620
All	All	0.67	0/3874	0.74	0/5234

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1858	0	1830	15	0
1	B	1872	0	1819	10	0
2	A	1	0	0	0	0
3	A	8	0	12	3	0
3	B	8	0	12	2	0
4	A	8	0	14	2	0
5	B	1	0	0	0	0
6	A	191	0	0	4	0
6	B	184	0	0	1	0
All	All	4131	0	3687	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:252:MPD:H52	4:A:252:MPD:HM1	1.74	0.69
1:B:32:MSE:SE	1:B:109[B]:THR:HG21	2.43	0.68
1:A:39:PHE:CD1	1:A:242:LEU:HD22	2.30	0.66
4:A:252:MPD:H52	4:A:252:MPD:CM	2.25	0.66
3:A:251:TRS:N	6:A:362:HOH:O	2.31	0.61
1:A:187:ARG:HG3	6:A:371:HOH:O	2.00	0.61
1:A:109[B]:THR:HG23	3:A:251:TRS:H11	1.84	0.60
1:A:109[B]:THR:HG23	3:A:251:TRS:C1	2.35	0.57
1:B:109[A]:THR:HG22	1:B:139:VAL:HB	1.87	0.56
1:A:169:ALA:HA	6:A:307:HOH:O	2.06	0.55
1:A:44:ASN:HD22	1:A:101[B]:ARG:CZ	2.22	0.51
1:A:208:TRP:HA	1:A:211:ILE:HG22	1.94	0.50
1:B:109[A]:THR:OG1	3:B:251:TRS:H11	2.13	0.49
1:B:114:HIS:HD2	6:B:361:HOH:O	1.96	0.49
1:B:154:MSE:HE2	1:B:156:ILE:HD11	1.95	0.47
1:A:107:VAL:HG13	1:A:109[A]:THR:HG23	1.97	0.46
1:A:124[B]:ARG:HD3	1:A:128:ASP:OD2	2.16	0.45
1:B:44:ASN:HD22	1:B:101:ARG:NH1	2.14	0.45
1:A:124[B]:ARG:NH1	1:A:128:ASP:OD2	2.50	0.44
1:A:83:LEU:HD13	1:A:87[B]:ILE:CD1	2.48	0.44
1:B:90:ILE:HD13	1:B:99:LEU:CD2	2.47	0.43
1:B:90:ILE:HD13	1:B:99:LEU:HD22	1.99	0.43
1:A:22:ASN:N	1:A:26:ASP:OD2	2.53	0.42
1:B:73:GLU:O	1:B:76:ILE:HG22	2.19	0.41
1:A:66:ILE:HB	1:A:87[A]:ILE:HD13	2.03	0.41
1:A:62:HIS:HE1	6:A:379:HOH:O	2.04	0.41
1:A:44:ASN:HD22	1:A:101[B]:ARG:NH2	2.19	0.41
1:B:32:MSE:SE	1:B:109[B]:THR:CG2	3.16	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/250 (94%)	233 (99%)	3 (1%)	0	100	100
1	B	235/250 (94%)	232 (99%)	3 (1%)	0	100	100
All	All	471/500 (94%)	465 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/209 (96%)	197 (98%)	3 (2%)	65	47
1	B	199/209 (95%)	198 (100%)	1 (0%)	88	83
All	All	399/418 (96%)	395 (99%)	4 (1%)	73	63

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	ARG
1	A	194	TYR
1	A	219	LYS
1	B	219	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	TRS	A	251	-	7,7,7	0.44	0	9,9,9	1.17	1 (11%)
3	TRS	B	251	-	7,7,7	0.42	0	9,9,9	1.22	1 (11%)
4	MPD	A	252	-	7,7,7	0.44	0	9,10,10	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	A	251	-	-	9/9/9/9	-
3	TRS	B	251	-	-	3/9/9/9	-
4	MPD	A	252	-	-	1/5/5/5	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	251	TRS	O1-C1-C	2.99	120.47	111.00
3	B	251	TRS	O1-C1-C	2.09	117.61	111.00

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	251	TRS	C1-C-C3-O3
3	A	251	TRS	C2-C-C3-O3
3	A	251	TRS	N-C-C3-O3
4	A	252	MPD	C2-C3-C4-O4
3	A	251	TRS	C2-C-C1-O1
3	A	251	TRS	C3-C-C1-O1
3	A	251	TRS	C1-C-C2-O2
3	A	251	TRS	N-C-C2-O2
3	B	251	TRS	C1-C-C2-O2
3	B	251	TRS	N-C-C2-O2
3	A	251	TRS	C3-C-C2-O2
3	B	251	TRS	C3-C-C2-O2
3	A	251	TRS	N-C-C1-O1

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	251	TRS	3	0
3	B	251	TRS	2	0
4	A	252	MPD	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	222/250 (88%)	0.12	16 (7%) 15 19	22, 29, 54, 63	0
1	B	226/250 (90%)	0.28	19 (8%) 11 14	23, 30, 53, 73	0
All	All	448/500 (89%)	0.20	35 (7%) 13 17	22, 29, 54, 73	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	LEU	7.4
1	A	214	THR	4.8
1	A	213	GLN	4.5
1	A	211	ILE	4.4
1	A	217	LEU	4.4
1	A	206	PHE	4.3
1	A	25	PHE	4.0
1	B	21	TYR	3.9
1	B	82	ARG	3.9
1	B	86	GLY	3.7
1	B	48	LEU	3.5
1	A	216	ILE	3.3
1	B	168	PHE	3.2
1	A	86	GLY	3.2
1	B	139	VAL	3.1
1	A	82	ARG	3.0
1	B	109[A]	THR	2.8
1	B	83	LEU	2.7
1	B	25[A]	PHE	2.7
1	B	165	GLU	2.6
1	B	166	ALA	2.6
1	A	83	LEU	2.6
1	A	85	ASP	2.6
1	A	209	ASP	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	22	ASN	2.4
1	B	76	ILE	2.3
1	B	107	VAL	2.3
1	B	79	ALA	2.3
1	B	170	HIS	2.2
1	B	111	VAL	2.2
1	A	77[A]	SER	2.2
1	B	108	LEU	2.2
1	B	78	HIS	2.1
1	B	138	LEU	2.1
1	A	108	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	TRS	A	251	8/8	0.77	0.22	43,66,68,69	0
3	TRS	B	251	8/8	0.78	0.19	51,63,66,70	0
4	MPD	A	252	8/8	0.89	0.17	45,53,66,74	0
5	NA	B	250	1/1	0.99	0.12	30,30,30,30	0
2	CL	A	250	1/1	0.99	0.08	32,32,32,32	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.