



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 10, 2020 – 12:50 PM BST

PDB ID : 3P7H  
Title : Structure of the human Langerin carbohydrate recognition domain in complex with maltose  
Authors : Skerra, A.; Schiefner, A.  
Deposited on : 2010-10-12  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	<div> <div>4%</div> <div> <div></div> <div>87%</div> <div>8%</div> </div> </div>
1	B	146	<div> <div>82%</div> <div>7%</div> <div>11%</div> </div>
1	C	146	<div> <div>%</div> <div>79%</div> <div>9%</div> <div>12%</div> </div>
1	D	146	<div> <div>2%</div> <div>87%</div> <div>11%</div> </div>
2	E	2	<div> <div>100%</div> </div>
2	F	2	<div> <div>50%</div> <div>50%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	G	2	 50%50%
2	H	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	F	1	-	-	-	X
2	GLC	G	1	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C-type lectin domain family 4 member K.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	135	Total	C	N	O	S	Se	0	0	0
			1097	713	178	201	4	1			
1	B	130	Total	C	N	O	S	Se	0	1	0
			1068	693	175	195	4	1			
1	C	129	Total	C	N	O	S	Se	0	0	0
			1058	691	172	190	4	1			
1	D	130	Total	C	N	O	S	Se	0	0	0
			1059	688	173	193	4	1			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	329	SER	-	expression tag	UNP Q9UJ71
A	330	ALA	-	expression tag	UNP Q9UJ71
A	331	TRP	-	expression tag	UNP Q9UJ71
A	332	SER	-	expression tag	UNP Q9UJ71
A	333	HIS	-	expression tag	UNP Q9UJ71
A	334	PRO	-	expression tag	UNP Q9UJ71
A	335	GLN	-	expression tag	UNP Q9UJ71
A	336	PHE	-	expression tag	UNP Q9UJ71
A	337	GLU	-	expression tag	UNP Q9UJ71
A	338	LYS	-	expression tag	UNP Q9UJ71
B	329	SER	-	expression tag	UNP Q9UJ71
B	330	ALA	-	expression tag	UNP Q9UJ71
B	331	TRP	-	expression tag	UNP Q9UJ71
B	332	SER	-	expression tag	UNP Q9UJ71
B	333	HIS	-	expression tag	UNP Q9UJ71
B	334	PRO	-	expression tag	UNP Q9UJ71
B	335	GLN	-	expression tag	UNP Q9UJ71
B	336	PHE	-	expression tag	UNP Q9UJ71
B	337	GLU	-	expression tag	UNP Q9UJ71
B	338	LYS	-	expression tag	UNP Q9UJ71
C	329	SER	-	expression tag	UNP Q9UJ71

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	330	ALA	-	expression tag	UNP Q9UJ71
C	331	TRP	-	expression tag	UNP Q9UJ71
C	332	SER	-	expression tag	UNP Q9UJ71
C	333	HIS	-	expression tag	UNP Q9UJ71
C	334	PRO	-	expression tag	UNP Q9UJ71
C	335	GLN	-	expression tag	UNP Q9UJ71
C	336	PHE	-	expression tag	UNP Q9UJ71
C	337	GLU	-	expression tag	UNP Q9UJ71
C	338	LYS	-	expression tag	UNP Q9UJ71
D	329	SER	-	expression tag	UNP Q9UJ71
D	330	ALA	-	expression tag	UNP Q9UJ71
D	331	TRP	-	expression tag	UNP Q9UJ71
D	332	SER	-	expression tag	UNP Q9UJ71
D	333	HIS	-	expression tag	UNP Q9UJ71
D	334	PRO	-	expression tag	UNP Q9UJ71
D	335	GLN	-	expression tag	UNP Q9UJ71
D	336	PHE	-	expression tag	UNP Q9UJ71
D	337	GLU	-	expression tag	UNP Q9UJ71
D	338	LYS	-	expression tag	UNP Q9UJ71

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	2	Total	C	O	0	0	0
			23	12	11			
2	F	2	Total	C	O	0	0	0
			23	12	11			
2	G	2	Total	C	O	0	0	0
			23	12	11			
2	H	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Ca 1	0	0
3	D	1	Total 1	Ca 1	0	0
3	C	1	Total 1	Ca 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	50	Total 50	O 50	0	0
4	B	61	Total 61	O 61	0	0
4	C	52	Total 52	O 52	0	0
4	D	41	Total 41	O 41	0	0



Chain F:  50% 50%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain G:  50% 50%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain H:  100%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.97Å 79.97Å 90.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 28.11 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.5 (30.00-2.30) 98.6 (28.11-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.56 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.181 , 0.229 0.176 , 0.224	Depositor DCC
$R_{free}$ test set	1280 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.4	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 31.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.047 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4582	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.54	0/1138	0.58	0/1550
1	B	0.57	0/1106	0.60	0/1504
1	C	0.60	0/1097	0.61	0/1492
1	D	0.55	0/1097	0.59	0/1492
All	All	0.57	0/4438	0.59	0/6038

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1097	0	1016	3	0
1	B	1068	0	993	6	0
1	C	1058	0	982	7	0
1	D	1059	0	986	2	0
2	E	23	0	20	0	0
2	F	23	0	20	0	0
2	G	23	0	19	1	0
2	H	23	0	21	0	0
3	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	50	0	0	0	0
4	B	61	0	0	0	0
4	C	52	0	0	0	0
4	D	41	0	0	0	0
All	All	4582	0	4057	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 15 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302:SER:O	1:A:305:ALA:HB2	1.95	0.67
1:A:276:GLN:HA	1:A:279:ARG:HH11	1.65	0.61
1:B:302:SER:O	1:B:305:ALA:HB2	2.02	0.58
1:C:325:PRO:HD2	1:D:274:LYS:HD3	1.87	0.56
1:C:268:ASP:O	1:C:269:ASP:HB2	2.09	0.53

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/146 (91%)	129 (97%)	4 (3%)	0	100	100
1	B	129/146 (88%)	126 (98%)	3 (2%)	0	100	100
1	C	126/146 (86%)	123 (98%)	3 (2%)	0	100	100
1	D	128/146 (88%)	125 (98%)	3 (2%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	516/584 (88%)	503 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/127 (92%)	113 (97%)	4 (3%)	37	51
1	B	114/127 (90%)	113 (99%)	1 (1%)	78	89
1	C	112/127 (88%)	108 (96%)	4 (4%)	35	49
1	D	113/127 (89%)	112 (99%)	1 (1%)	78	89
All	All	456/508 (90%)	446 (98%)	10 (2%)	52	69

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	243	TYR
1	C	243	TYR
1	C	291	ASN
1	A	327	GLU
1	C	261	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	E	1	2	12,12,12	1.72	5 (41%)	17,17,17	0.98	1 (5%)
2	GLC	E	2	3,2	11,11,12	1.58	2 (18%)	15,15,17	1.60	3 (20%)
2	GLC	F	1	2	12,12,12	1.69	4 (33%)	17,17,17	2.29	7 (41%)
2	GLC	F	2	3,2	11,11,12	0.83	0	15,15,17	0.91	0
2	GLC	G	1	2	12,12,12	1.47	1 (8%)	17,17,17	1.42	3 (17%)
2	GLC	G	2	3,2	11,11,12	1.00	1 (9%)	15,15,17	2.07	2 (13%)
2	GLC	H	1	2	12,12,12	1.64	3 (25%)	17,17,17	1.36	2 (11%)
2	GLC	H	2	3,2	11,11,12	1.05	1 (9%)	15,15,17	1.95	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	1	2	-	2/2/22/22	0/1/1/1
2	GLC	E	2	3,2	-	0/2/19/22	0/1/1/1
2	GLC	F	1	2	-	0/2/22/22	0/1/1/1
2	GLC	F	2	3,2	-	2/2/19/22	0/1/1/1
2	GLC	G	1	2	-	2/2/22/22	0/1/1/1
2	GLC	G	2	3,2	-	2/2/19/22	0/1/1/1
2	GLC	H	1	2	-	0/2/22/22	0/1/1/1
2	GLC	H	2	3,2	-	2/2/19/22	0/1/1/1

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	2	GLC	O5-C1	4.00	1.50	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	1	GLC	O5-C1	3.04	1.50	1.42
2	H	2	GLC	O5-C1	3.01	1.48	1.43
2	F	1	GLC	O4-C4	2.67	1.49	1.43
2	E	1	GLC	C4-C5	2.63	1.58	1.53

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	2	GLC	C1-O5-C5	6.43	120.90	112.19
2	F	1	GLC	C3-C4-C5	4.58	118.41	110.24
2	F	1	GLC	O5-C5-C4	4.42	117.71	109.69
2	H	2	GLC	C1-O5-C5	4.42	118.17	112.19
2	G	2	GLC	O5-C1-C2	4.05	117.02	110.77

There are no chirality outliers.

5 of 10 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	2	GLC	O5-C5-C6-O6
2	E	1	GLC	O5-C5-C6-O6
2	G	2	GLC	O5-C5-C6-O6
2	F	2	GLC	C4-C5-C6-O6
2	H	2	GLC	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	2	GLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	134/146 (91%)	-0.25	6 (4%) 33 40	12, 25, 61, 107	0
1	B	129/146 (88%)	-0.51	0 100 100	11, 23, 39, 57	0
1	C	128/146 (87%)	-0.46	2 (1%) 72 77	11, 22, 43, 67	0
1	D	129/146 (88%)	-0.28	3 (2%) 60 67	12, 27, 50, 68	0
All	All	520/584 (89%)	-0.37	11 (2%) 63 70	11, 25, 46, 107	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	327	GLU	3.8
1	C	291	ASN	3.7
1	A	328	PRO	3.7
1	A	332	SER	3.0
1	D	291	ASN	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	H	1	12/12	0.52	0.40	60,67,68,69	0
2	GLC	E	1	12/12	0.54	0.34	63,72,76,76	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GLC	G	1	12/12	0.65	0.49	59,76,79,79	0
2	GLC	F	1	12/12	0.70	0.45	54,65,68,70	0
2	GLC	E	2	11/12	0.88	0.21	42,52,57,62	0
2	GLC	H	2	11/12	0.89	0.17	40,47,53,55	0
2	GLC	G	2	11/12	0.92	0.12	35,42,47,49	0
2	GLC	F	2	11/12	0.93	0.13	24,35,41,43	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	D	1	1/1	0.96	0.08	33,33,33,33	0
3	CA	C	1	1/1	0.97	0.03	29,29,29,29	0
3	CA	A	1	1/1	0.99	0.03	26,26,26,26	0
3	CA	B	1	1/1	0.99	0.03	24,24,24,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.