



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 24, 2021 – 12:12 PM EST

PDB ID : 2P8E
Title : Crystal structure of the serine/threonine phosphatase domain of human PPM1B
Authors : Bonanno, J.B.; Freeman, J.; Bain, K.T.; Lau, C.; Xu, W.; Smith, D.; Wasserman, S.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2007-03-22
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.16
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	
1	B	307	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4871 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PPM1B beta isoform variant 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	278	Total	C	N	O	S	0	4	0
			2212	1376	387	432	17			
1	B	281	Total	C	N	O	S	0	3	0
			2226	1383	391	434	18			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP Q4J6C0
A	0	SER	-	expression tag	UNP Q4J6C0
A	1	LEU	-	expression tag	UNP Q4J6C0
A	298	GLU	-	expression tag	UNP Q4J6C0
A	299	GLY	-	expression tag	UNP Q4J6C0
A	300	HIS	-	expression tag	UNP Q4J6C0
A	301	HIS	-	expression tag	UNP Q4J6C0
A	302	HIS	-	expression tag	UNP Q4J6C0
A	303	HIS	-	expression tag	UNP Q4J6C0
A	304	HIS	-	expression tag	UNP Q4J6C0
A	305	HIS	-	expression tag	UNP Q4J6C0
B	-1	MET	-	expression tag	UNP Q4J6C0
B	0	SER	-	expression tag	UNP Q4J6C0
B	1	LEU	-	expression tag	UNP Q4J6C0
B	298	GLU	-	expression tag	UNP Q4J6C0
B	299	GLY	-	expression tag	UNP Q4J6C0
B	300	HIS	-	expression tag	UNP Q4J6C0
B	301	HIS	-	expression tag	UNP Q4J6C0
B	302	HIS	-	expression tag	UNP Q4J6C0
B	303	HIS	-	expression tag	UNP Q4J6C0
B	304	HIS	-	expression tag	UNP Q4J6C0
B	305	HIS	-	expression tag	UNP Q4J6C0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Mg 1	0	0
2	A	1	Total 1	Mg 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	218	Total 218	O 218	0	0
3	B	213	Total 213	O 213	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PPM1B beta isoform variant 6



- Molecule 1: PPM1B beta isoform variant 6



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.40Å 92.31Å 118.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.82 26.06 – 1.82	Depositor EDS
% Data completeness (in resolution range)	93.8 (20.00-1.82) 93.8 (26.06-1.82)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 1.82Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.186 , 0.246 0.189 , 0.247	Depositor DCC
R_{free} test set	2525 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.359	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 53.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4871	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OCS, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/2257	0.77	2/3047 (0.1%)
1	B	0.75	0/2269	0.75	2/3064 (0.1%)
All	All	0.75	0/4526	0.76	4/6111 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	118	ARG	NE-CZ-NH2	-8.24	116.18	120.30
1	A	5	LEU	CA-CB-CG	6.07	129.25	115.30
1	A	118	ARG	NE-CZ-NH2	-5.19	117.70	120.30
1	B	286	ASP	CB-CG-OD1	5.16	122.94	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2212	0	2117	24	0
1	B	2226	0	2130	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	218	0	0	5	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	213	0	0	8	1
All	All	4871	0	4247	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:GLU:HG3	3:A:463:HOH:O	1.71	0.90
1:B:277:ASP:O	1:B:281:HIS:ND1	2.12	0.82
1:A:233:GLU:H	1:A:233:GLU:CD	1.92	0.73
1:B:123:LEU:HA	1:B:127:MET:HG3	1.75	0.67
1:B:52[B]:ASP:OD1	3:B:467:HOH:O	2.13	0.67
1:A:256[A]:GLU:HG2	3:A:500:HOH:O	1.97	0.65
1:B:119:ASN:CB	3:B:316:HOH:O	2.48	0.61
1:B:77:GLU:HG3	3:B:475:HOH:O	2.01	0.60
1:B:119:ASN:HB3	3:B:316:HOH:O	2.04	0.56
1:B:146:PHE:CE1	1:B:229:ILE:HD12	2.41	0.56
1:A:228:GLU:HB3	3:A:321:HOH:O	2.07	0.54
1:B:78:HIS:HE1	3:B:365:HOH:O	1.91	0.53
1:A:169:HIS:HE2	1:A:225:GLU:CD	2.13	0.51
1:B:155[A]:VAL:HG23	1:B:162:VAL:HG13	1.94	0.49
1:A:120:PHE:HB2	1:A:123:LEU:HD23	1.94	0.49
1:B:188:MET:HE3	3:B:373:HOH:O	2.12	0.49
1:A:100:GLU:H	1:A:100:GLU:CD	2.16	0.49
1:A:66:ARG:HD2	3:A:334:HOH:O	2.13	0.48
1:A:150:GLY:HA2	1:A:198[B]:VAL:CG2	2.43	0.47
1:A:85:PHE:HE2	1:A:138:MET:HE1	1.79	0.46
1:B:77:GLU:CG	3:B:475:HOH:O	2.63	0.46
1:A:53:TRP:HZ3	1:A:138:MET:HE2	1.81	0.45
1:B:234:GLU:HG3	3:B:349:HOH:O	2.16	0.45
1:B:59:TYR:CE2	1:B:72:SER:HA	2.51	0.45
1:A:59:TYR:CE2	1:A:72:SER:HA	2.52	0.45
1:A:108:THR:O	1:A:112:LYS:HG3	2.17	0.45
1:A:112:LYS:HB3	1:A:112:LYS:HE3	1.66	0.44
1:B:153:ARG:HB3	1:B:246:TRP:NE1	2.33	0.44
1:A:157:TYR:CD1	1:A:255:CYS:HB3	2.52	0.44
1:A:60:ASP:HB3	1:A:133:THR:OG1	2.18	0.44
1:B:46:ILE:HB	1:B:50:LEU:HB2	2.00	0.44
1:A:66:ARG:NE	1:A:122:ASP:OD2	2.48	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:285:ARG:NH2	3:A:443:HOH:O	2.50	0.43
1:A:5:LEU:HD13	1:A:34:VAL:HA	2.01	0.43
1:A:172:CYS:SG	1:A:190:GLN:HG2	2.58	0.43
1:A:150:GLY:HA2	1:A:198[B]:VAL:HG21	2.02	0.42
1:B:234:GLU:CD	1:B:234:GLU:H	2.23	0.41
1:A:110:PHE:CD1	1:A:224:PRO:HG3	2.56	0.41
1:A:233:GLU:CD	1:A:233:GLU:N	2.67	0.41
1:A:169:HIS:CE1	1:A:225:GLU:HG3	2.55	0.41
1:B:118:ARG:NH2	1:B:209:CYS:SG	2.95	0.40
1:B:164:PHE:CG	1:B:229:ILE:HD11	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:454:HOH:O	3:B:512:HOH:O[3_655]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/307 (90%)	268 (98%)	7 (2%)	0	100	100
1	B	279/307 (91%)	273 (98%)	6 (2%)	0	100	100
All	All	554/614 (90%)	541 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/260 (94%)	239 (98%)	5 (2%)	55	43
1	B	245/260 (94%)	241 (98%)	4 (2%)	62	53
All	All	489/520 (94%)	480 (98%)	9 (2%)	57	48

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	112	LYS
1	A	129	ARG
1	A	138	MET
1	A	195	SER
1	B	6	ASP
1	B	129	ARG
1	B	186	SER
1	B	266	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	ASN
1	A	101	ASN
1	A	104	ASN
1	A	119	ASN
1	A	143	HIS
1	A	161	GLN
1	A	167	GLN
1	A	181	GLN
1	A	182	ASN
1	B	78	HIS
1	B	101	ASN
1	B	181	GLN
1	B	182	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OCS	A	242	1,2	7,8,9	1.23	0	6,11,13	1.25	1 (16%)
1	OCS	B	242	1,2	7,8,9	1.21	1 (14%)	6,11,13	1.74	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	A	242	1,2	-	3/4/7/9	-
1	OCS	B	242	1,2	-	0/4/7/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	242	OCS	CB-CA	-2.02	1.51	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	242	OCS	OD2-SG-CB	3.61	111.49	105.74
1	A	242	OCS	OD2-SG-CB	2.56	109.82	105.74

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	242	OCS	CA-CB-SG-OD2
1	A	242	OCS	CA-CB-SG-OD1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	242	OCS	CA-CB-SG-OD3

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	277/307 (90%)	0.08	9 (3%)	47	42	12, 20, 33, 40	0
1	B	280/307 (91%)	0.05	8 (2%)	51	46	11, 19, 31, 45	0
All	All	557/614 (90%)	0.06	17 (3%)	49	43	11, 19, 33, 45	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	123	LEU	4.9
1	A	4	PHE	4.2
1	A	189	ILE	3.4
1	A	86	ARG	3.2
1	A	83	GLU	2.8
1	B	86	ARG	2.7
1	B	6	ASP	2.7
1	A	100	GLU	2.4
1	A	120	PHE	2.2
1	B	4	PHE	2.2
1	B	7	LYS	2.2
1	A	135	VAL	2.2
1	A	281	HIS	2.1
1	B	120	PHE	2.1
1	B	126	GLY	2.1
1	A	58	VAL	2.0
1	B	48	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	OCS	A	242	9/10	0.98	0.11	16,18,23,25	0
1	OCS	B	242	9/10	0.98	0.08	15,17,20,21	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	A	306	1/1	0.98	0.11	16,16,16,16	0
2	MG	B	306	1/1	0.99	0.11	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.