



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 02:14 am BST

PDB ID : 2P91  
Title : Crystal structure of Enoyl-[acyl-carrier-protein] reductase (NADH) from Aquifex aeolicus VF5  
Authors : Chen, L.; Li, Y.; Ebihara, A.; Shinkai, A.; Kuramitsu, S.; Yokoyama, S.; Zhao, M.; Rose, J.P.; Wang, B.-C.; Southeast Collaboratory for Structural Genomics (SECSG); RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2007-03-23  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>5%</div> <div>11%</div> </div> </div>
1	B	285	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>6%</div> <div>12%</div> </div> </div>
1	C	285	<div> <div></div> <div> <div></div> <div>76%</div> <div>•</div> <div>20%</div> </div> </div>
1	D	285	<div> <div></div> <div> <div></div> <div>74%</div> <div>8%</div> <div>18%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enoyl-[acyl-carrier-protein] reductase [NADH].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	0	0
			1960	1259	334	360	7			
1	B	252	Total	C	N	O	S	0	0	0
			1941	1244	332	358	7			
1	C	228	Total	C	N	O	S	0	0	0
			1759	1126	300	327	6			
1	D	233	Total	C	N	O	S	0	0	0
			1796	1151	307	332	6			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	SEE REMARK 999	UNP O67505
A	2	LEU	-	SEE REMARK 999	UNP O67505
A	3	ARG	-	SEE REMARK 999	UNP O67505
A	4	LYS	-	SEE REMARK 999	UNP O67505
A	5	LEU	-	SEE REMARK 999	UNP O67505
A	6	SER	-	SEE REMARK 999	UNP O67505
A	7	LYS	-	SEE REMARK 999	UNP O67505
A	8	PHE	-	SEE REMARK 999	UNP O67505
A	9	SER	-	SEE REMARK 999	UNP O67505
A	10	ASN	-	SEE REMARK 999	UNP O67505
A	11	LYS	-	SEE REMARK 999	UNP O67505
A	12	GLY	-	SEE REMARK 999	UNP O67505
A	13	GLU	-	SEE REMARK 999	UNP O67505
A	14	VAL	-	SEE REMARK 999	UNP O67505
A	15	PHE	-	SEE REMARK 999	UNP O67505
B	1	MET	-	SEE REMARK 999	UNP O67505
B	2	LEU	-	SEE REMARK 999	UNP O67505
B	3	ARG	-	SEE REMARK 999	UNP O67505
B	4	LYS	-	SEE REMARK 999	UNP O67505
B	5	LEU	-	SEE REMARK 999	UNP O67505
B	6	SER	-	SEE REMARK 999	UNP O67505

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	7	LYS	-	SEE REMARK 999	UNP O67505
B	8	PHE	-	SEE REMARK 999	UNP O67505
B	9	SER	-	SEE REMARK 999	UNP O67505
B	10	ASN	-	SEE REMARK 999	UNP O67505
B	11	LYS	-	SEE REMARK 999	UNP O67505
B	12	GLY	-	SEE REMARK 999	UNP O67505
B	13	GLU	-	SEE REMARK 999	UNP O67505
B	14	VAL	-	SEE REMARK 999	UNP O67505
B	15	PHE	-	SEE REMARK 999	UNP O67505
C	1	MET	-	SEE REMARK 999	UNP O67505
C	2	LEU	-	SEE REMARK 999	UNP O67505
C	3	ARG	-	SEE REMARK 999	UNP O67505
C	4	LYS	-	SEE REMARK 999	UNP O67505
C	5	LEU	-	SEE REMARK 999	UNP O67505
C	6	SER	-	SEE REMARK 999	UNP O67505
C	7	LYS	-	SEE REMARK 999	UNP O67505
C	8	PHE	-	SEE REMARK 999	UNP O67505
C	9	SER	-	SEE REMARK 999	UNP O67505
C	10	ASN	-	SEE REMARK 999	UNP O67505
C	11	LYS	-	SEE REMARK 999	UNP O67505
C	12	GLY	-	SEE REMARK 999	UNP O67505
C	13	GLU	-	SEE REMARK 999	UNP O67505
C	14	VAL	-	SEE REMARK 999	UNP O67505
C	15	PHE	-	SEE REMARK 999	UNP O67505
D	1	MET	-	SEE REMARK 999	UNP O67505
D	2	LEU	-	SEE REMARK 999	UNP O67505
D	3	ARG	-	SEE REMARK 999	UNP O67505
D	4	LYS	-	SEE REMARK 999	UNP O67505
D	5	LEU	-	SEE REMARK 999	UNP O67505
D	6	SER	-	SEE REMARK 999	UNP O67505
D	7	LYS	-	SEE REMARK 999	UNP O67505
D	8	PHE	-	SEE REMARK 999	UNP O67505
D	9	SER	-	SEE REMARK 999	UNP O67505
D	10	ASN	-	SEE REMARK 999	UNP O67505
D	11	LYS	-	SEE REMARK 999	UNP O67505
D	12	GLY	-	SEE REMARK 999	UNP O67505
D	13	GLU	-	SEE REMARK 999	UNP O67505
D	14	VAL	-	SEE REMARK 999	UNP O67505
D	15	PHE	-	SEE REMARK 999	UNP O67505

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	153	Total 153	O 153	0	0
2	B	184	Total 184	O 184	0	0
2	C	150	Total 150	O 150	0	0
2	D	157	Total 157	O 157	0	0



- Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]



HIS	THR	THR	LYS	VAL	ASN	PRO	F231	G232	K233	T236	V245	G265	I268	V271	PHE	GLY	ARG	GLU	GLU	GLU	ILE	LYS	LYS	GLU	VAL	TYR	GLY	ASP
-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.08 Å   119.02 Å   119.20 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	30.00 – 2.00 23.38 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-2.00) 99.7 (23.38-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.89 (at 1.99 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.167 , 0.197 0.169 , 0.198	Depositor DCC
$R_{free}$ test set	3528 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.6	Xtriage
Anisotropy	0.385	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 35.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.467 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8100	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/1998	0.60	2/2700 (0.1%)
1	B	0.52	0/1978	0.56	0/2673
1	C	0.52	0/1791	0.55	1/2420 (0.0%)
1	D	0.49	0/1829	0.55	0/2469
All	All	0.51	0/7596	0.57	3/10262 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	186	ARG	NE-CZ-NH1	-8.16	116.22	120.30
1	A	186	ARG	NE-CZ-NH2	6.89	123.75	120.30
1	C	18	LEU	CA-CB-CG	5.94	128.96	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1960	0	1989	14	0
1	B	1941	0	1969	12	0
1	C	1759	0	1782	9	0
1	D	1796	0	1822	15	0
2	A	153	0	0	3	0
2	B	184	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	150	0	0	4	0
2	D	157	0	0	3	0
All	All	8100	0	7562	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:ILE:HG21	1:B:159:LEU:HD21	1.24	1.09
1:B:35:ILE:HG21	1:B:159:LEU:CD2	2.03	0.89
1:A:62:ARG:HD3	2:A:396:HOH:O	1.73	0.86
1:B:35:ILE:CG2	1:B:159:LEU:HD21	2.09	0.79
1:B:146:LEU:HA	1:B:149:MET:HE3	1.70	0.71
1:A:62:ARG:HG3	1:A:62:ARG:HH11	1.57	0.69
1:C:18:LEU:HD13	1:C:20:GLU:H	1.57	0.68
2:B:329:HOH:O	1:D:128:LYS:HG3	1.92	0.68
1:C:90:LYS:HE2	1:C:94:GLU:OE2	1.97	0.65
1:D:231:PHE:O	1:D:233:LYS:N	2.32	0.62
1:B:230:PRO:HD3	1:B:269:MET:HE2	1.83	0.60
1:A:62:ARG:CG	1:A:62:ARG:HH11	2.17	0.58
1:D:207:VAL:HG12	1:D:208:LYS:HG2	1.89	0.54
1:A:62:ARG:CD	2:A:396:HOH:O	2.45	0.53
1:B:124:ARG:HG2	1:B:128:LYS:HE3	1.92	0.52
1:A:31:ASN:HD21	1:A:210:LEU:HB2	1.75	0.52
1:C:18:LEU:HD22	1:C:46:GLU:O	2.12	0.49
1:D:64:ARG:NH2	2:D:419:HOH:O	2.45	0.49
1:B:230:PRO:HD3	1:B:269:MET:CE	2.43	0.48
1:A:224:HIS:CE1	1:B:271:VAL:HG12	2.49	0.47
1:A:128:LYS:HG3	2:C:308:HOH:O	2.14	0.47
1:C:46:GLU:HG3	1:C:245:VAL:HG11	1.97	0.47
1:C:18:LEU:HD11	1:C:20:GLU:HB2	1.97	0.46
1:B:81:SER:HB3	1:B:133:ILE:HD11	1.98	0.46
1:B:226:THR:O	1:B:232:GLY:HA2	2.17	0.45
1:C:234:PRO:N	2:C:401:HOH:O	2.48	0.45
1:D:208:LYS:HD2	1:D:236:THR:HA	1.98	0.45
1:A:81:SER:HB3	1:A:133:ILE:HD11	1.98	0.45
1:A:62:ARG:CG	1:A:62:ARG:NH1	2.80	0.44
1:A:226:THR:O	1:A:232:GLY:HA2	2.18	0.44
1:A:101:ASP:OD1	1:A:152:ARG:HD2	2.17	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:54:TYR:CE1	1:C:60:GLU:HB2	2.54	0.43
1:D:46:GLU:HG3	1:D:245:VAL:HG11	1.99	0.43
1:A:46:GLU:HG3	1:A:245:VAL:HG11	2.01	0.42
1:A:128:LYS:CG	2:C:308:HOH:O	2.66	0.42
1:D:35:ILE:HG21	1:D:159:LEU:HD11	2.02	0.42
1:D:146:LEU:HA	1:D:149:MET:HE3	2.00	0.42
1:B:32:GLU:CD	1:B:32:GLU:H	2.23	0.42
1:C:90:LYS:HE3	2:C:391:HOH:O	2.19	0.42
1:D:172:ASN:HB3	2:D:309:HOH:O	2.19	0.42
1:A:62:ARG:NE	2:A:396:HOH:O	2.53	0.41
1:C:171:TYR:CZ	1:C:174:MET:HG3	2.55	0.41
2:B:329:HOH:O	1:D:128:LYS:CG	2.61	0.41
1:D:265:GLY:O	1:D:268:ILE:HG12	2.21	0.41
1:B:17:GLY:N	1:B:46:GLU:OE1	2.54	0.40
1:D:87:LYS:HD3	1:D:144:GLU:OE2	2.21	0.40
1:D:207:VAL:CG1	1:D:208:LYS:HG2	2.51	0.40
1:D:160:SER:HA	1:D:178:LYS:HD2	2.03	0.40
1:D:23:ARG:HD2	2:D:348:HOH:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/285 (88%)	243 (97%)	7 (3%)	0	100	100
1	B	248/285 (87%)	242 (98%)	6 (2%)	0	100	100
1	C	224/285 (79%)	217 (97%)	7 (3%)	0	100	100
1	D	229/285 (80%)	219 (96%)	9 (4%)	1 (0%)	34	30
All	All	951/1140 (83%)	921 (97%)	29 (3%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	232	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/233 (88%)	203 (99%)	2 (1%)	76	81
1	B	203/233 (87%)	203 (100%)	0	100	100
1	C	184/233 (79%)	184 (100%)	0	100	100
1	D	187/233 (80%)	187 (100%)	0	100	100
All	All	779/932 (84%)	777 (100%)	2 (0%)	92	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	LYS
1	A	62	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	254/285 (89%)	0.07	6 (2%) 59 57	7, 15, 28, 35	0
1	B	252/285 (88%)	0.07	5 (1%) 65 63	7, 15, 27, 37	0
1	C	228/285 (80%)	-0.11	0 100 100	7, 17, 28, 34	0
1	D	233/285 (81%)	-0.05	0 100 100	7, 17, 29, 40	0
All	All	967/1140 (84%)	-0.00	11 (1%) 80 79	7, 16, 28, 40	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	210	LEU	5.2
1	A	210	LEU	3.6
1	B	211	ALA	3.0
1	B	98	GLY	3.0
1	A	215	ILE	2.7
1	A	271	VAL	2.5
1	A	62	ARG	2.3
1	A	211	ALA	2.2
1	B	62	ARG	2.2
1	B	17	GLY	2.1
1	A	217	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.