



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 03:59 am BST

PDB ID : 4PBE
Title : Phosphotriesterase Variant Rev6
Authors : Campbell, E.; Kaltenbach, M.; Tokuriki, N.; Jackson, C.J.
Deposited on : 2014-04-12
Resolution : 1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	<div> <div>8%</div> <div>91%</div> <div>8% ..</div> </div>
1	G	333	<div> <div>5%</div> <div>87%</div> <div>11% ..</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5712 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphotriesterase variant PTE-revR6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	0	6	0
			2582	1627	465	480	10			
1	G	331	Total	C	N	O	S	0	5	0
			2581	1627	463	480	11			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is CACODYLATE ION (three-letter code: CAC) (formula: C₂H₆AsO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	G	1	Total	C	O	0	0
			8	6	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	275	Total	O	0	0
			275	275		
5	G	257	Total	O	0	0
			257	257		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphotriesterase variant PTE-revR6



- Molecule 1: Phosphotriesterase variant PTE-revR6



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.64Å 85.97Å 89.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.59 – 1.51 42.98 – 1.51	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.59-1.51) 92.6 (42.98-1.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 1.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.210 , 0.249 0.211 , 0.249	Depositor DCC
R_{free} test set	5162 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	13.6	Xtriage
Anisotropy	0.347	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.017 for -h,l,k 0.018 for -l,-k,-h 0.019 for k,h,-l 0.007 for k,l,h 0.007 for l,h,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5712	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, ZN, MPD, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2615	0.56	0/3546
1	G	0.37	0/2615	0.54	0/3546
All	All	0.36	0/5230	0.55	0/7092

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2582	0	2604	20	0
1	G	2581	0	2599	22	0
2	A	2	0	0	0	0
2	G	2	0	0	0	0
3	A	5	0	0	0	0
4	G	8	0	14	0	0
5	A	275	0	0	3	0
5	G	257	0	0	2	0
All	All	5712	0	5217	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:167:ILE:HD12	1:G:197:PRO:HG2	1.69	0.75
1:A:67[B]:ARG:NH2	5:A:2775:HOH:O	2.23	0.70
1:G:344:GLU:O	5:G:2726:HOH:O	2.13	0.66
1:G:167:ILE:HD11	1:G:199:ILE:HG13	1.80	0.63
1:A:67[A]:ARG:NH1	5:A:2684:HOH:O	2.32	0.62
1:A:308:CYS:HB2	1:G:136:LEU:HD13	1.82	0.61
1:G:203:GLU:HG2	1:G:206:GLN:HG2	1.84	0.58
1:G:57:HIS:HB2	1:G:303:LEU:HB3	1.86	0.57
1:A:87:LEU:HD12	1:A:116[B]:VAL:HG12	1.88	0.55
1:G:172:THR:HG22	1:G:180:HIS:HB3	1.90	0.54
1:G:302:TRP:CH2	1:G:321:ASN:HB3	2.42	0.53
1:A:110[B]:VAL:HG23	5:A:2560:HOH:O	2.09	0.53
1:A:110[B]:VAL:HG21	1:A:158:ILE:HG21	1.91	0.52
1:A:203[A]:GLU:O	1:A:205:SER:N	2.43	0.51
1:G:337:ARG:HD3	1:G:346:LEU:HD12	1.93	0.50
1:A:302:TRP:CH2	1:A:321:ASN:HB3	2.46	0.49
1:G:57:HIS:O	1:G:303:LEU:HA	2.11	0.49
1:G:333:ILE:HG23	1:G:346:LEU:HD13	1.95	0.49
1:A:57:HIS:HB2	1:A:303:LEU:HB3	1.96	0.48
1:G:91:ARG:NH2	1:G:121:ASP:OD2	2.46	0.48
1:A:363:ARG:HG2	1:A:364:ALA:HA	1.96	0.47
1:A:57:HIS:O	1:A:303:LEU:HA	2.15	0.46
1:A:363:ARG:HB3	1:A:363:ARG:CZ	2.44	0.46
1:A:260:ILE:HD13	1:A:260:ILE:HA	1.69	0.45
1:G:176:MET:HG3	1:G:180:HIS:HB2	1.98	0.45
1:A:244:ALA:O	1:A:295[B]:GLN:NE2	2.30	0.44
1:G:353:ASN:HB2	1:G:354:PRO:HD3	1.98	0.44
1:G:337:ARG:HD2	1:G:341:THR:O	2.18	0.43
1:G:48:GLU:OE2	5:G:2716:HOH:O	2.21	0.43
1:G:98:ILE:HG21	1:G:325[B]:MET:HE1	2.00	0.43
1:G:339:LYS:HA	1:G:339:LYS:HD3	1.77	0.42
1:A:136:LEU:HD13	1:G:308:CYS:HB2	2.02	0.41
1:A:333:ILE:HB	1:A:334:PRO:HD3	2.01	0.41
1:A:353:ASN:HB2	1:A:354:PRO:HD3	2.03	0.41
1:G:300:ASN:OD1	1:G:328:ILE:HG12	2.21	0.41
1:A:363:ARG:HE	1:A:364:ALA:HB2	1.86	0.41
1:G:60:GLY:O	1:G:306:ILE:HA	2.20	0.41
1:A:131:TRP:CG	1:A:132:LEU:N	2.89	0.40
1:G:135:SER:O	1:G:139:ARG:HG3	2.22	0.40
1:A:181:GLU:HG3	1:A:215:ILE:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/333 (100%)	323 (97%)	11 (3%)	0	100	100
1	G	333/333 (100%)	323 (97%)	10 (3%)	0	100	100
All	All	667/666 (100%)	646 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/268 (101%)	264 (97%)	7 (3%)	46	16
1	G	271/268 (101%)	260 (96%)	11 (4%)	30	6
All	All	542/536 (101%)	524 (97%)	18 (3%)	39	10

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ASP
1	A	51	PHE
1	A	136	LEU
1	A	260	ILE
1	A	299	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	303	LEU
1	A	363	ARG
1	G	51	PHE
1	G	77[A]	GLU
1	G	77[B]	GLU
1	G	136	LEU
1	G	167	ILE
1	G	211	GLN
1	G	215	ILE
1	G	299	SER
1	G	303	LEU
1	G	362	LEU
1	G	363	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	KCX	G	169	1,2	7,11,12	1.17	1 (14%)	4,12,14	0.70	0
1	KCX	A	169	1,2	7,11,12	0.95	1 (14%)	4,12,14	1.19	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	G	169	1,2	-	0/7/10/12	-
1	KCX	A	169	1,2	-	0/7/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	169	KCX	CE-NZ	2.62	1.51	1.45
1	A	169	KCX	CE-NZ	2.09	1.50	1.45

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	169	KCX	CE-NZ-CX	-2.18	119.26	122.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CAC	A	2403	2	0,4,4	0.00	-	0,6,6	0.00	-
4	MPD	G	2403	-	7,7,7	0.53	0	9,10,10	0.77	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	G	2403	-	-	2/5/5/5	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	2403	MPD	O2-C2-C3	2.02	117.40	109.80

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	2403	MPD	C1-C2-C3-C4
4	G	2403	MPD	C2-C3-C4-O4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/333 (99%)	0.51	26 (7%) 12 13	9, 16, 64, 98	0
1	G	330/333 (99%)	0.45	16 (4%) 30 33	11, 21, 39, 102	0
All	All	660/666 (99%)	0.48	42 (6%) 19 20	9, 19, 45, 102	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	262	LEU	19.2
1	G	363	ARG	12.6
1	A	271	LEU	12.1
1	G	364	ALA	11.6
1	A	265	ASN	11.2
1	A	260	ILE	10.6
1	A	364	ALA	9.6
1	A	266	ALA	9.4
1	A	261	GLY	9.2
1	G	362	LEU	8.0
1	A	34	GLY	7.1
1	A	363	ARG	6.9
1	A	273	GLY	6.4
1	A	270	ALA	5.0
1	A	264	ASP	5.0
1	A	204[A]	GLY	4.9
1	A	274	SER	4.9
1	A	362	LEU	4.7
1	G	35	ASP	4.7
1	G	206	GLN	4.3
1	A	203[A]	GLU	4.1
1	A	269	THR	3.6
1	A	259	ALA	3.5
1	G	361	THR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	272	MET	3.3
1	A	268	ALA	3.3
1	A	205	SER	3.2
1	A	35	ASP	3.1
1	G	345	THR	3.1
1	A	263	GLU	3.1
1	G	34	GLY	3.1
1	G	156[A]	TYR	3.0
1	G	334	PRO	2.8
1	G	203	GLU	2.4
1	G	262	LEU	2.3
1	A	258	SER	2.3
1	A	175	LYS	2.2
1	G	292	TYR	2.2
1	G	205	SER	2.1
1	G	330	LEU	2.1
1	A	267	SER	2.0
1	G	175	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	G	169	12/13	0.90	0.10	12,15,20,20	0
1	KCX	A	169	12/13	0.94	0.09	8,10,13,14	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MPD	G	2403	8/8	0.81	0.19	15,27,31,34	0
3	CAC	A	2403	5/5	0.93	0.14	10,12,21,26	5
2	ZN	G	2402	1/1	0.97	0.07	19,19,19,19	1
2	ZN	A	2402	1/1	0.98	0.05	17,17,17,17	0
2	ZN	G	2401	1/1	0.99	0.05	15,15,15,15	0
2	ZN	A	2401	1/1	0.99	0.06	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.