



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 07:06 PM BST

PDB ID : 6PBJ
Title : The structure of 3-deoxy-d-arabino-heptulosonate 7-phosphate synthase with Gly190Pro mutation
Authors : Jiao, W.; Fan, Y.; Blackmore, N.J.; Parker, E.J.
Deposited on : 2019-06-13
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>9%</div> <div></div> </div> </div>
1	B	464	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>7%</div> <div>10%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PEG	B	509	-	-	X	-
6	PEG	B	510	-	-	X	-
7	PG4	A	510	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7214 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phospho-2-dehydro-3-deoxyheptonate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	9	0
			3411	2140	617	636	18			
1	B	416	Total	C	N	O	S	0	4	0
			3185	2005	578	586	16			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP A0A0T9FW33
A	0	ALA	-	expression tag	UNP A0A0T9FW33
A	190	PRO	GLY	engineered mutation	UNP A0A0T9FW33
B	-1	GLY	-	expression tag	UNP A0A0T9FW33
B	0	ALA	-	expression tag	UNP A0A0T9FW33
B	190	PRO	GLY	engineered mutation	UNP A0A0T9FW33

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mn	0	0
			1	1		
4	A	1	Total	Mn	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	3	Total	Cl	0	0
			3	3		
5	A	3	Total	Cl	0	0
			3	3		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 7 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			13	8	5		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	240	Total	O	0	0
			240	240		
8	B	289	Total	O	0	0
			289	289		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phospho-2-dehydro-3-deoxyheptonate aldolase



- Molecule 1: Phospho-2-dehydro-3-deoxyheptonate aldolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	204.25Å 204.25Å 66.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.83 – 1.90 36.83 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (36.83-1.90) 99.9 (36.83-1.90)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.158 , 0.180 0.159 , 0.178	Depositor DCC
R_{free} test set	6234 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.004 for -h,-k,l	Xtriage
Reported twinning fraction	0.611 for h,-h-k,-l 0.389 for h+k,-k,-l	Depositor
Outliers	0 of 124853 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7214	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PO4, MN, PG4, SO4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/3478	0.77	0/4741
1	B	0.77	0/3251	0.77	0/4429
All	All	0.77	0/6729	0.77	0/9170

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3411	0	3336	47	0
1	B	3185	0	3117	19	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	15	0	0	1	0
3	B	15	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	3	0	0	1	0
5	B	3	0	0	0	0
6	A	7	0	10	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	21	0	30	10	0
7	A	13	0	18	8	0
8	A	240	0	0	14	0
8	B	289	0	0	5	0
All	All	7214	0	6511	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:509:PEG:H42	8:A:678:HOH:O	1.68	0.92
1:B:96:GLU:OE1	6:B:510:PEG:H32	1.72	0.88
1:A:171:ARG:HH21	7:A:510:PG4:H82	1.36	0.88
1:A:5:VAL:HG13	1:B:9:ILE:HD11	1.54	0.87
1:A:100[B]:ARG:HG2	1:A:100[B]:ARG:HH21	1.46	0.81
1:A:171:ARG:HH21	7:A:510:PG4:C8	1.92	0.81
6:A:509:PEG:H12	8:A:805:HOH:O	1.83	0.78
1:A:100[B]:ARG:HH22	1:A:191:LEU:HD13	1.48	0.78
1:B:35:ALA:HA	6:B:509:PEG:H12	1.71	0.73
1:A:135:ARG:NH2	1:A:137:ALA:H	1.86	0.72
1:A:171:ARG:HE	7:A:510:PG4:C8	2.03	0.71
1:A:23:ARG:NH2	8:A:601:HOH:O	2.07	0.68
6:B:509:PEG:H12	6:B:509:PEG:H42	1.75	0.67
1:A:171:ARG:NH2	7:A:510:PG4:H82	2.10	0.67
1:A:237:ASN:CB	8:A:728:HOH:O	2.43	0.66
1:B:96:GLU:OE1	6:B:510:PEG:C3	2.41	0.65
1:A:111[B]:VAL:HG23	8:A:659:HOH:O	1.96	0.65
1:A:210:ALA:H	1:A:424:ALA:HB1	1.62	0.65
1:A:415:GLU:HB2	1:A:417:VAL:HG13	1.79	0.64
1:A:148[B]:ARG:NH1	8:A:604:HOH:O	2.30	0.64
1:A:284:ARG:HD3	8:A:741:HOH:O	1.97	0.63
1:B:445:ASN:H	1:B:448:GLN:HE21	1.47	0.62
1:A:100[B]:ARG:CG	1:A:100[B]:ARG:HH21	2.13	0.62
1:A:267:GLY:N	8:A:606:HOH:O	2.34	0.61
1:A:23:ARG:NH1	8:A:605:HOH:O	2.34	0.60
1:B:83:GLN:HA	1:B:124:VAL:O	2.02	0.59
1:B:220:GLU:CD	1:B:461:ARG:HH12	2.06	0.58
6:B:509:PEG:C1	6:B:509:PEG:H42	2.33	0.58
1:A:100[B]:ARG:NH2	1:A:191:LEU:HD13	2.18	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:ARG:HD3	8:B:858:HOH:O	2.06	0.55
1:B:77:GLY:HA2	1:B:404[B]:HIS:CD2	2.42	0.54
1:B:69:GLU:OE2	8:B:601:HOH:O	2.19	0.53
1:A:210:ALA:HB3	1:A:424:ALA:HB2	1.89	0.53
1:A:108:GLN:HG2	1:A:200:TRP:CE3	2.44	0.53
1:A:171:ARG:NH2	7:A:510:PG4:C8	2.69	0.51
1:A:73:GLN:NE2	8:A:614:HOH:O	2.44	0.51
1:A:341:HIS:HB2	5:A:506:CL:CL	2.48	0.51
1:A:135:ARG:HB2	1:A:148[A]:ARG:HD3	1.92	0.50
1:A:100[B]:ARG:NH2	1:A:100[B]:ARG:CG	2.75	0.50
1:B:369:HIS:CE1	1:B:411:GLU:OE2	2.65	0.49
1:A:317:VAL:HG21	1:A:353:LYS:HG2	1.95	0.49
1:B:220:GLU:OE1	1:B:461:ARG:NH1	2.46	0.49
1:A:83:GLN:HA	1:A:124:VAL:O	2.13	0.48
1:A:96[B]:GLU:OE2	1:A:100[B]:ARG:HD2	2.13	0.48
1:A:100[B]:ARG:NH2	8:A:619:HOH:O	2.47	0.47
1:A:108:GLN:O	1:A:111[A]:VAL:HG12	2.15	0.47
1:B:204:PHE:O	1:B:208:SER:HB2	2.14	0.47
1:A:171:ARG:HH21	7:A:510:PG4:H81	1.79	0.47
6:B:511:PEG:H11	8:B:731:HOH:O	2.15	0.46
1:A:100[B]:ARG:NH2	1:A:100[B]:ARG:HG2	2.24	0.46
1:B:445:ASN:H	1:B:448:GLN:NE2	2.11	0.46
1:A:260:ARG:HD2	6:B:510:PEG:H42	1.98	0.46
1:A:263:ASP:C	1:A:270:GLN:NE2	2.70	0.45
1:A:148[A]:ARG:NH2	8:A:604:HOH:O	2.34	0.45
1:A:108:GLN:HG2	1:A:200:TRP:HE3	1.81	0.45
1:B:100:ARG:HD3	8:B:777:HOH:O	2.17	0.45
1:A:12:LEU:HB3	1:A:13:PRO:HD2	1.99	0.45
1:B:32:LYS:HE3	8:B:828:HOH:O	2.16	0.45
1:A:96[B]:GLU:HB3	1:A:97:PRO:HD3	2.00	0.44
3:A:504:SO4:O1	8:A:602:HOH:O	2.21	0.43
1:A:135:ARG:HH21	1:A:137:ALA:H	1.65	0.42
6:B:510:PEG:C3	6:B:510:PEG:O1	2.67	0.42
1:A:11:GLN:HG2	1:B:3:TRP:CZ3	2.54	0.42
1:A:263:ASP:C	1:A:270:GLN:HE21	2.22	0.42
1:A:294:ALA:HA	1:A:297:GLN:NE2	2.35	0.42
1:A:77:GLY:HA2	1:A:404[A]:HIS:CE1	2.54	0.42
7:A:510:PG4:H62	7:A:510:PG4:O5	2.19	0.42
6:B:509:PEG:C4	6:B:509:PEG:C1	2.94	0.42
6:B:511:PEG:H32	6:B:511:PEG:H11	1.87	0.41
1:A:171:ARG:NE	7:A:510:PG4:C8	2.78	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16:PRO:HA	1:B:17:PRO:HD3	1.88	0.41
6:A:509:PEG:C4	8:A:678:HOH:O	2.48	0.41
1:A:100[B]:ARG:HH22	1:A:191:LEU:CD1	2.23	0.40
1:A:381:THR:HA	1:A:442:PRO:HG3	2.03	0.40
1:B:84:GLY:HA2	1:B:410:VAL:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/464 (96%)	438 (98%)	6 (1%)	1 (0%)	47	38
1	B	410/464 (88%)	405 (99%)	4 (1%)	1 (0%)	47	38
All	All	855/928 (92%)	843 (99%)	10 (1%)	2 (0%)	47	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	440	CYS
1	B	129	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	350/377 (93%)	344 (98%)	6 (2%)	60	57
1	B	326/377 (86%)	319 (98%)	7 (2%)	53	48
All	All	676/754 (90%)	663 (98%)	13 (2%)	55	53

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	TRP
1	A	106	LEU
1	A	135	ARG
1	A	283	GLU
1	A	337	ARG
1	A	461	ARG
1	B	2	ASN
1	B	40	TRP
1	B	106	LEU
1	B	240	THR
1	B	298	VAL
1	B	337	ARG
1	B	450	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	GLN
1	A	270	GLN
1	A	297	GLN
1	A	447	GLN
1	B	447	GLN
1	B	448	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 8 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PO4	B	501	-	4,4,4	1.92	1 (25%)	6,6,6	0.55	0
3	SO4	B	504	-	4,4,4	0.37	0	6,6,6	0.12	0
6	PEG	B	511	-	6,6,6	0.29	0	5,5,5	0.27	0
3	SO4	A	502	-	4,4,4	0.48	0	6,6,6	0.20	0
2	PO4	A	501	-	4,4,4	1.11	0	6,6,6	0.48	0
3	SO4	B	502	-	4,4,4	0.42	0	6,6,6	0.28	0
6	PEG	B	510	-	6,6,6	0.22	0	5,5,5	0.16	0
6	PEG	B	509	-	6,6,6	0.38	0	5,5,5	0.14	0
3	SO4	A	504	-	4,4,4	0.55	0	6,6,6	0.12	0
3	SO4	A	503	-	4,4,4	0.33	0	6,6,6	0.05	0
3	SO4	B	503	-	4,4,4	0.45	0	6,6,6	0.15	0
7	PG4	A	510	-	12,12,12	0.45	0	11,11,11	0.25	0
6	PEG	A	509	-	6,6,6	0.47	0	5,5,5	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	B	511	-	-	2/4/4/4	-
6	PEG	B	509	-	-	1/4/4/4	-
6	PEG	B	510	-	-	3/4/4/4	-
6	PEG	A	509	-	-	2/4/4/4	-
7	PG4	A	510	-	-	3/10/10/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	PO4	P-O2	-2.65	1.46	1.54

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	510	PG4	O3-C5-C6-O4
6	B	510	PEG	C1-C2-O2-C3
6	B	510	PEG	O1-C1-C2-O2
6	A	509	PEG	C4-C3-O2-C2
7	A	510	PG4	O2-C3-C4-O3
6	A	509	PEG	O1-C1-C2-O2
6	B	510	PEG	C4-C3-O2-C2
7	A	510	PG4	C4-C3-O2-C2
6	B	511	PEG	O2-C3-C4-O4
6	B	509	PEG	C1-C2-O2-C3
6	B	511	PEG	C1-C2-O2-C3

There are no ring outliers.

6 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	511	PEG	2	0
6	B	510	PEG	4	0
6	B	509	PEG	4	0
3	A	504	SO4	1	0
7	A	510	PG4	8	0
6	A	509	PEG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/464 (95%)	-0.03	23 (5%) 27 30	8, 22, 48, 75	0
1	B	416/464 (89%)	-0.33	10 (2%) 59 62	9, 17, 43, 72	0
All	All	860/928 (92%)	-0.18	33 (3%) 40 43	8, 19, 45, 75	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	426	ASP	5.2
1	A	209	PRO	4.6
1	A	381	THR	4.5
1	A	418	THR	4.4
1	A	428	SER	4.0
1	A	416	ASN	3.6
1	A	424	ALA	3.6
1	A	417	VAL	3.6
1	A	421	LEU	3.5
1	A	425	GLN	3.5
1	A	207	THR	3.5
1	B	209	PRO	3.4
1	A	423	GLY	3.3
1	A	440	CYS	3.1
1	B	267	GLY	3.0
1	A	234	ALA	3.0
1	A	427	ILE	2.9
1	B	210	ALA	2.8
1	A	210	ALA	2.8
1	A	438	THR	2.8
1	B	207	THR	2.8
1	B	237	ASN	2.8
1	A	136	SER	2.7
1	A	341	HIS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	422	GLY	2.5
1	A	199	ASP	2.4
1	B	212	ALA	2.4
1	B	266	ASP	2.2
1	A	212	ALA	2.2
1	B	371	ASN	2.2
1	B	155	PHE	2.2
1	B	236	ARG	2.1
1	A	420	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CL	B	507	1/1	0.83	0.09	58,58,58,58	0
5	CL	A	507	1/1	0.85	0.06	61,61,61,61	0
6	PEG	B	509	7/7	0.87	0.16	20,31,33,33	7
5	CL	B	506	1/1	0.88	0.06	55,55,55,55	0
3	SO4	A	503	5/5	0.92	0.19	54,56,61,65	0
6	PEG	B	511	7/7	0.92	0.11	21,23,25,27	0
6	PEG	B	510	7/7	0.92	0.15	18,22,27,35	7
5	CL	A	506	1/1	0.93	0.11	54,54,54,54	0
7	PG4	A	510	13/13	0.93	0.13	22,26,28,28	0
3	SO4	B	503	5/5	0.94	0.15	11,20,23,28	5
3	SO4	A	502	5/5	0.94	0.15	9,12,13,15	5
5	CL	A	508	1/1	0.94	0.05	38,38,38,38	0
3	SO4	B	504	5/5	0.96	0.18	38,50,59,67	0
6	PEG	A	509	7/7	0.96	0.14	18,22,25,25	7

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	504	5/5	0.97	0.11	16,20,21,22	5
4	MN	A	505	1/1	0.97	0.08	39,39,39,39	1
5	CL	B	508	1/1	0.97	0.06	49,49,49,49	0
3	SO4	B	502	5/5	0.98	0.08	24,29,29,31	0
4	MN	B	505	1/1	0.98	0.15	33,33,33,33	1
2	PO4	B	501	5/5	0.99	0.06	19,19,24,26	0
2	PO4	A	501	5/5	0.99	0.07	26,26,30,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.