



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 06:29 am BST

PDB ID : 6PEU  
Title : Structure of YcaO enzyme from *Methanocaldococcus jannaschii* in complex with peptide  
Authors : Dong, S.-H.; Nair, S.K.  
Deposited on : 2019-06-20  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	<div> <div>8%</div> <div>93%</div> <div>6%</div> </div>
1	B	385	<div> <div>8%</div> <div>92%</div> <div>8%</div> </div>
1	C	385	<div> <div>8%</div> <div>89%</div> <div>9%</div> </div>
1	D	385	<div> <div>2%</div> <div>86%</div> <div>11%</div> </div>
2	M	12	<div> <div>83%</div> <div>17%</div> </div>
2	N	12	<div> <div>8%</div> <div>83%</div> <div>8%</div> <div>8%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	P	12	<div><div></div><div>17%</div><div>83%</div><div>8%</div><div>8%</div></div>
2	Q	12	<div><div></div><div>8%</div><div>83%</div><div>8%</div><div>8%</div></div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Uncharacterized protein MJ1094.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	S	0	0	0
			3100	1981	535	575	9			
1	B	385	Total	C	N	O	S	0	0	0
			3109	1988	536	577	8			
1	C	383	Total	C	N	O	S	0	0	0
			3092	1976	534	574	8			
1	D	380	Total	C	N	O	S	0	0	0
			3063	1957	529	569	8			

- Molecule 2 is a protein called GLY-ARG-LEU-GLY-PHE-TYR-GLY-TYR-ASP-LEU-GLN-ASP.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	M	12	Total	C	N	O	0	0	0
			99	64	16	19			
2	N	12	Total	C	N	O	0	0	0
			99	64	16	19			
2	P	12	Total	C	N	O	0	0	0
			99	64	16	19			
2	Q	12	Total	C	N	O	0	0	0
			99	64	16	19			

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Mg	0	0
			2	2		
4	A	2	Total	Mg	0	0
			2	2		
4	D	2	Total	Mg	0	0
			2	2		
4	C	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	217	Total	O	0	0
			217	217		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	250	Total 250	O 250	0	0
5	C	238	Total 238	O 238	0	0
5	D	210	Total 210	O 210	0	0
5	M	11	Total 11	O 11	0	0
5	N	12	Total 12	O 12	0	0
5	P	12	Total 12	O 12	0	0
5	Q	12	Total 12	O 12	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Uncharacterized protein MJ1094





#### • Molecule 1: Uncharacterized protein MJ1094




#### • Molecule 1: Uncharacterized protein MJ1094




#### • Molecule 1: Uncharacterized protein MJ1094





- Molecule 2: GLY-ARG-LEU-GLY-PHE-TYR-GLY-TYR-ASP-LEU-GLN-ASP

Chain M:  83% 17%





- Molecule 2: GLY-ARG-LEU-GLY-PHE-TYR-GLY-TYR-ASP-LEU-GLN-ASP

Chain N:  8% 83% 8% 8%





- Molecule 2: GLY-ARG-LEU-GLY-PHE-TYR-GLY-TYR-ASP-LEU-GLN-ASP

Chain P:  17% 83% 8% 8%



- Molecule 2: GLY-ARG-LEU-GLY-PHE-TYR-GLY-TYR-ASP-LEU-GLN-ASP

Chain Q:  8% 83% 8% 8%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.32Å 149.75Å 105.67Å 90.00° 103.57° 90.00°	Depositor
Resolution (Å)	25.00 – 1.95 102.72 – 1.95	Depositor EDS
% Data completeness (in resolution range)	95.8 (25.00-1.95) 95.9 (102.72-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.195 , 0.237 0.197 , 0.240	Depositor DCC
$R_{free}$ test set	6298 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.6	Xtriage
Anisotropy	0.459	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 43.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	13853	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/3156	0.66	1/4256 (0.0%)
1	B	0.46	0/3166	0.69	4/4271 (0.1%)
1	C	0.45	0/3148	0.67	2/4246 (0.0%)
1	D	0.44	0/3119	0.68	0/4209
2	M	0.56	0/101	0.75	0/134
2	N	0.66	0/101	0.74	0/134
2	P	0.63	0/101	0.76	0/134
2	Q	0.57	0/101	0.80	0/134
All	All	0.46	0/12993	0.68	7/17518 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	50	LEU	CA-CB-CG	7.87	133.39	115.30
1	B	50	LEU	CB-CG-CD1	6.18	121.50	111.00
1	B	86	ARG	NE-CZ-NH1	-6.14	117.23	120.30
1	C	342	LEU	CA-CB-CG	5.81	128.67	115.30
1	C	356	ARG	NE-CZ-NH2	-5.69	117.45	120.30
1	A	278	ARG	NE-CZ-NH2	-5.27	117.67	120.30
1	B	86	ARG	NE-CZ-NH2	5.26	122.93	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	6	LYS	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3100	0	3179	16	0
1	B	3109	0	3181	20	0
1	C	3092	0	3167	23	0
1	D	3063	0	3128	33	0
2	M	99	0	86	1	0
2	N	99	0	86	1	0
2	P	99	0	86	1	0
2	Q	99	0	86	1	0
3	A	31	0	12	0	0
3	B	31	0	12	2	0
3	C	31	0	12	0	0
3	D	31	0	12	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	1	0	0	0	0
4	D	2	0	0	0	0
5	A	217	0	0	3	0
5	B	250	0	0	1	0
5	C	238	0	0	3	0
5	D	210	0	0	1	0
5	M	11	0	0	0	0
5	N	12	0	0	0	0
5	P	12	0	0	0	0
5	Q	12	0	0	0	0
All	All	13853	0	13047	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:22:ILE:HG12	1:D:81:MET:HE1	1.41	1.02
1:C:145:THR:HG21	5:C:640:HOH:O	1.69	0.91
1:D:52:ARG:HE	1:D:65:HIS:HE1	1.15	0.88
1:B:145:THR:HG21	5:B:602:HOH:O	1.73	0.87
1:D:52:ARG:HE	1:D:65:HIS:CE1	2.00	0.79
1:D:81:MET:SD	5:D:747:HOH:O	2.40	0.79
1:D:56:VAL:N	1:D:57:ASP:HB2	2.01	0.75
1:D:252:HIS:HD2	1:D:254:HIS:H	1.36	0.73
1:A:252:HIS:HD2	1:A:254:HIS:H	1.37	0.73
1:B:252:HIS:HD2	1:B:254:HIS:H	1.37	0.72
1:C:36:ASN:OD1	1:C:38:GLN:HB2	1.90	0.72
1:B:106:VAL:HG12	1:B:111:LEU:HD11	1.72	0.71
1:D:56:VAL:H	1:D:57:ASP:HB2	1.57	0.69
1:A:66:TYR:OH	1:A:275:HIS:HD2	1.76	0.68
1:A:36:ASN:HD21	1:A:38:GLN:HE21	1.39	0.68
1:C:153:ASN:HD22	1:C:155:ASN:H	1.41	0.67
1:C:128:ILE:HG23	1:C:331:LYS:HD3	1.74	0.67
1:B:145:THR:HG23	1:B:147:GLY:H	1.59	0.66
1:C:254:HIS:HD2	1:C:256:GLU:H	1.43	0.66
1:B:74:GLN:HE22	3:B:500:ATP:HN61	1.44	0.66
1:A:254:HIS:HD2	1:A:256:GLU:H	1.44	0.64
1:C:252:HIS:HD2	1:C:254:HIS:H	1.44	0.64
1:C:257:ILE:O	1:C:261:ARG:HG3	2.01	0.61
1:D:254:HIS:HD2	1:D:256:GLU:H	1.50	0.59
1:A:71:ASN:C	1:A:71:ASN:HD22	2.05	0.59
1:B:284:ARG:NH1	2:N:443:ARG:O	2.29	0.59
1:B:273:GLN:NE2	1:B:284:ARG:HD3	2.17	0.59
1:B:169:HIS:HE1	3:B:500:ATP:O1B	1.85	0.58
1:A:83:ALA:HB3	5:A:602:HOH:O	2.03	0.58
1:D:81:MET:CE	1:D:84:ILE:HD12	2.34	0.57
1:C:270:ARG:NH1	5:C:601:HOH:O	2.29	0.57
1:B:254:HIS:HD2	1:B:256:GLU:H	1.52	0.57
1:B:145:THR:HG23	1:B:147:GLY:N	2.20	0.56
1:A:151:ARG:HD2	2:M:452:GLN:O	2.05	0.56
1:D:362:MET:HE2	1:D:362:MET:HA	1.89	0.55
1:D:81:MET:HE1	1:D:84:ILE:HD12	1.90	0.54
1:C:273:GLN:NE2	1:C:284:ARG:HE	2.06	0.54
1:B:208:GLU:O	1:B:212:LYS:HG2	2.07	0.54
1:B:128:ILE:HG23	1:B:331:LYS:HD3	1.91	0.53
1:A:68:LYS:HE3	5:A:746:HOH:O	2.09	0.53
1:B:185:LEU:HD11	1:B:372:LEU:HD11	1.90	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:66:TYR:OH	1:D:275:HIS:HD2	1.92	0.52
1:D:22:ILE:HG12	1:D:81:MET:CE	2.26	0.52
1:C:169:HIS:HE1	1:C:261:ARG:NH2	2.08	0.52
1:D:25:ALA:O	1:D:29:ILE:HG12	2.09	0.52
1:B:194:ASN:HB3	1:B:196:GLU:OE1	2.10	0.51
1:D:167:ILE:HD13	1:D:355:VAL:HG21	1.93	0.51
1:C:141:VAL:HA	1:C:356:ARG:HD3	1.93	0.51
1:D:93:LYS:O	1:D:96:VAL:HG22	2.11	0.50
1:C:66:TYR:OH	1:C:275:HIS:HD2	1.93	0.50
1:A:185:LEU:HD11	1:A:372:LEU:HD11	1.94	0.49
1:D:6:LYS:H	1:D:7:LEU:CB	2.25	0.49
1:D:194:ASN:OD1	1:D:196:GLU:HG2	2.12	0.49
1:D:252:HIS:CD2	1:D:254:HIS:H	2.23	0.48
1:C:156:GLY:HA3	1:C:174:ILE:HD11	1.95	0.47
1:D:298:ARG:O	1:D:301:ARG:HB2	2.14	0.47
1:D:6:LYS:H	1:D:7:LEU:HB2	1.79	0.47
1:D:300:HIS:CD2	2:Q:444:LEU:HD13	2.49	0.47
1:D:326:ASP:O	1:D:330:ILE:HG12	2.14	0.46
1:C:141:VAL:O	1:C:356:ARG:HD3	2.15	0.46
1:D:56:VAL:CA	1:D:57:ASP:HB2	2.46	0.46
1:D:254:HIS:HB3	1:D:257:ILE:HG12	1.98	0.45
1:B:5:TYR:HD2	1:B:9:SER:HB2	1.80	0.45
1:D:320:ARG:HD2	1:D:325:LYS:HB3	1.98	0.45
1:A:106:VAL:HG13	1:A:111:LEU:HD11	1.99	0.45
1:A:36:ASN:HD21	1:A:38:GLN:NE2	2.12	0.45
1:C:106:VAL:HG13	1:C:111:LEU:HD11	1.99	0.45
1:C:128:ILE:HG22	1:C:342:LEU:HG	1.99	0.45
1:C:326:ASP:O	1:C:330:ILE:HG12	2.17	0.44
1:A:66:TYR:OH	1:A:275:HIS:CD2	2.63	0.44
1:C:358:ILE:HG22	1:C:360:PRO:HD3	1.99	0.44
1:B:252:HIS:CD2	1:B:254:HIS:H	2.24	0.43
1:D:163:LEU:O	1:D:167:ILE:HG12	2.17	0.43
1:C:145:THR:HG23	1:C:147:GLY:H	1.83	0.43
1:B:376:ALA:O	1:B:380:VAL:HG23	2.19	0.42
1:D:81:MET:HE2	1:D:84:ILE:HD12	2.01	0.42
1:D:104:ILE:HG13	1:D:105:ASN:N	2.35	0.42
1:D:273:GLN:NE2	1:D:284:ARG:HD3	2.35	0.42
1:D:234:ILE:HD13	1:D:247:VAL:HG23	2.02	0.42
1:C:252:HIS:HE1	5:C:726:HOH:O	2.02	0.41
1:C:247:VAL:HB	2:P:444:LEU:HD23	2.02	0.41
1:A:39:HIS:HE1	1:C:307:GLU:O	2.02	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:254:HIS:HE1	5:A:799:HOH:O	2.04	0.41
1:B:273:GLN:HE22	1:B:284:ARG:HD3	1.85	0.41
1:D:254:HIS:CD2	1:D:256:GLU:HB2	2.56	0.40
1:B:218:ILE:HG12	1:B:220:LYS:HD3	2.02	0.40
1:A:252:HIS:CD2	1:A:254:HIS:H	2.27	0.40
1:A:293:TYR:OH	1:A:297:LYS:HD2	2.22	0.40
1:B:208:GLU:HG2	1:B:212:LYS:HE2	2.03	0.40
1:D:171:THR:HG23	1:D:359:ILE:HD11	2.01	0.40
1:C:252:HIS:CD2	1:C:254:HIS:H	2.32	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/385 (99%)	376 (98%)	4 (1%)	2 (0%)	29	17
1	B	383/385 (100%)	377 (98%)	5 (1%)	1 (0%)	41	30
1	C	381/385 (99%)	375 (98%)	5 (1%)	1 (0%)	41	30
1	D	378/385 (98%)	366 (97%)	9 (2%)	3 (1%)	19	9
2	M	10/12 (83%)	9 (90%)	1 (10%)	0	100	100
2	N	10/12 (83%)	9 (90%)	1 (10%)	0	100	100
2	P	10/12 (83%)	9 (90%)	1 (10%)	0	100	100
2	Q	10/12 (83%)	9 (90%)	1 (10%)	0	100	100
All	All	1564/1588 (98%)	1530 (98%)	27 (2%)	7 (0%)	34	22

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	57	ASP

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	57	ASP
1	D	7	LEU
1	B	110	ILE
1	A	110	ILE
1	C	110	ILE
1	D	110	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/339 (100%)	332 (98%)	6 (2%)	59	53
1	B	338/339 (100%)	333 (98%)	5 (2%)	65	60
1	C	337/339 (99%)	327 (97%)	10 (3%)	41	30
1	D	333/339 (98%)	325 (98%)	8 (2%)	49	40
2	M	9/9 (100%)	8 (89%)	1 (11%)	6	1
2	N	9/9 (100%)	7 (78%)	2 (22%)	1	0
2	P	9/9 (100%)	7 (78%)	2 (22%)	1	0
2	Q	9/9 (100%)	7 (78%)	2 (22%)	1	0
All	All	1382/1392 (99%)	1346 (97%)	36 (3%)	46	36

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	VAL
1	A	60	GLU
1	A	71	ASN
1	A	106	VAL
1	A	253	LEU
1	A	374	ARG
1	B	50	LEU
1	B	86	ARG
1	B	90	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	216	LYS
1	B	285	GLU
1	C	6	LYS
1	C	21	LYS
1	C	128	ILE
1	C	153	ASN
1	C	182	LEU
1	C	185	LEU
1	C	340	ASP
1	C	342	LEU
1	C	356	ARG
1	C	381	LYS
1	D	7	LEU
1	D	22	ILE
1	D	49	TYR
1	D	55	VAL
1	D	133	THR
1	D	185	LEU
1	D	307	GLU
1	D	342	LEU
2	M	444	LEU
2	N	443	ARG
2	N	444	LEU
2	P	444	LEU
2	P	453	ASP
2	Q	444	LEU
2	Q	453	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (36) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	GLN
1	A	38	GLN
1	A	39	HIS
1	A	71	ASN
1	A	74	GLN
1	A	130	ASN
1	A	161	ASN
1	A	204	HIS
1	A	252	HIS
1	A	254	HIS
1	A	275	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	23	GLN
1	B	74	GLN
1	B	130	ASN
1	B	161	ASN
1	B	169	HIS
1	B	252	HIS
1	B	254	HIS
1	B	273	GLN
1	C	23	GLN
1	C	131	ASN
1	C	153	ASN
1	C	161	ASN
1	C	252	HIS
1	C	254	HIS
1	C	273	GLN
1	C	275	HIS
1	D	38	GLN
1	D	39	HIS
1	D	65	HIS
1	D	74	GLN
1	D	161	ASN
1	D	252	HIS
1	D	254	HIS
1	D	273	GLN
1	D	275	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 11 ligands modelled in this entry, 7 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ATP	B	500	4	26,33,33	0.96	2 (7%)	31,52,52	1.36	5 (16%)
3	ATP	D	500	4	26,33,33	1.07	3 (11%)	31,52,52	1.39	6 (19%)
3	ATP	A	500	4	26,33,33	1.02	2 (7%)	31,52,52	1.32	4 (12%)
3	ATP	C	500	4	26,33,33	0.99	1 (3%)	31,52,52	1.37	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	B	500	4	-	1/18/38/38	0/3/3/3
3	ATP	D	500	4	-	3/18/38/38	0/3/3/3
3	ATP	A	500	4	-	2/18/38/38	0/3/3/3
3	ATP	C	500	4	-	3/18/38/38	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	ATP	C5-C4	2.79	1.48	1.40
3	C	500	ATP	C5-C4	2.69	1.48	1.40
3	B	500	ATP	C5-C4	2.48	1.47	1.40
3	D	500	ATP	O4'-C1'	2.36	1.44	1.41
3	D	500	ATP	C5-C4	2.32	1.47	1.40
3	D	500	ATP	C2-N3	2.26	1.35	1.32
3	B	500	ATP	O4'-C1'	2.05	1.43	1.41
3	A	500	ATP	C2-N3	2.04	1.35	1.32

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	500	ATP	N3-C2-N1	-3.76	122.80	128.68
3	C	500	ATP	N3-C2-N1	-3.70	122.89	128.68
3	D	500	ATP	N3-C2-N1	-3.63	123.00	128.68
3	A	500	ATP	N3-C2-N1	-3.39	123.38	128.68
3	C	500	ATP	PB-O3B-PG	-3.07	122.30	132.83
3	B	500	ATP	PB-O3B-PG	-2.70	123.56	132.83
3	A	500	ATP	PB-O3B-PG	-2.68	123.62	132.83
3	C	500	ATP	C2-N1-C6	2.57	123.16	118.75
3	B	500	ATP	C4-C5-N7	-2.53	106.76	109.40
3	A	500	ATP	C2-N1-C6	2.47	122.98	118.75
3	D	500	ATP	O3G-PG-O2G	2.27	116.32	107.64
3	D	500	ATP	PB-O3B-PG	-2.27	125.04	132.83
3	D	500	ATP	C4-C5-N7	-2.26	107.04	109.40
3	B	500	ATP	C2-N1-C6	2.25	122.61	118.75
3	A	500	ATP	C4-C5-N7	-2.09	107.22	109.40
3	B	500	ATP	PA-O3A-PB	-2.07	125.71	132.83
3	D	500	ATP	PA-O3A-PB	-2.07	125.73	132.83
3	D	500	ATP	C2-N1-C6	2.03	122.22	118.75
3	C	500	ATP	C4-C5-N7	-2.01	107.30	109.40

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	500	ATP	C5'-O5'-PA-O2A
3	D	500	ATP	C5'-O5'-PA-O2A
3	C	500	ATP	C5'-O5'-PA-O3A
3	D	500	ATP	C5'-O5'-PA-O3A
3	D	500	ATP	C5'-O5'-PA-O1A
3	A	500	ATP	C5'-O5'-PA-O3A
3	B	500	ATP	C5'-O5'-PA-O3A
3	A	500	ATP	C5'-O5'-PA-O1A
3	C	500	ATP	C5'-O5'-PA-O1A


There are no ring outliers.


1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	500	ATP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	384/385 (99%)	-0.11	5 (1%) 77 83	23, 33, 51, 93	0
1	B	385/385 (100%)	-0.12	4 (1%) 82 87	20, 32, 50, 65	0
1	C	383/385 (99%)	-0.05	3 (0%) 86 90	21, 32, 54, 80	0
1	D	380/385 (98%)	-0.04	9 (2%) 59 68	23, 36, 55, 92	0
2	M	12/12 (100%)	0.19	0 100 100	28, 32, 55, 63	0
2	N	12/12 (100%)	0.31	1 (8%) 11 17	25, 28, 51, 62	0
2	P	12/12 (100%)	0.38	2 (16%) 1 2	26, 30, 54, 61	0
2	Q	12/12 (100%)	0.19	1 (8%) 11 17	26, 30, 50, 58	0
All	All	1580/1588 (99%)	-0.07	25 (1%) 72 79	20, 33, 53, 93	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	5	TYR	8.7
1	D	56	VAL	5.1
1	A	56	VAL	4.7
1	D	7	LEU	4.4
1	A	57	ASP	3.8
1	B	287	PHE	3.6
2	P	442	GLY	3.5
1	B	5	TYR	3.3
2	N	443	ARG	3.3
1	D	287	PHE	3.2
1	D	29	ILE	2.9
2	P	443	ARG	2.7
1	A	287	PHE	2.6
1	A	59	LYS	2.5
1	D	374	ARG	2.5
2	Q	443	ARG	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	287	PHE	2.3
1	A	374	ARG	2.3
1	D	59	LYS	2.2
1	B	57	ASP	2.2
1	C	128	ILE	2.2
1	B	134	ILE	2.1
1	D	5	TYR	2.1
1	D	21	LYS	2.0
1	D	57	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ATP	C	500	31/31	0.94	0.12	19,23,39,40	0
3	ATP	D	500	31/31	0.95	0.10	26,31,40,43	0
4	MG	B	502	1/1	0.96	0.11	34,34,34,34	0
4	MG	D	502	1/1	0.96	0.20	34,34,34,34	0
3	ATP	B	500	31/31	0.96	0.09	21,25,39,42	0
4	MG	A	501	1/1	0.96	0.06	33,33,33,33	0
3	ATP	A	500	31/31	0.97	0.09	23,25,36,36	0
4	MG	A	502	1/1	0.97	0.08	29,29,29,29	0
4	MG	B	501	1/1	0.97	0.06	29,29,29,29	0
4	MG	D	501	1/1	0.98	0.03	36,36,36,36	0
4	MG	C	501	1/1	0.99	0.05	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around ATP C 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ATP D 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ATP B 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ATP A 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.