



wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 07:34 am BST

PDB ID : 1PG7
Title : Murine 6A6 Fab in complex with humanized anti-Tissue Factor D3H44 Fab
Authors : Eigenbrot, C.; Meng, Y.G.; Krishnamurthy, R.; Lipari, M.T.; Presta, L.; Devaux, B.; Wong, T.; Moran, P.; Bullens, S.; Kirchhofer, D.
Deposited on : 2003-05-27
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	H	217	
1	I	217	
2	L	213	
2	M	213	
3	W	210	
3	Y	210	
4	X	220	
4	Z	220	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13317 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called humanized antibody D3H44.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	213	Total	C	N	O	S	7	0	0
			1600	1011	266	317	6			
1	I	213	Total	C	N	O	S	3	0	0
			1600	1011	266	317	6			

- Molecule 2 is a protein called humanized antibody D3H44.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	213	Total	C	N	O	S	11	1	0
			1658	1038	280	335	5			
2	M	213	Total	C	N	O	S	0	0	0
			1647	1032	276	334	5			

- Molecule 3 is a protein called murine antibody 6A6 Fab fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	W	210	Total	C	N	O	S	5	0	0
			1581	988	265	323	5			
3	Y	210	Total	C	N	O	S	5	0	0
			1581	988	265	323	5			

- Molecule 4 is a protein called murine antibody 6A6 Fab fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	X	220	Total	C	N	O	S	1	2	0
			1688	1068	282	330	8			
4	Z	220	Total	C	N	O	S	7	2	0
			1688	1068	282	330	8			

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	26	Total 26	O 26	0	0
5	I	33	Total 33	O 33	0	0
5	L	33	Total 33	O 33	0	0
5	M	43	Total 43	O 43	0	0
5	W	41	Total 41	O 41	0	0
5	X	26	Total 26	O 26	0	0
5	Y	38	Total 38	O 38	0	0
5	Z	34	Total 34	O 34	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: humanized antibody D3H44




- Molecule 1: humanized antibody D3H44



- Molecule 2: humanized antibody D3H44



- Molecule 2: humanized antibody D3H44





- Molecule 3: murine antibody 6A6 Fab fragment







- Molecule 3: murine antibody 6A6 Fab fragment



- Molecule 4: murine antibody 6A6 Fab fragment



- Molecule 4: murine antibody 6A6 Fab fragment



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	78.89 Å 85.78 Å 92.37 Å 77.48° 75.92° 63.33°	Depositor
Resolution (Å)	50.00 – 2.50	Depositor
% Data completeness (in resolution range)	98.9 (50.00-2.50)	Depositor
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 98.1	Depositor
R, R_{free}	0.217 , 0.265	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	13317	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.46	0/1639	0.74	1/2236 (0.0%)
1	I	0.48	0/1639	0.75	2/2236 (0.1%)
2	L	0.49	0/1695	0.72	0/2301
2	M	0.50	0/1684	0.74	0/2287
3	W	0.46	0/1619	0.73	0/2211
3	Y	0.46	0/1619	0.73	0/2211
4	X	0.46	0/1733	0.77	1/2369 (0.0%)
4	Z	0.48	0/1733	0.79	1/2369 (0.0%)
All	All	0.48	0/13361	0.75	5/18220 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	95	ASP	CB-CG-OD1	6.92	124.53	118.30
4	Z	177	LEU	CA-CB-CG	6.44	130.12	115.30
4	X	177	LEU	CA-CB-CG	5.47	127.89	115.30
1	I	95	ASP	CB-CG-OD1	5.36	123.12	118.30
1	I	124	LEU	N-CA-C	-5.17	97.03	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	140	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1600	0	1549	19	0
1	I	1600	0	1549	17	0
2	L	1658	0	1610	25	0
2	M	1647	0	1598	23	0
3	W	1581	0	1519	15	0
3	Y	1581	0	1519	19	0
4	X	1688	0	1650	23	0
4	Z	1688	0	1650	31	0
5	H	26	0	0	0	0
5	I	33	0	0	1	0
5	L	33	0	0	1	0
5	M	43	0	0	2	0
5	W	41	0	0	3	0
5	X	26	0	0	0	0
5	Y	38	0	0	1	0
5	Z	34	0	0	3	0
All	All	13317	0	12644	158	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 158 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:X:70:THR:HG23	4:X:79:TYR:HB2	1.50	0.91
3:W:28:THR:HG23	3:W:30:SER:H	1.39	0.86
2:L:193:ALA:HB2	2:L:208:SER:HB3	1.60	0.83
2:L:166:GLN:HG3	2:L:173:TYR:CZ	2.23	0.74
2:M:166:GLN:HG3	2:M:173:TYR:CZ	2.24	0.73

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	209/217 (96%)	199 (95%)	9 (4%)	1 (0%)	29	48
1	I	209/217 (96%)	199 (95%)	10 (5%)	0	100	100
2	L	212/213 (100%)	207 (98%)	5 (2%)	0	100	100
2	M	211/213 (99%)	204 (97%)	7 (3%)	0	100	100
3	W	208/210 (99%)	197 (95%)	9 (4%)	2 (1%)	15	28
3	Y	208/210 (99%)	197 (95%)	9 (4%)	2 (1%)	15	28
4	X	220/220 (100%)	204 (93%)	13 (6%)	3 (1%)	11	20
4	Z	220/220 (100%)	205 (93%)	12 (6%)	3 (1%)	11	20
All	All	1697/1720 (99%)	1612 (95%)	74 (4%)	11 (1%)	22	43

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	X	44	ASN
4	X	127	VAL
4	Z	127	VAL
3	Y	41	ASP
4	Z	44	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	176/180 (98%)	170 (97%)	6 (3%)	37	63

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	I	176/180 (98%)	168 (96%)	8 (4%)	27	51
2	L	188/187 (100%)	180 (96%)	8 (4%)	29	53
2	M	187/187 (100%)	176 (94%)	11 (6%)	19	37
3	W	176/176 (100%)	165 (94%)	11 (6%)	18	34
3	Y	176/176 (100%)	171 (97%)	5 (3%)	43	70
4	X	195/193 (101%)	178 (91%)	17 (9%)	10	20
4	Z	195/193 (101%)	180 (92%)	15 (8%)	13	25
All	All	1469/1472 (100%)	1388 (94%)	81 (6%)	22	41

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	W	52	ASN
4	X	13	LYS
4	Z	128	CYS
3	W	69	ASP
3	W	154	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	199	GLN
3	W	52	ASN
4	Z	5	GLN
2	L	210	ASN
1	H	192	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.