



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 06:51 am BST

PDB ID : 1PH5  
Title : CRYSTAL STRUCTURE OF THE OXYTRICHA NOVA TELOMERE  
END-BINDING PROTEIN COMPLEXED WITH NONCOGNATE SSDNA  
GGGGTTTTG(3DR)GG  
Authors : Theobald, D.L.; Schultz, S.C.  
Deposited on : 2003-05-29  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	13	<div> <div>62%</div> <div>31%</div> <div>8%</div> </div>
1	H	13	<div> <div>46%</div> <div>46%</div> <div>8%</div> </div>
2	D	11	<div> <div>27%</div> <div>64%</div> <div>9%</div> </div>
3	A	459	<div> <div>3%</div> <div>80%</div> <div>19%</div> <div>.</div> </div>
4	B	216	<div> <div>9%</div> <div>73%</div> <div>26%</div> <div>.</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6658 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*GP\*GP\*GP\*GP\*TP\*TP\*TP\*TP\*GP\*GP\*GP\*GP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	12	Total	C	N	O	P	0	0	0
			253	120	48	74	11			
1	H	12	Total	C	N	O	P	0	0	0
			253	120	48	74	11			

- Molecule 2 is a DNA chain called 5'-D(\*GP\*GP\*GP\*GP\*TP\*TP\*TP\*TP\*GP\*(3DR)P\*GP\*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	11	Total	C	N	O	P	0	0	0
			219	105	38	66	10			

- Molecule 3 is a protein called Telomere-binding protein alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	459	Total	C	N	O	S	0	0	0
			3693	2346	636	709	2			

- Molecule 4 is a protein called Telomere-binding protein beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	216	Total	C	N	O	S	0	0	0
			1732	1112	294	325	1			

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	3	Total	Na	0	0
			3	3		
5	G	1	Total	Na	0	0
			1	1		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	G	34	Total O 34 34	0	0
7	D	30	Total O 30 30	0	0
7	H	34	Total O 34 34	0	0
7	A	330	Total O 330 330	0	0
7	B	75	Total O 75 75	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(\*GP\*GP\*GP\*GP\*TP\*TP\*TP\*TP\*GP\*GP\*GP\*GP\*T)-3'

Chain G: 




- Molecule 1: 5'-D(\*GP\*GP\*GP\*GP\*TP\*TP\*TP\*TP\*GP\*GP\*GP\*GP\*T)-3'

Chain H: 





- Molecule 2: 5'-D(\*GP\*GP\*GP\*GP\*TP\*TP\*TP\*TP\*GP\*(3DR)P\*GP\*G)-3'

Chain D: 





- Molecule 3: Telomere-binding protein alpha subunit

Chain A: 



- Molecule 4: Telomere-binding protein beta subunit

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.35Å 93.35Å 423.23Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.70 – 2.30 39.70 – 2.10	Depositor EDS
% Data completeness (in resolution range)	83.5 (39.70-2.30) 79.7 (39.70-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 2.10Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.228 , 0.258 0.222 , 0.256	Depositor DCC
$R_{free}$ test set	5207 reflections (10.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.5	Xtriage
Anisotropy	0.314	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 38.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6658	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, 3DR, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	G	0.63	0/284	0.70	0/440
1	H	0.57	0/284	0.67	0/440
2	D	0.93	1/232 (0.4%)	0.83	0/357
3	A	0.53	0/3763	0.61	1/5086 (0.0%)
4	B	0.44	0/1770	0.51	0/2391
All	All	0.54	1/6333 (0.0%)	0.61	1/8714 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	12	DG	C3'-O3'	-5.80	1.36	1.44

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	493	LEU	N-CA-C	9.64	137.02	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	253	0	138	4	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	253	0	138	4	0
2	D	219	0	121	12	0
3	A	3693	0	3690	61	0
4	B	1732	0	1718	41	1
5	G	1	0	0	0	0
5	H	3	0	0	0	0
6	A	1	0	0	1	0
7	A	330	0	0	5	2
7	B	75	0	0	0	0
7	D	30	0	0	1	0
7	G	34	0	0	1	0
7	H	34	0	0	0	0
All	All	6658	0	5805	112	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (112) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:285:LYS:HB3	3:A:331:ASN:HD21	1.27	0.97
4:B:10:GLN:HE21	4:B:12:SER:H	1.14	0.94
4:B:80:ILE:HD12	4:B:218:ILE:HD11	1.61	0.81
3:A:345:LEU:HD12	3:A:346:PRO:HD2	1.61	0.81
3:A:493:LEU:C	3:A:494:ILE:HG13	2.01	0.80
2:D:6:DT:H2''	2:D:7:DT:H5'	1.63	0.79
4:B:90:ILE:HG23	4:B:118:VAL:HG13	1.66	0.76
3:A:284:GLN:HG3	3:A:330:LEU:HD12	1.67	0.76
4:B:28:PHE:O	4:B:31:VAL:HG12	1.94	0.67
1:H:11:DG:N3	1:H:11:DG:H5'	2.10	0.67
3:A:76:LEU:HD23	3:A:76:LEU:N	2.11	0.66
3:A:229:LYS:HE3	3:A:310:ARG:HA	1.81	0.63
3:A:422:THR:HB	3:A:425:ASN:HD21	1.62	0.63
3:A:76:LEU:HA	7:A:1233:HOH:O	1.99	0.63
3:A:458:ARG:O	3:A:462:GLU:HG3	1.99	0.62
3:A:285:LYS:HB3	3:A:331:ASN:ND2	2.08	0.61
2:D:6:DT:O2	2:D:8:DT:H73	2.01	0.60
2:D:8:DT:H2''	2:D:9:DG:H5'	1.84	0.60
2:D:12:DG:H2'	2:D:12:DG:N3	2.16	0.59
3:A:155:SER:H	3:A:158:GLN:HE21	1.50	0.59
1:H:4:DG:H1'	1:H:5:DT:H5'	1.83	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:477:ALA:HB2	3:A:491:THR:HG21	1.84	0.58
3:A:155:SER:H	3:A:158:GLN:NE2	2.02	0.57
2:D:8:DT:H71	2:D:12:DG:H2''	1.85	0.57
4:B:20:GLU:HB3	4:B:31:VAL:HG23	1.86	0.57
3:A:183:LYS:HG2	7:A:1317:HOH:O	2.05	0.56
3:A:61:ALA:HB2	3:A:76:LEU:HB3	1.86	0.56
3:A:434:TYR:CD2	4:B:175:VAL:HG21	2.40	0.56
4:B:71:PHE:CE1	4:B:123:PRO:HD3	2.41	0.56
3:A:270:VAL:HG21	3:A:309:LEU:HB3	1.90	0.54
2:D:2:DG:H3'	2:D:3:DG:H5''	1.90	0.53
3:A:108:GLU:CD	3:A:108:GLU:H	2.11	0.53
4:B:21:LEU:HD22	4:B:95:LEU:HB3	1.91	0.53
3:A:459:LYS:HE3	3:A:463:ASP:OD2	2.09	0.53
3:A:283:THR:HG21	4:B:9:GLN:O	2.09	0.52
3:A:61:ALA:CB	3:A:76:LEU:HB3	2.39	0.52
4:B:215:ILE:O	4:B:218:ILE:HG22	2.10	0.52
3:A:418:LYS:HD3	4:B:156:LEU:HD21	1.92	0.52
4:B:52:VAL:O	4:B:58:PHE:HA	2.10	0.52
3:A:428:THR:HG22	4:B:190:VAL:O	2.09	0.51
3:A:228:ALA:HB2	3:A:247:ASP:HB3	1.92	0.51
3:A:200:PHE:CZ	3:A:205:VAL:HG11	2.46	0.51
4:B:27:ASP:OD2	4:B:30:LYS:HE3	2.12	0.51
3:A:335:LEU:O	3:A:494:ILE:N	2.44	0.50
3:A:149:PHE:CE2	3:A:171:PRO:HG3	2.46	0.49
2:D:2:DG:H3'	2:D:3:DG:C5'	2.43	0.49
3:A:435:THR:HG21	3:A:441:ALA:HA	1.95	0.49
1:G:3:DG:N3	1:G:3:DG:H2'	2.28	0.48
3:A:414:GLN:OE1	4:B:170:VAL:HG23	2.13	0.48
4:B:87:VAL:HG11	4:B:127:GLU:HB3	1.96	0.48
3:A:444:PHE:O	3:A:445:ASN:HB2	2.14	0.48
4:B:10:GLN:HG2	4:B:16:GLN:HB2	1.95	0.48
4:B:32:SER:HB2	4:B:35:LEU:HD12	1.95	0.48
3:A:285:LYS:HD3	3:A:331:ASN:ND2	2.29	0.47
3:A:48:SER:C	3:A:50:GLN:H	2.16	0.47
3:A:85:LEU:C	3:A:96:ASP:HB2	2.34	0.47
4:B:81:VAL:HG11	4:B:219:PHE:HD1	1.79	0.47
1:G:3:DG:H1'	1:G:4:DG:H5'	1.96	0.47
3:A:194:LYS:O	3:A:198:GLN:HG3	2.15	0.47
6:A:1005:CL:CL	4:B:182:LYS:HD2	2.51	0.47
3:A:149:PHE:CZ	3:A:171:PRO:HG3	2.48	0.47
3:A:434:TYR:HB3	3:A:436:GLN:OE1	2.15	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:1:DG:H1'	1:H:2:DG:H5'	1.97	0.47
3:A:323:SER:HB2	3:A:329:SER:OG	2.14	0.47
3:A:67:THR:HG21	3:A:73:ILE:HD12	1.97	0.46
2:D:5:DT:C7	7:A:1335:HOH:O	2.63	0.46
4:B:42:TYR:HB2	4:B:129:LEU:HD23	1.97	0.46
3:A:86:LYS:O	3:A:88:GLN:N	2.49	0.46
3:A:357:ALA:HB3	4:B:185:VAL:HG21	1.98	0.46
3:A:351:GLN:O	3:A:355:HIS:HB2	2.16	0.45
3:A:389:GLY:HA3	3:A:406:LYS:HB2	1.99	0.45
3:A:146:TRP:NE1	3:A:175:SER:HB3	2.32	0.45
3:A:281:THR:HG23	4:B:133:ARG:HH12	1.81	0.45
4:B:96:GLU:OE2	4:B:210:THR:HG22	2.17	0.45
4:B:216:ALA:O	4:B:220:VAL:HG23	2.17	0.45
4:B:87:VAL:CG1	4:B:127:GLU:HB3	2.47	0.45
3:A:57:VAL:HG23	3:A:192:LEU:HD11	2.00	0.44
1:G:4:DG:H1'	1:G:5:DT:H5'	1.99	0.44
3:A:277:THR:OG1	3:A:289:ILE:HB	2.18	0.44
3:A:338:VAL:HG12	3:A:477:ALA:HA	1.98	0.44
2:D:8:DT:H5'	7:D:341:HOH:O	2.16	0.43
3:A:373:PHE:CE1	3:A:417:VAL:HG11	2.53	0.43
2:D:5:DT:H1'	2:D:6:DT:H71	1.98	0.43
3:A:324:LEU:HD13	3:A:467:LEU:HD21	2.01	0.43
4:B:24:ASN:O	4:B:25:GLU:HG2	2.19	0.43
1:G:10:DG:H5''	7:G:489:HOH:O	2.19	0.43
3:A:204:SER:O	3:A:205:VAL:HB	2.19	0.42
3:A:48:SER:C	3:A:50:GLN:N	2.73	0.42
4:B:12:SER:HB3	4:B:15:LYS:HB3	2.01	0.42
4:B:27:ASP:CG	4:B:30:LYS:HG3	2.39	0.42
4:B:42:TYR:HB2	4:B:129:LEU:CD2	2.50	0.42
3:A:319:VAL:HB	7:A:1321:HOH:O	2.19	0.42
1:H:3:DG:H2'	1:H:3:DG:N3	2.35	0.42
3:A:434:TYR:HD2	4:B:175:VAL:HG21	1.85	0.42
2:D:6:DT:H2''	2:D:7:DT:C5'	2.44	0.42
3:A:390:TYR:HB2	3:A:410:ILE:HG12	2.01	0.42
3:A:226:VAL:HG12	3:A:227:VAL:N	2.35	0.42
4:B:178:ALA:HA	4:B:181:LYS:HD2	2.01	0.42
3:A:429:TYR:CD1	3:A:429:TYR:N	2.88	0.41
3:A:121:ARG:O	3:A:146:TRP:HA	2.21	0.41
3:A:281:THR:HG23	4:B:133:ARG:NH1	2.35	0.41
4:B:90:ILE:CG2	4:B:118:VAL:HG13	2.44	0.41
3:A:467:LEU:HD13	7:A:1213:HOH:O	2.19	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:209:ASN:O	4:B:209:ASN:CG	2.59	0.41
2:D:7:DT:H2'	2:D:7:DT:H6	1.73	0.41
3:A:281:THR:HG1	4:B:57:PHE:HZ	1.67	0.41
4:B:61:PRO:HB3	4:B:116:LEU:HD23	2.03	0.41
4:B:148:ILE:O	4:B:151:PHE:HB3	2.21	0.40
4:B:58:PHE:CD1	4:B:58:PHE:C	2.95	0.40
3:A:358:ASP:OD1	3:A:481:ARG:NH2	2.43	0.40
4:B:75:PHE:CZ	4:B:123:PRO:HD2	2.56	0.40
4:B:80:ILE:HA	4:B:83:LEU:HG	2.04	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1258:HOH:O	7:A:1258:HOH:O[12_555]	0.91	1.29
7:A:1256:HOH:O	7:A:1256:HOH:O[12_555]	1.06	1.14
4:B:130:ASN:ND2	4:B:130:ASN:ND2[12_555]	1.81	0.39

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	457/459 (100%)	432 (94%)	22 (5%)	3 (1%)	22	26
4	B	214/216 (99%)	200 (94%)	13 (6%)	1 (0%)	29	35
All	All	671/675 (99%)	632 (94%)	35 (5%)	4 (1%)	25	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	87	GLN
4	B	223	LYS

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
3	A	89	LYS
3	A	404	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	407/407 (100%)	395 (97%)	12 (3%)	42	58
4	B	189/189 (100%)	182 (96%)	7 (4%)	34	48
All	All	596/596 (100%)	577 (97%)	19 (3%)	39	54

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	76	LEU
3	A	129	LEU
3	A	231	LEU
3	A	243	LEU
3	A	251	GLN
3	A	283	THR
3	A	309	LEU
3	A	353	LEU
3	A	362	GLU
3	A	398	SER
3	A	482	ARG
3	A	494	ILE
4	B	9	GLN
4	B	41	CYS
4	B	53	THR
4	B	66	GLU
4	B	92	ASN
4	B	128	ARG
4	B	215	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	87	GLN
3	A	158	GLN
3	A	327	ASN
3	A	331	ASN
3	A	351	GLN
3	A	372	GLN
3	A	426	ASN
3	A	445	ASN
4	B	10	GLN
4	B	79	ASN
4	B	92	ASN
4	B	149	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	3DR	D	10	2	8,11,12	0.69	0	9,14,17	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3DR	D	10	2	-	1/3/15/16	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	10	3DR	C4'-C5'-O5'-P

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	G	12/13 (92%)	-0.20	0	100	100	33, 39, 48, 51	0
1	H	12/13 (92%)	-0.09	0	100	100	38, 46, 50, 51	0
2	D	10/11 (90%)	-0.70	0	100	100	23, 30, 44, 46	0
3	A	459/459 (100%)	-0.21	13 (2%)	53	60	19, 30, 54, 82	0
4	B	216/216 (100%)	0.26	19 (8%)	10	13	25, 47, 75, 82	0
All	All	709/712 (99%)	-0.07	32 (4%)	33	40	19, 33, 69, 82	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	89	LYS	7.7
4	B	9	GLN	7.4
3	A	91	ALA	7.2
3	A	88	GLN	6.6
3	A	405	GLY	6.3
3	A	92	GLY	6.2
4	B	125	LEU	6.2
3	A	90	GLY	6.1
4	B	219	PHE	5.6
4	B	216	ALA	5.2
4	B	196	LYS	4.2
4	B	78	VAL	3.9
3	A	69	GLN	3.8
3	A	404	SER	3.8
3	A	403	ALA	3.7
4	B	126	GLN	3.6
4	B	220	VAL	3.4
3	A	494	ILE	3.0
4	B	77	ASN	3.0
3	A	87	GLN	2.6

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
4	B	223	LYS	2.6
4	B	35	LEU	2.5
4	B	122	LYS	2.5
4	B	213	LEU	2.4
4	B	81	VAL	2.4
4	B	211	ALA	2.4
4	B	224	GLY	2.2
3	A	70	GLU	2.2
4	B	217	ASP	2.1
3	A	93	ASP	2.1
4	B	65	LYS	2.1
4	B	82	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	3DR	D	10	11/12	0.97	0.10	27,30,34,35	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NA	H	1004	1/1	0.87	0.22	43,43,43,43	0
5	NA	H	1003	1/1	0.89	0.07	46,46,46,46	0
5	NA	G	1001	1/1	0.95	0.11	34,34,34,34	0
5	NA	H	1002	1/1	0.98	0.09	39,39,39,39	0
6	CL	A	1005	1/1	0.99	0.05	42,42,42,42	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.