



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2020 – 10:06 PM BST

PDB ID : 6PH5
Title : Binary product complex crystal structure of DNA polymerase Beta with an extra-helical template base
Authors : Batra, V.K.; Wilson, S.H.
Deposited on : 2019-06-25
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.13
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	<div> <div>7%</div> <div> <div></div> <div>71%</div> <div>24%</div> <div>• •</div> </div> </div>
2	T	17	<div> <div>12%</div> <div> <div></div> <div>65%</div> <div>29%</div> <div>6%</div> </div> </div>
3	P	11	<div> <div>9%</div> <div> <div></div> <div>64%</div> <div>36%</div> </div> </div>
4	D	5	<div> <div></div> <div> <div></div> <div>80%</div> <div>20%</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3302 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	0	0	0
			2576	1631	451	485	9			

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*CP*GP*AP*CP*GP*CP*GP*CP*GP*CP*AP*TP*CP*AP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	17	Total	C	N	O	P	0	0	0
			342	162	66	98	16			

- Molecule 3 is a DNA chain called DNA (5'-D(*GP*CP*TP*GP*AP*TP*GP*CP*GP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	11	Total	C	N	O	P	0	0	0
			222	106	41	65	10			

- Molecule 4 is a DNA chain called DNA (5'-D(P*GP*TP*CP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	5	Total	C	N	O	P	0	0	0
			106	49	20	32	5			

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Na	0	0
			2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	36	Total 36	O 36	0	0
6	T	10	Total 10	O 10	0	0
6	P	6	Total 6	O 6	0	0
6	D	2	Total 2	O 2	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.13 Å 79.12 Å 55.20 Å 90.00° 106.61° 90.00°	Depositor
Resolution (Å)	22.60 – 2.60 22.60 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (22.60-2.60) 88.7 (22.60-2.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 2.60 Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.205 , 0.279 0.205 , 0.279	Depositor DCC
R_{free} test set	1380 reflections (10.03%)	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtriage
Anisotropy	0.298	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.043 for l,-k,h	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3302	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	5/2624 (0.2%)	0.71	2/3527 (0.1%)
2	T	0.91	0/383	0.99	1/588 (0.2%)
3	P	1.01	0/248	1.03	0/381
4	D	1.23	1/118 (0.8%)	1.01	0/179
All	All	0.72	6/3373 (0.2%)	0.79	3/4675 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	320	PHE	CE1-CZ	9.38	1.55	1.37
1	A	325	TRP	NE1-CE2	-7.59	1.27	1.37
4	D	1	DG	OP3-P	-7.56	1.52	1.61
1	A	325	TRP	CD2-CE3	-6.49	1.30	1.40
1	A	325	TRP	CG-CD1	-5.62	1.28	1.36
1	A	325	TRP	CZ2-CH2	-5.36	1.27	1.37

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	40	ARG	NE-CZ-NH1	7.35	123.97	120.30
2	T	7	DC	C1'-O4'-C4'	-5.63	104.47	110.10
1	A	287	LEU	CA-CB-CG	-5.50	102.64	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2576	0	2583	83	0
2	T	342	0	190	4	0
3	P	222	0	125	3	0
4	D	106	0	57	1	0
5	A	2	0	0	0	0
6	A	36	0	0	4	0
6	D	2	0	0	0	0
6	P	6	0	0	0	0
6	T	10	0	0	1	0
All	All	3302	0	2955	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:THR:HB	1:A:301:LEU:HD21	1.59	0.84
1:A:171:SER:OG	6:A:501:HOH:O	1.96	0.82
2:T:13:DT:OP1	6:T:101:HOH:O	1.97	0.81
1:A:25:PHE:CE1	1:A:88:ILE:HD11	2.23	0.74
1:A:326:LYS:HG2	1:A:327:TYR:H	1.53	0.73
1:A:317:LYS:HA	1:A:320:PHE:HD1	1.53	0.72
1:A:317:LYS:HA	1:A:320:PHE:CD1	2.25	0.71
1:A:286:ALA:HA	1:A:323:ILE:HD12	1.75	0.69
1:A:328:ARG:HB3	1:A:328:ARG:NH1	2.09	0.68
1:A:316:GLU:O	1:A:319:ILE:HG22	1.95	0.67
1:A:292:THR:N	1:A:299:ARG:O	2.24	0.67
1:A:279:ASN:HA	1:A:282:MET:HB3	1.77	0.66
1:A:60:LYS:NZ	1:A:67:THR:OG1	2.29	0.65
1:A:151:PRO:HG2	1:A:154:GLU:HG3	1.79	0.64
1:A:212:HIS:NE2	1:A:216:GLU:OE1	2.30	0.64
1:A:124:ASP:O	1:A:128:ASN:ND2	2.25	0.64
1:A:123:GLU:HG3	1:A:126:ARG:HH12	1.66	0.61
1:A:282:MET:O	1:A:286:ALA:HB2	2.02	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:ALA:HA	1:A:323:ILE:CD1	2.32	0.59
1:A:276:ASP:N	6:A:502:HOH:O	2.09	0.56
1:A:25:PHE:CD1	1:A:88:ILE:HD11	2.40	0.56
1:A:128:ASN:HB3	1:A:131:LYS:HZ2	1.72	0.55
1:A:129:GLU:OE2	1:A:140:LEU:HD23	2.09	0.53
1:A:323:ILE:CD1	1:A:325:TRP:CZ2	2.92	0.53
2:T:7:DC:H5''	2:T:8:DG:OP2	2.09	0.53
1:A:298:ILE:HG23	1:A:311:LEU:HD23	1.90	0.53
1:A:251:PRO:HG2	1:A:253:ARG:CZ	2.39	0.53
1:A:311:LEU:HG	1:A:322:TYR:CE1	2.44	0.53
1:A:129:GLU:OE1	1:A:137:ARG:HG2	2.09	0.53
1:A:197:HIS:ND1	1:A:199:SER:OG	2.34	0.53
1:A:123:GLU:HG3	1:A:126:ARG:NH1	2.25	0.52
1:A:292:THR:O	1:A:298:ILE:HA	2.11	0.51
3:P:1:DG:H2'	3:P:2:DC:C6	2.45	0.51
2:T:15:DA:H1'	2:T:16:DG:H5'	1.93	0.51
1:A:289:LYS:HG3	1:A:323:ILE:HG13	1.93	0.50
2:T:6:DG:H1	3:P:11:DC:H42	1.59	0.49
1:A:311:LEU:HD23	1:A:311:LEU:H	1.78	0.49
1:A:316:GLU:CD	1:A:333:ARG:HH22	2.16	0.49
1:A:287:LEU:HD21	1:A:291:PHE:C	2.33	0.49
1:A:85:LEU:O	1:A:88:ILE:HG22	2.13	0.48
1:A:197:HIS:CG	1:A:198:PRO:HD2	2.48	0.48
1:A:287:LEU:HA	1:A:287:LEU:HD23	1.37	0.48
1:A:323:ILE:HD13	1:A:325:TRP:CZ2	2.49	0.48
1:A:326:LYS:HE2	1:A:326:LYS:HB3	1.61	0.47
1:A:317:LYS:NZ	1:A:327:TYR:HA	2.29	0.47
1:A:328:ARG:HB3	1:A:328:ARG:CZ	2.44	0.47
1:A:204:SER:O	1:A:204:SER:OG	2.28	0.47
1:A:276:ASP:CG	1:A:277:ILE:HD12	2.35	0.47
1:A:85:LEU:O	1:A:89:ARG:HB2	2.16	0.46
1:A:167:LYS:HG3	1:A:171:SER:HA	1.97	0.46
1:A:25:PHE:CD1	1:A:29:VAL:HB	2.51	0.46
1:A:283:ARG:HG2	1:A:293:ILE:CG2	2.46	0.45
1:A:277:ILE:HA	1:A:280:LYS:HG2	1.97	0.45
1:A:291:PHE:CE1	1:A:322:TYR:O	2.69	0.45
1:A:323:ILE:O	1:A:323:ILE:HG12	2.17	0.45
1:A:207:GLN:HA	1:A:208:PRO:HD3	1.83	0.45
1:A:24:ASN:OD1	6:A:503:HOH:O	2.21	0.45
1:A:218:LEU:HB2	1:A:224:ILE:HD12	1.99	0.44
1:A:25:PHE:CZ	1:A:88:ILE:HD11	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:THR:O	1:A:97:ILE:HG13	2.17	0.44
1:A:170:ASP:HB3	1:A:173:TYR:CD2	2.52	0.44
1:A:183:ARG:CZ	1:A:275:SER:HB3	2.48	0.43
1:A:292:THR:HG22	1:A:299:ARG:HB2	1.98	0.43
1:A:11:LEU:O	1:A:52:LYS:HE2	2.17	0.43
1:A:301:LEU:HD22	1:A:301:LEU:N	2.34	0.43
1:A:196:THR:HG22	1:A:197:HIS:N	2.34	0.43
1:A:326:LYS:HG2	1:A:327:TYR:N	2.29	0.43
1:A:123:GLU:N	1:A:123:GLU:OE1	2.31	0.43
1:A:319:ILE:O	1:A:323:ILE:HG22	2.19	0.43
1:A:331:LYS:HE3	1:A:331:LYS:HB3	1.73	0.43
1:A:288:GLU:O	1:A:289:LYS:HG2	2.20	0.42
1:A:236:MET:HG2	1:A:256:ASP:OD1	2.20	0.42
1:A:298:ILE:HG23	1:A:311:LEU:CD2	2.50	0.41
1:A:328:ARG:HH11	1:A:328:ARG:HB3	1.78	0.41
1:A:128:ASN:HB3	1:A:131:LYS:NZ	2.36	0.41
1:A:292:THR:HG22	1:A:299:ARG:O	2.20	0.41
1:A:68:LYS:NZ	4:D:1:DG:OP1	2.41	0.41
1:A:131:LYS:HZ2	1:A:131:LYS:HG3	1.50	0.41
1:A:329:GLU:HA	6:A:511:HOH:O	2.20	0.41
1:A:266:TYR:HB3	1:A:316:GLU:HA	2.03	0.41
1:A:196:THR:OG1	1:A:265:TYR:CD1	2.74	0.40
1:A:292:THR:O	1:A:298:ILE:HD12	2.21	0.40
1:A:113:LYS:HE2	1:A:113:LYS:HB3	1.75	0.40
1:A:209:LYS:HB2	1:A:209:LYS:HE2	1.94	0.40
1:A:239:CYS:SG	1:A:255:ILE:HB	2.62	0.40
1:A:69:ILE:O	1:A:73:ILE:HG13	2.20	0.40
3:P:10:DC:H2'	3:P:11:DC:C5	2.56	0.40
1:A:276:ASP:C	1:A:280:LYS:HZ2	2.24	0.40
1:A:196:THR:CG2	1:A:197:HIS:N	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/335 (96%)	298 (93%)	20 (6%)	2 (1%)	25	47

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	289	LYS
1	A	330	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/295 (95%)	274 (98%)	5 (2%)	59	80

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	ARG
1	A	243	SER
1	A	317	LYS
1	A	325	TRP
1	A	328	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/335 (96%)	0.25	24 (7%) 14 10	29, 44, 93, 123	0
2	T	17/17 (100%)	0.33	2 (11%) 4 3	33, 46, 91, 120	0
3	P	11/11 (100%)	0.07	1 (9%) 9 6	36, 43, 63, 79	0
4	D	5/5 (100%)	-0.66	0 100 100	31, 33, 38, 45	0
All	All	357/368 (97%)	0.23	27 (7%) 13 10	29, 44, 93, 123	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	327	TYR	8.2
2	T	7	DC	6.5
1	A	245	ASN	5.7
1	A	10	THR	4.8
1	A	300	PRO	4.1
1	A	207	GLN	3.7
1	A	291	PHE	3.5
1	A	306	VAL	3.5
3	P	11	DC	3.5
1	A	11	LEU	3.4
1	A	208	PRO	3.2
1	A	312	PRO	3.0
1	A	335	GLU	3.0
1	A	326	LYS	3.0
1	A	324	GLN	2.9
1	A	314	ASP	2.9
1	A	290	GLY	2.8
1	A	304	THR	2.7
1	A	308	GLY	2.4
2	T	8	DG	2.3
1	A	303	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	246	ASP	2.2
1	A	305	GLY	2.2
1	A	321	ASP	2.2
1	A	322	TYR	2.2
1	A	302	GLY	2.0
1	A	334	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NA	A	402	1/1	0.94	0.10	35,35,35,35	0
5	NA	A	401	1/1	0.97	0.05	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.