



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 11:24 PM BST

PDB ID : 3PIJ
Title : beta-fructofuranosidase from Bifidobacterium longum - complex with fructose
Authors : Bujacz, A.; Bujacz, G.; Redzyna, I.; Krzepkowska-Jedrzejczak, M.; Bielecki, S.
Deposited on : 2010-11-06
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>9%</div> <div>..</div> </div> </div>
1	B	526	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>12%</div> <div>.</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9378 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-fructofuranosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	523	Total	C	N	O	S	0	4	0
			4152	2619	710	796	27			
1	B	526	Total	C	N	O	S	0	9	0
			4208	2652	722	808	26			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	240	SER	PHE	engineered mutation	UNP A2TLS9
A	519	LEU	-	expression tag	UNP A2TLS9
A	520	GLU	-	expression tag	UNP A2TLS9
A	521	HIS	-	expression tag	UNP A2TLS9
A	522	HIS	-	expression tag	UNP A2TLS9
A	523	HIS	-	expression tag	UNP A2TLS9
A	524	HIS	-	expression tag	UNP A2TLS9
A	525	HIS	-	expression tag	UNP A2TLS9
A	526	HIS	-	expression tag	UNP A2TLS9
B	240	SER	PHE	engineered mutation	UNP A2TLS9
B	519	LEU	-	expression tag	UNP A2TLS9
B	520	GLU	-	expression tag	UNP A2TLS9
B	521	HIS	-	expression tag	UNP A2TLS9
B	522	HIS	-	expression tag	UNP A2TLS9
B	523	HIS	-	expression tag	UNP A2TLS9
B	524	HIS	-	expression tag	UNP A2TLS9
B	525	HIS	-	expression tag	UNP A2TLS9
B	526	HIS	-	expression tag	UNP A2TLS9

- Molecule 2 is beta-D-fructofuranose (three-letter code: FRU) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Cl	0	0
			3	3		
3	A	2	Total	Cl	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	548	Total	O	0	0
			548	548		
4	B	441	Total	O	0	0
			441	441		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-fructofuranosidase



• Molecule 1: Beta-fructofuranosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	86.77Å 86.77Å 223.87Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.40 – 1.80 28.40 – 1.80	Depositor EDS
% Data completeness (in resolution range)	92.1 (28.40-1.80) 92.2 (28.40-1.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0091	Depositor
R, R_{free}	0.153 , 0.208 0.164 , 0.217	Depositor DCC
R_{free} test set	1683 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	21.8	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.030 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9378	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FRU, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	0/4283	0.90	5/5817 (0.1%)
1	B	0.93	2/4357 (0.0%)	0.86	2/5918 (0.0%)
All	All	0.98	2/8640 (0.0%)	0.88	7/11735 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	462	VAL	CB-CG2	5.83	1.65	1.52
1	B	462	VAL	CB-CG1	5.26	1.63	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	96	ARG	NE-CZ-NH1	-5.98	117.31	120.30
1	A	341	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	A	312	ARG	NE-CZ-NH2	-5.67	117.47	120.30
1	A	385	ASP	CB-CG-OD1	5.42	123.17	118.30
1	B	341	ARG	NE-CZ-NH2	-5.21	117.70	120.30
1	A	16	ASP	CB-CG-OD1	5.20	122.98	118.30
1	B	525	HIS	CB-CA-C	5.07	120.53	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4152	0	3910	30	0
1	B	4208	0	3956	64	0
2	A	12	0	12	0	0
2	B	12	0	12	0	0
3	A	2	0	0	0	0
3	B	3	0	0	0	0
4	A	548	0	0	10	0
4	B	441	0	0	8	0
All	All	9378	0	7890	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:284:GLY:HA3	4:B:986:HOH:O	1.37	1.24
1:B:166:MET:HE3	1:B:169:ASP:HA	1.36	1.07
1:B:166:MET:CE	1:B:169:ASP:HA	1.90	1.02
1:B:166:MET:CE	1:B:169:ASP:CA	2.45	0.94
1:B:17:HIS:HE1	1:B:102:ALA:H	1.12	0.91
1:A:77:GLN:HG3	4:A:877:HOH:O	1.69	0.91
1:A:312:ARG:HD2	1:A:342:GLU:OE1	1.73	0.87
1:A:436[B]:MET:HE3	1:A:485:TYR:H	1.44	0.83
1:B:166:MET:CE	1:B:169:ASP:CB	2.57	0.83
1:B:166:MET:HE1	1:B:169:ASP:CB	2.11	0.79
1:B:36[A]:ASN:HD22	1:B:331:ASP:HA	1.51	0.74
1:B:178:HIS:HD2	1:B:180:ARG:HH12	1.36	0.73
1:B:1:MET:HG3	1:B:1:MET:O	1.89	0.71
1:B:166:MET:HE1	1:B:169:ASP:CG	2.10	0.71
1:A:77:GLN:CG	4:A:877:HOH:O	2.34	0.71
1:A:436[B]:MET:CE	1:A:485:TYR:H	2.05	0.70
1:B:17:HIS:CE1	1:B:102:ALA:H	2.03	0.69
1:B:155:ASN:HD21	1:B:159:THR:H	1.41	0.68
1:B:166:MET:CE	1:B:169:ASP:CG	2.61	0.68
1:B:166:MET:HE2	1:B:169:ASP:CA	2.24	0.67
1:A:300:ASN:HD22	1:A:483:TYR:H	1.43	0.67
1:A:155:ASN:HD21	1:A:159:THR:H	1.42	0.66
1:B:300:ASN:HD21	1:B:433:ARG:HH12	1.44	0.65
1:A:256:SER:HB3	1:A:302:TYR:CZ	2.34	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:178:HIS:CD2	1:B:180:ARG:HH12	2.17	0.62
1:B:166:MET:HE1	1:B:169:ASP:HB2	1.81	0.62
1:B:8:THR:HG22	1:B:216[B]:ARG:NH2	2.16	0.60
1:A:300:ASN:ND2	1:A:483:TYR:H	1.99	0.59
1:A:392:GLU:OE1	1:A:510:HIS:HE1	1.86	0.59
1:B:221:ARG:HD2	1:B:283:GLY:O	2.03	0.58
1:B:300:ASN:HD22	1:B:483:TYR:H	1.49	0.58
1:A:365:ASP:OD2	1:A:511:HIS:HD2	1.86	0.58
1:B:256:SER:HB3	1:B:302:TYR:CZ	2.38	0.58
1:A:373[B]:THR:HG23	4:A:985:HOH:O	2.02	0.58
1:B:526:HIS:CD2	1:B:526:HIS:OXT	2.57	0.58
1:B:8:THR:HG22	1:B:216[B]:ARG:HH22	1.68	0.58
1:B:8:THR:CG2	1:B:216[B]:ARG:NH2	2.66	0.57
1:A:44:HIS:NE2	1:A:470:GLU:OE2	2.31	0.57
1:B:166:MET:HE3	1:B:169:ASP:CA	2.14	0.57
1:B:526:HIS:HD2	1:B:526:HIS:OXT	1.86	0.57
1:B:379:GLU:HG2	1:B:496:VAL:HG22	1.86	0.57
1:B:525:HIS:O	1:B:526:HIS:HB2	2.05	0.56
1:B:166:MET:HE2	1:B:169:ASP:N	2.20	0.56
1:A:77:GLN:HG3	4:A:774:HOH:O	2.05	0.56
1:B:300:ASN:ND2	1:B:483:TYR:H	2.03	0.56
1:A:312:ARG:CD	1:A:342:GLU:OE1	2.51	0.55
1:A:391:ILE:HD11	1:A:493:ILE:HD13	1.89	0.55
1:B:392:GLU:OE1	1:B:510:HIS:HE1	1.90	0.55
1:B:7:GLU:O	1:B:216[B]:ARG:NE	2.41	0.53
1:B:166:MET:HE3	1:B:169:ASP:CG	2.30	0.52
1:B:173:ASP:HA	4:B:960:HOH:O	2.09	0.52
1:B:457:LYS:NZ	4:B:599:HOH:O	2.43	0.51
1:A:300:ASN:HD21	1:A:433:ARG:HH12	1.59	0.50
1:A:4:PHE:H	1:A:4:PHE:HD2	1.60	0.49
1:B:374:LEU:HD21	1:B:380:GLN:HG2	1.93	0.49
1:A:436[B]:MET:HE2	4:A:1030:HOH:O	2.12	0.49
1:B:108:GLU:OE1	1:B:163:LYS:NZ	2.39	0.49
1:B:274:GLY:HA3	1:B:292:PHE:CE1	2.49	0.48
1:B:120:ASP:HB2	1:B:214:MET:SD	2.53	0.48
1:B:7:GLU:O	1:B:216[B]:ARG:NH2	2.47	0.48
1:B:23:LYS:HE3	4:B:673:HOH:O	2.14	0.48
1:A:88:SER:HB2	1:A:94:TRP:CD2	2.50	0.47
1:B:7:GLU:O	1:B:216[B]:ARG:CZ	2.63	0.47
1:A:503:LYS:HD3	4:A:1008:HOH:O	2.16	0.46
1:B:313:GLN:NE2	4:B:698:HOH:O	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:391:ILE:CD1	1:B:493:ILE:HG21	2.46	0.46
1:A:77:GLN:CD	4:A:877:HOH:O	2.55	0.45
1:A:436[B]:MET:HG2	1:A:438:ASN:O	2.17	0.44
1:B:274:GLY:HA3	1:B:292:PHE:CZ	2.52	0.44
1:B:525:HIS:O	1:B:526:HIS:CB	2.63	0.44
1:B:256:SER:HB3	1:B:302:TYR:CE1	2.53	0.44
1:A:347[B]:ASP:OD1	4:A:721:HOH:O	2.21	0.44
1:A:436[B]:MET:CE	4:A:1030:HOH:O	2.65	0.44
1:A:404:ARG:NH2	1:B:434:GLN:HB3	2.33	0.44
1:B:393:MET:HB3	1:B:507[B]:LEU:HD22	2.00	0.43
1:B:166:MET:HE3	1:B:169:ASP:OD1	2.18	0.43
1:B:379:GLU:HA	1:B:495:LEU:O	2.18	0.43
1:B:72:HIS:HE1	4:B:866:HOH:O	2.01	0.43
1:B:139[A]:ASP:OD2	1:B:140:ASN:N	2.51	0.43
1:B:47:SER:HB2	1:B:52:ILE:CG2	2.48	0.43
1:B:166:MET:HE2	1:B:169:ASP:HA	1.84	0.43
1:A:128:TYR:OH	1:A:214:MET:HB3	2.20	0.42
1:B:8:THR:CG2	1:B:216[B]:ARG:HH22	2.28	0.42
1:B:301:TYR:OH	1:B:304:PRO:HB3	2.19	0.42
1:B:364:GLU:HG2	1:B:513:LYS:HG3	2.02	0.42
1:A:523:HIS:HB2	4:A:949:HOH:O	2.20	0.41
1:B:8:THR:HG23	1:B:216[B]:ARG:NH2	2.34	0.41
1:B:64:ARG:HD2	4:B:976:HOH:O	2.20	0.41
1:B:139[A]:ASP:CG	1:B:140:ASN:H	2.24	0.41
1:A:303:ALA:N	1:A:304:PRO:CD	2.84	0.41
1:B:382:ILE:HD12	1:B:507[A]:LEU:HD21	2.03	0.41
1:A:364:GLU:HG2	1:A:513[A]:LYS:HG2	2.02	0.40
1:B:246:ASP:HB2	4:B:977:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	525/526 (100%)	504 (96%)	19 (4%)	2 (0%)	34	21
1	B	533/526 (101%)	514 (96%)	17 (3%)	2 (0%)	34	21
All	All	1058/1052 (101%)	1018 (96%)	36 (3%)	4 (0%)	34	21

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	181	ASP
1	B	321	PRO
1	B	181	ASP
1	A	321	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	438/437 (100%)	427 (98%)	11 (2%)	47	34
1	B	446/437 (102%)	434 (97%)	12 (3%)	44	31
All	All	884/874 (101%)	861 (97%)	23 (3%)	46	32

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	2	THR
1	A	3	ASP
1	A	4	PHE
1	A	95	LYS
1	A	139	ASP
1	A	155	ASN
1	A	281	GLU
1	A	287	LYS
1	A	440	ASP
1	A	522	HIS
1	B	1	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	2	THR
1	B	7	GLU
1	B	54	ASP
1	B	100	MET
1	B	155	ASN
1	B	173	ASP
1	B	212	LYS
1	B	221	ARG
1	B	455	SER
1	B	459	ASP
1	B	526	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	GLN
1	A	155	ASN
1	A	300	ASN
1	A	313	GLN
1	A	369	HIS
1	A	380	GLN
1	A	510	HIS
1	A	511	HIS
1	B	17	HIS
1	B	72	HIS
1	B	155	ASN
1	B	178	HIS
1	B	300	ASN
1	B	313	GLN
1	B	510	HIS
1	B	526	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FRU	B	527	-	11,12,12	0.69	0	10,18,18	0.91	0
2	FRU	A	527	-	11,12,12	0.79	0	10,18,18	1.22	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	B	527	-	-	1/5/24/24	0/1/1/1
2	FRU	A	527	-	-	1/5/24/24	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	527	FRU	O2-C2-O5	-2.35	104.98	109.50

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	527	FRU	O1-C1-C2-C3
2	A	527	FRU	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	523/526 (99%)	-0.23	8 (1%) 73 70	13, 21, 35, 64	0
1	B	526/526 (100%)	-0.10	12 (2%) 60 56	15, 25, 41, 67	0
All	All	1049/1052 (99%)	-0.17	20 (1%) 66 63	13, 23, 39, 67	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	THR	8.7
1	A	2	THR	8.5
1	A	1	MET	7.8
1	B	1	MET	7.5
1	A	522	HIS	5.1
1	B	525	HIS	5.1
1	A	523	HIS	4.8
1	B	3	ASP	4.2
1	B	526	HIS	4.1
1	B	247	GLY	3.4
1	B	246	ASP	3.3
1	A	521	HIS	3.0
1	A	3	ASP	2.7
1	B	173	ASP	2.5
1	B	201	ASP	2.5
1	A	189	ASP	2.4
1	B	141	THR	2.3
1	A	519	LEU	2.3
1	B	4	PHE	2.2
1	B	521	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FRU	A	527	12/12	0.94	0.09	19,24,26,26	0
2	FRU	B	527	12/12	0.95	0.10	27,29,31,32	0
3	CL	B	529	1/1	0.97	0.05	32,32,32,32	0
3	CL	B	528	1/1	0.97	0.07	37,37,37,37	0
3	CL	A	528	1/1	0.98	0.06	30,30,30,30	0
3	CL	A	529	1/1	0.99	0.13	38,38,38,38	0
3	CL	B	530	1/1	0.99	0.02	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.