



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 04:09 am BST

PDB ID : 2PNV  
Title : Crystal Structure of the leucine zipper domain of small-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  (SKCa) channel from *Rattus norvegicus*  
Authors : Kim, J.Y.; Kim, M.K.; Kang, G.B.; Park, C.S.; Eom, S.H.  
Deposited on : 2007-04-25  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	43	
1	B	43	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 638 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Small conductance calcium-activated potassium channel protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	39	Total	C	N	O	S	0	0	0
			317	199	51	65	2			
1	B	39	Total	C	N	O	S	0	0	0
			317	199	51	65	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	484	GLY	-	EXPRESSION TAG	UNP P70604
A	485	SER	-	EXPRESSION TAG	UNP P70604
A	486	HIS	-	EXPRESSION TAG	UNP P70604
A	487	MET	-	EXPRESSION TAG	UNP P70604
B	484	GLY	-	EXPRESSION TAG	UNP P70604
B	485	SER	-	EXPRESSION TAG	UNP P70604
B	486	HIS	-	EXPRESSION TAG	UNP P70604
B	487	MET	-	EXPRESSION TAG	UNP P70604

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	O	0	0
			1	1		
2	B	3	Total	O	0	0
			3	3		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Small conductance calcium-activated potassium channel protein 2

Chain A: 



- Molecule 1: Small conductance calcium-activated potassium channel protein 2

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	29.20 Å 29.20 Å 192.90 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.10 25.29 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-2.10) 94.7 (25.29-2.10)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.61 (at 2.10 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.210 , 0.278 0.385 , 0.328	Depositor DCC
$R_{free}$ test set	240 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.5	Xtriage
Anisotropy	0.387	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 106.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.388 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.350 for h,-h-k,-l	Depositor
Outliers	0 of 5159 reflections	Xtriage
$F_o, F_c$ correlation	0.78	EDS
Total number of atoms	638	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/320	0.58	0/429
1	B	0.71	2/320 (0.6%)	0.91	3/429 (0.7%)
All	All	0.57	2/640 (0.3%)	0.76	3/858 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	525	LEU	C-O	-8.50	1.07	1.23
1	B	525	LEU	C-N	-6.74	1.21	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	525	LEU	O-C-N	9.62	139.39	121.10
1	B	525	LEU	CA-C-N	-7.29	96.70	117.10
1	B	525	LEU	CA-CB-CG	6.26	129.70	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	317	0	318	11	0
1	B	317	0	318	8	0
2	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	3	0	0	0	0
All	All	638	0	636	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:505:GLU:O	1:A:509:VAL:HG23	1.73	0.88
1:B:525:LEU:O	1:B:526:PRO:C	2.06	0.86
1:B:508:ILE:O	1:B:512:GLU:HG3	1.76	0.84
1:A:510:THR:HG22	1:A:514:LYS:HE3	1.66	0.77
1:A:504:PHE:O	1:A:508:ILE:HG12	1.85	0.76
1:B:525:LEU:O	1:B:526:PRO:OXT	2.11	0.68
1:B:521:SER:O	1:B:524:ALA:HB3	1.96	0.64
1:B:522:ILE:O	1:B:525:LEU:HD22	2.00	0.62
1:A:518:LEU:O	1:A:522:ILE:HG13	2.01	0.60
1:B:521:SER:O	1:B:525:LEU:HD13	2.03	0.58
1:B:512:GLU:O	1:B:516:GLU:HG3	2.09	0.51
1:A:525:LEU:HB3	1:A:526:PRO:HD2	1.94	0.50
1:A:510:THR:CG2	1:A:514:LYS:HE3	2.40	0.48
1:B:518:LEU:O	1:B:522:ILE:HG13	2.14	0.47
1:A:508:ILE:O	1:A:512:GLU:HG3	2.17	0.45
1:A:495:SER:HA	1:A:498:ASN:HD22	1.82	0.44
1:A:523:HIS:O	1:A:524:ALA:C	2.56	0.43
1:A:525:LEU:O	1:A:526:PRO:C	2.57	0.42
1:A:515:LEU:CD2	1:A:519:ILE:HD11	2.50	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	37/43 (86%)	33 (89%)	2 (5%)	2 (5%)	2	0
1	B	37/43 (86%)	36 (97%)	1 (3%)	0	100	100
All	All	74/86 (86%)	69 (93%)	3 (4%)	2 (3%)	5	1

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	524	ALA
1	A	515	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	37/40 (92%)	36 (97%)	1 (3%)	44	48
1	B	37/40 (92%)	34 (92%)	3 (8%)	11	8
All	All	74/80 (92%)	70 (95%)	4 (5%)	22	20

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	500	ARG
1	B	497	LEU
1	B	518	LEU
1	B	525	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	498	ASN



### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.