



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 06:28 am BST

PDB ID : 6PN4
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with 7-(3-(Aminomethyl)-4-(cyclopropylmethoxy)phenyl)-4-methylquinolin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2019-07-02
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>6%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>••</div> </div> </div>
1	B	422	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div>••</div> </div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7297 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	0	3	0
			3357	2149	573	613	22			
1	B	411	Total	C	N	O	S	0	3	0
			3353	2145	574	612	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 7-[3-(aminomethyl)-4-(cyclopropylmethoxy)phenyl]-4-methylquinolin-2-amine (three-letter code: OU1) (formula: C₂₁H₂₃N₃O) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	1
			26	21	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	1
			26	21	4	1		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	160	Total	O	0	0
			160	160		
7	B	246	Total	O	0	0
			246	246		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.76Å 111.65Å 164.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.32 – 1.90 92.32 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.7 (92.32-1.90) 97.4 (92.32-1.90)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	0.24	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 1.90Å)	Xtriage
Refinement program	PHENIX (1.11.1-2575_1496: ???)	Depositor
R, R_{free}	0.216 , 0.262 0.214 , 0.260	Depositor DCC
R_{free} test set	3670 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	1.119	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7297	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, OU1, H4B, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3457	0.52	0/4690
1	B	0.39	0/3456	0.53	0/4685
All	All	0.38	0/6913	0.52	0/9375

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3357	0	3266	35	0
1	B	3353	0	3270	22	0
2	A	43	0	30	8	0
2	B	43	0	30	3	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	26	0	0	2	0
4	B	26	0	0	0	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	160	0	0	4	0
7	B	246	0	0	4	0
All	All	7297	0	6632	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:801:HEM:O1D	7:A:901:HOH:O	1.94	0.84
2:B:801:HEM:HBB2	2:B:801:HEM:HHC	1.61	0.82
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.63	0.80
2:A:801:HEM:HHC	2:A:801:HEM:HBB2	1.68	0.76
1:B:692:HIS:ND1	7:B:901:HOH:O	2.06	0.73
1:A:667:ARG:NH1	1:A:668[A]:CYS:SG	2.64	0.69
1:B:414:ARG:HB2	2:B:801:HEM:HBD2	1.72	0.69
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.76	0.66
2:A:801:HEM:HBC2	2:A:801:HEM:HMC2	1.79	0.63
1:A:465:ARG:NH2	1:A:471:ASP:OD2	2.31	0.63
2:B:801:HEM:HBC2	2:B:801:HEM:HMC2	1.81	0.62
1:B:525:GLN:HE21	1:B:528:GLY:HA2	1.68	0.58
1:A:628:GLN:NE2	1:B:632:GLU:OE2	2.38	0.56
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.88	0.55
1:B:593:ILE:HA	1:B:597:ASP:HB2	1.90	0.54
1:A:330:ILE:HD11	1:B:696:LEU:HB3	1.90	0.53
1:A:391:THR:O	1:A:392:SER:OG	2.26	0.53
1:A:496:PRO:HA	1:A:499:VAL:HG23	1.90	0.52
1:A:598:TYR:HA	1:A:604:TYR:HB2	1.92	0.51
1:A:510:TRP:CE2	1:A:521:PRO:HD3	2.46	0.50
1:A:316:LEU:HD12	1:A:319:LYS:HE2	1.93	0.50
2:A:801:HEM:O2A	4:A:803[B]:OU1:N28	2.45	0.50
1:A:450:THR:HA	1:A:455:LEU:HD22	1.92	0.49
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.92	0.49
1:B:322:LEU:HD13	1:B:699:ARG:NH2	2.27	0.49
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.48	0.49
1:A:631:VAL:HG11	1:B:628:GLN:HG2	1.93	0.49
1:B:516:ARG:NH1	7:B:907:HOH:O	2.42	0.48
1:A:414:ARG:HB2	2:A:801:HEM:HAD1	1.97	0.47
1:B:686:SER:HA	1:B:691:PHE:CG	2.49	0.47
1:A:676:TRP:CE2	1:A:680:VAL:HG21	2.50	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:596:ARG:NH2	1:A:600:ASP:OD2	2.43	0.46
1:A:370:LYS:HE3	1:A:370:LYS:HB2	1.75	0.45
1:B:572:LEU:HB3	1:B:579:PHE:HB2	1.98	0.45
1:A:464:GLN:HB3	1:A:579:PHE:CE2	2.51	0.45
1:A:480:ILE:HD13	1:A:541:VAL:HG13	1.99	0.45
1:B:510:TRP:CE2	1:B:521:PRO:HD3	2.51	0.45
1:B:350:THR:OG1	1:B:353:GLN:OE1	2.27	0.45
1:A:548:HIS:CE1	1:A:550:LYS:HB2	2.52	0.44
1:A:455:LEU:HD12	1:A:587:TRP:HB3	2.00	0.44
1:A:706[A]:TYR:OH	2:A:801:HEM:O2D	2.30	0.44
1:A:676:TRP:CZ2	1:A:680:VAL:HG21	2.53	0.44
4:A:803[B]:OU1:N28	7:A:904:HOH:O	2.37	0.44
1:A:493:LEU:HA	7:A:903:HOH:O	2.18	0.43
2:A:801:HEM:HBD2	2:A:801:HEM:HMD2	2.00	0.43
1:A:588:TYR:CD1	1:A:593:ILE:HD11	2.54	0.43
1:A:430:ARG:NH1	7:A:912:HOH:O	2.46	0.43
1:B:596:ARG:O	1:B:600:ASP:HB2	2.19	0.43
2:A:801:HEM:CMD	2:A:801:HEM:HBD2	2.49	0.43
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.54	0.42
1:A:595:VAL:O	1:A:599:CYS:HB2	2.20	0.42
1:B:336:MET:HE2	1:B:336:MET:HB2	1.87	0.42
1:A:473:ARG:HD2	1:A:580:SER:HB2	2.02	0.42
1:A:382:GLU:HG3	1:A:383:GLU:N	2.33	0.42
1:A:593:ILE:O	1:A:597:ASP:HB2	2.20	0.42
1:B:573:GLU:OE1	7:B:902:HOH:O	2.22	0.41
1:A:714:HIS:CD2	1:A:716:TRP:HA	2.55	0.41
1:B:325:GLY:O	1:B:332:MET:HG3	2.21	0.41
1:A:328:GLU:H	1:A:328:GLU:CD	2.25	0.40
1:B:559:LEU:HD11	1:B:606:ILE:HD13	2.03	0.40
1:B:440:ASN:ND2	7:B:909:HOH:O	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/422 (97%)	392 (96%)	15 (4%)	2 (0%)	29	18
1	B	410/422 (97%)	400 (98%)	10 (2%)	0	100	100
All	All	819/844 (97%)	792 (97%)	25 (3%)	2 (0%)	47	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	338	PRO
1	A	715	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/377 (98%)	355 (96%)	14 (4%)	33	24
1	B	369/377 (98%)	360 (98%)	9 (2%)	49	43
All	All	738/754 (98%)	715 (97%)	23 (3%)	40	32

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	337	LEU
1	A	382	GLU
1	A	469	LYS
1	A	508	GLN
1	A	511	LYS
1	A	523	LEU
1	A	547	ARG
1	A	555	LYS
1	A	608	GLU
1	A	609	GLU
1	A	620	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	645	LYS
1	A	663	GLU
1	A	715	VAL
1	B	332	MET
1	B	351	LYS
1	B	390	SER
1	B	392	SER
1	B	516	ARG
1	B	540	LEU
1	B	547	ARG
1	B	573	GLU
1	B	608	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	801	1	27,50,50	1.85	5 (18%)	17,82,82	2.16	8 (47%)
5	ACT	A	804	-	1,3,3	1.78	0	0,3,3	0.00	-
5	ACT	B	804	-	1,3,3	1.41	0	0,3,3	0.00	-
4	OU1	B	803[A]	-	28,28,28	0.93	1 (3%)	39,40,40	1.80	10 (25%)
4	OU1	A	803[B]	-	28,28,28	0.98	1 (3%)	39,40,40	1.64	7 (17%)
4	OU1	B	803[B]	-	28,28,28	0.93	1 (3%)	39,40,40	1.80	10 (25%)
2	HEM	B	801	1	27,50,50	1.95	4 (14%)	17,82,82	2.37	7 (41%)
3	H4B	B	802	-	16,18,18	0.80	0	11,26,26	2.52	5 (45%)
3	H4B	A	802	-	16,18,18	0.87	0	11,26,26	2.61	7 (63%)
4	OU1	A	803[A]	-	28,28,28	0.98	1 (3%)	39,40,40	1.64	7 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	801	1	-	4/6/54/54	-
4	OU1	B	803[A]	-	-	3/11/13/13	0/4/4/4
4	OU1	A	803[B]	-	-	4/11/13/13	0/4/4/4
4	OU1	B	803[B]	-	-	3/11/13/13	0/4/4/4
2	HEM	B	801	1	-	1/6/54/54	-
3	H4B	B	802	-	-	2/8/17/17	0/2/2/2
3	H4B	A	802	-	-	3/8/17/17	0/2/2/2
4	OU1	A	803[A]	-	-	3/11/13/13	0/4/4/4

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	HEM	C3B-C2B	-4.95	1.33	1.40
2	A	801	HEM	C3B-C2B	-4.73	1.33	1.40
2	B	801	HEM	C3B-CAB	3.67	1.55	1.47
2	B	801	HEM	C3C-C2C	-3.62	1.35	1.40
2	A	801	HEM	C3C-C2C	-3.61	1.35	1.40
2	B	801	HEM	C3C-CAC	3.33	1.54	1.47
2	A	801	HEM	C3B-CAB	3.31	1.54	1.47
2	A	801	HEM	C3C-CAC	3.15	1.54	1.47
4	A	803[B]	OU1	C02-N01	2.92	1.37	1.33
4	A	803[A]	OU1	C02-N01	2.92	1.37	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEM	CMC-C2C	2.40	1.57	1.51
4	B	803[A]	OU1	C05-C10	-2.19	1.38	1.42
4	B	803[B]	OU1	C05-C10	-2.19	1.38	1.42

All (61) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	HEM	CBD-CAD-C3D	-5.72	101.93	112.48
3	B	802	H4B	C4-C4A-C8A	4.90	118.92	114.57
3	A	802	H4B	C4-C4A-C8A	4.64	118.69	114.57
4	B	803[A]	OU1	C04-C05-C10	4.52	120.46	118.01
4	B	803[B]	OU1	C04-C05-C10	4.52	120.46	118.01
4	A	803[B]	OU1	C04-C05-C10	4.44	120.42	118.01
4	A	803[A]	OU1	C04-C05-C10	4.44	120.42	118.01
4	A	803[B]	OU1	C26-C21-C08	-4.44	113.53	120.86
4	A	803[A]	OU1	C26-C21-C08	-4.44	113.53	120.86
4	B	803[A]	OU1	C26-C21-C08	-3.91	114.39	120.86
4	B	803[B]	OU1	C26-C21-C08	-3.91	114.39	120.86
2	B	801	HEM	CAD-CBD-CGD	3.73	118.92	112.67
4	B	803[A]	OU1	C05-C10-N01	-3.64	118.95	122.81
4	B	803[B]	OU1	C05-C10-N01	-3.64	118.95	122.81
3	B	802	H4B	C4-N3-C2	3.63	121.69	115.93
2	A	801	HEM	CBD-CAD-C3D	3.34	118.63	112.48
2	B	801	HEM	CBA-CAA-C2A	-3.33	106.34	112.49
2	A	801	HEM	CMD-C2D-C1D	-3.27	123.44	128.46
4	B	803[A]	OU1	C08-C09-C10	-3.25	118.60	121.44
4	B	803[B]	OU1	C08-C09-C10	-3.25	118.60	121.44
3	A	802	H4B	N3-C2-N1	-3.22	120.38	125.42
2	A	801	HEM	CMA-C3A-C4A	-3.19	123.56	128.46
2	B	801	HEM	CMC-C2C-C3C	3.09	130.46	124.68
4	A	803[B]	OU1	C22-C21-C08	3.03	126.60	121.36
4	A	803[A]	OU1	C22-C21-C08	3.03	126.60	121.36
3	B	802	H4B	N3-C2-N1	-3.00	120.71	125.42
3	A	802	H4B	C4-N3-C2	3.00	120.70	115.93
3	A	802	H4B	N2-C2-N3	2.96	121.86	117.25
3	A	802	H4B	C4-C4A-N5	2.93	121.58	119.12
2	A	801	HEM	CMC-C2C-C3C	2.92	130.15	124.68
4	A	803[B]	OU1	C05-C10-N01	-2.85	119.79	122.81
4	A	803[A]	OU1	C05-C10-N01	-2.85	119.79	122.81
3	A	802	H4B	C2-N1-C8A	2.81	120.83	114.54
4	B	803[A]	OU1	C06-C05-C04	-2.72	118.46	123.66
4	B	803[B]	OU1	C06-C05-C04	-2.72	118.46	123.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	803[A]	OU1	N02-C02-N01	2.72	120.51	118.26
4	B	803[B]	OU1	N02-C02-N01	2.72	120.51	118.26
2	B	801	HEM	CMA-C3A-C4A	-2.72	124.29	128.46
2	A	801	HEM	CAD-CBD-CGD	-2.59	108.32	112.67
4	B	803[A]	OU1	C06-C05-C10	2.46	121.08	118.33
4	B	803[B]	OU1	C06-C05-C10	2.46	121.08	118.33
3	B	802	H4B	C4-C4A-N5	2.46	121.18	119.12
4	B	803[A]	OU1	O29-C24-C25	2.37	118.88	115.78
4	B	803[B]	OU1	O29-C24-C25	2.37	118.88	115.78
3	A	802	H4B	C4A-N5-C6	-2.36	114.75	121.16
4	B	803[A]	OU1	C22-C21-C08	2.35	125.42	121.36
4	B	803[B]	OU1	C22-C21-C08	2.35	125.42	121.36
4	A	803[B]	OU1	C06-C05-C04	-2.28	119.30	123.66
4	A	803[A]	OU1	C06-C05-C04	-2.28	119.30	123.66
2	B	801	HEM	C1D-C2D-C3D	2.28	108.58	107.00
3	B	802	H4B	C2-N1-C8A	2.27	119.62	114.54
4	A	803[B]	OU1	N02-C02-N01	2.25	120.12	118.26
4	A	803[A]	OU1	N02-C02-N01	2.25	120.12	118.26
2	A	801	HEM	CBA-CAA-C2A	-2.21	108.42	112.49
2	A	801	HEM	C4C-C3C-C2C	2.20	108.43	106.90
2	B	801	HEM	CMD-C2D-C1D	-2.18	125.11	128.46
4	B	803[A]	OU1	C33-C31-C30	-2.16	107.67	119.43
4	B	803[B]	OU1	C33-C31-C30	-2.16	107.67	119.43
4	A	803[B]	OU1	C32-C31-C30	-2.03	108.37	119.43
4	A	803[A]	OU1	C32-C31-C30	-2.03	108.37	119.43
2	A	801	HEM	CAA-CBA-CGA	-2.02	109.28	112.67

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	HEM	C2A-CAA-CBA-CGA
2	A	801	HEM	C2D-C3D-CAD-CBD
2	A	801	HEM	C4D-C3D-CAD-CBD
2	A	801	HEM	C3D-CAD-CBD-CGD
2	B	801	HEM	C2A-CAA-CBA-CGA
3	A	802	H4B	C7-C6-C9-C10
4	A	803[B]	OU1	C31-C30-O29-C24
4	A	803[A]	OU1	C31-C30-O29-C24
4	B	803[A]	OU1	C24-C25-C27-N28
4	A	803[B]	OU1	C24-C25-C27-N28
4	B	803[A]	OU1	O29-C30-C31-C32

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	B	803[B]	OU1	O29-C30-C31-C32
3	A	802	H4B	C7-C6-C9-O9
4	B	803[B]	OU1	C26-C25-C27-N28
3	B	802	H4B	C7-C6-C9-C10
4	A	803[B]	OU1	C07-C08-C21-C26
4	A	803[A]	OU1	C07-C08-C21-C26
3	B	802	H4B	N5-C6-C9-O9
3	A	802	H4B	N5-C6-C9-O9
4	B	803[A]	OU1	O29-C30-C31-C33
4	A	803[B]	OU1	O29-C30-C31-C33
4	B	803[B]	OU1	O29-C30-C31-C33
4	A	803[A]	OU1	O29-C30-C31-C33



There are no ring outliers.


3 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	HEM	8	0
4	A	803[B]	OU1	2	0
2	B	801	HEM	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	411/422 (97%)	0.41	24 (5%) 23 25	23, 54, 105, 150	0
1	B	411/422 (97%)	0.12	6 (1%) 73 76	22, 41, 78, 113	0
All	All	822/844 (97%)	0.27	30 (3%) 42 45	22, 47, 98, 150	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PHE	6.7
1	A	322	LEU	4.4
1	A	604	TYR	4.3
1	A	716	TRP	4.2
1	A	488	PRO	4.1
1	A	373	GLY	4.0
1	B	348	VAL	3.9
1	A	348	VAL	3.9
1	A	553	TRP	3.6
1	A	347	ASP	3.4
1	A	339	SER	3.2
1	B	299	ARG	3.1
1	A	599	CYS	3.1
1	A	706[A]	TYR	2.9
1	A	715	VAL	2.9
1	A	349	ARG	2.9
1	A	300	PHE	2.9
1	B	352	ASP	2.8
1	B	350	THR	2.7
1	A	602	SER	2.6
1	B	351	LYS	2.5
1	A	486	LYS	2.4
1	A	595	VAL	2.4
1	A	382	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	597	ASP	2.3
1	A	506	ILE	2.2
1	A	552	ASP	2.2
1	A	507	GLN	2.2
1	A	551	PHE	2.1
1	A	299	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	H4B	B	802	17/17	0.90	0.12	29,37,42,48	0
4	OU1	B	803[B]	25/25	0.91	0.15	28,35,56,61	1
4	OU1	B	803[A]	25/25	0.91	0.15	28,35,56,61	1
3	H4B	A	802	17/17	0.92	0.15	28,42,52,55	0
5	ACT	B	804	4/4	0.93	0.15	63,66,66,69	0
4	OU1	A	803[B]	25/25	0.94	0.16	26,40,83,87	1
5	ACT	A	804	4/4	0.94	0.17	55,55,57,58	0
4	OU1	A	803[A]	25/25	0.94	0.16	26,40,83,87	1
2	HEM	B	801	43/43	0.97	0.13	20,32,54,63	0
2	HEM	A	801	43/43	0.97	0.12	20,30,58,66	0
6	ZN	B	805	1/1	0.99	0.12	35,35,35,35	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OU1 B 803 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OU1 B 803 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OU1 A 803 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OU1 A 803 (A):**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.