



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 02:30 pm BST

PDB ID : 6PNB
Title : Structure of human neuronal nitric oxide synthase R354A/G357D mutant heme domain in complex with 7-(4-(Aminomethyl)phenyl)-4-methylquinolin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2019-07-02
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 86%, grey 11%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 86% 11% .. </div> </div>
1	B	421	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 4%, orange 1%, yellow 1%, green 86%, grey 11%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 4% 86% 11% . </div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7466 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	414	Total	C	N	O	S	0	3	0
			3383	2166	575	619	23			
1	B	409	Total	C	N	O	S	0	2	0
			3343	2142	570	610	21			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	354	ALA	ARG	engineered mutation	UNP P29475
A	357	ASP	GLY	engineered mutation	UNP P29475
B	354	ALA	ARG	engineered mutation	UNP P29475
B	357	ASP	GLY	engineered mutation	UNP P29475

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	B	1	Total	C	Fe	N	O	
			43	34	1	4	4	

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O		
			17	9	5	3		
3	B	1	Total	C	N	O		
			17	9	5	3		

- Molecule 4 is 7-[4-(aminomethyl)phenyl]-4-methylquinolin-2-amine (three-letter code: OSY) (formula: $C_{17}H_{17}N_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			20	17	3		
4	B	1	Total	C	N	0	0
			20	17	3		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total 1	Zn 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	314	Total 314	O 314	0	0
7	B	253	Total 253	O 253	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.36Å 123.21Å 165.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.36 – 2.05 49.36 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.6 (49.36-2.05) 99.8 (49.36-2.05)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.05Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.185 , 0.233 0.184 , 0.233	Depositor DCC
R_{free} test set	3327 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.688	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7466	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OSY, GOL, ZN, H4B, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/3488	0.51	0/4732
1	B	0.36	0/3444	0.48	0/4671
All	All	0.37	0/6932	0.49	0/9403

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3383	0	3293	31	0
1	B	3343	0	3257	28	0
2	A	43	0	30	4	0
2	B	43	0	30	2	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	20	0	0	2	0
4	B	20	0	0	1	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	314	0	0	2	0
7	B	253	0	0	2	0
All	All	7466	0	6656	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:528:LEU:HD22	1:A:536:PRO:HB2	1.58	0.85
1:B:674:ARG:NH2	7:B:901:HOH:O	2.19	0.74
1:B:528:LEU:HD22	1:B:536:PRO:HB2	1.69	0.72
1:A:610:ASN:O	7:A:901:HOH:O	2.09	0.69
2:B:801:HEM:HBB2	2:B:801:HEM:HHC	1.76	0.68
1:A:711:TYR:OH	2:A:801:HEM:O2D	2.12	0.67
1:B:572:VAL:HG21	4:B:803:OSY:C07	2.33	0.58
2:A:801:HEM:HBC2	2:A:801:HEM:HMC2	1.84	0.58
1:B:711:TYR:OH	2:B:801:HEM:O2D	2.22	0.58
1:A:326:THR:OG1	1:A:704:ARG:NH1	2.39	0.56
1:A:491:LYS:HG2	1:A:504:VAL:HG11	1.88	0.55
1:A:480:TRP:HB2	1:A:528:LEU:HB3	1.89	0.55
1:B:483:GLN:HB2	1:B:486:ARG:HG3	1.89	0.54
1:A:490:TYR:CE1	1:A:519:ARG:HA	2.42	0.54
1:A:326:THR:HG21	1:A:706:THR:HG22	1.89	0.54
1:A:555:LYS:NZ	7:A:910:HOH:O	2.41	0.53
1:B:390:ASN:ND2	7:B:907:HOH:O	2.42	0.53
1:A:636:VAL:HG11	1:B:633:GLN:HG3	1.90	0.53
1:B:480:TRP:HB2	1:B:528:LEU:HB3	1.90	0.52
2:A:801:HEM:HHC	2:A:801:HEM:HBB2	1.91	0.52
1:A:552:ARG:NH1	1:A:649:ASP:OD1	2.28	0.52
1:A:483:GLN:HB2	1:A:486:ARG:HG3	1.92	0.51
1:B:414:TRP:CE3	1:B:426:TRP:HA	2.46	0.50
1:B:327:LEU:HB2	1:B:704:ARG:HG2	1.93	0.50
1:A:633:GLN:NE2	1:B:637:GLU:OE2	2.44	0.48
1:A:491:LYS:HD2	1:A:491:LYS:HA	1.68	0.48
1:B:501:PRO:HA	1:B:504:VAL:HG23	1.95	0.48
1:A:414:TRP:CE3	1:A:426:TRP:HA	2.49	0.48
1:A:667:MET:HE1	1:A:698:GLN:HG2	1.97	0.47
1:A:572:VAL:HG21	4:A:803:OSY:C07	2.45	0.47
1:A:327:LEU:HB2	1:A:704:ARG:HH11	1.80	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:509:ILE:O	1:B:513:GLN:HG2	2.15	0.46
1:B:356:LYS:HG2	1:B:397:SER:HA	1.96	0.46
1:A:603:TYR:O	1:A:610:ASN:N	2.49	0.45
1:B:598:ILE:HA	1:B:602:ASP:OD2	2.16	0.45
1:A:509:ILE:O	1:A:513:GLN:HG2	2.17	0.45
1:B:363:ALA:O	1:B:367:ILE:HG12	2.17	0.44
1:A:478:ARG:HD2	1:A:585:SER:HB2	1.98	0.44
1:A:680:ASP:O	1:A:684:ILE:HG12	2.18	0.44
1:B:490:TYR:CE1	1:B:519:ARG:HA	2.53	0.43
1:A:691:SER:HA	1:A:696:PHE:CG	2.54	0.43
1:A:327:LEU:HD13	1:A:328:GLU:H	1.83	0.43
1:B:685:VAL:HA	1:B:686:PRO:HD3	1.88	0.43
1:B:689:SER:HB3	1:B:692:ILE:HD11	2.00	0.43
1:A:600:VAL:O	1:A:604:CYS:HB2	2.19	0.42
1:B:598:ILE:O	1:B:602:ASP:HB2	2.19	0.42
1:A:327:LEU:HB2	1:A:704:ARG:NH1	2.35	0.42
1:B:491:LYS:HA	1:B:491:LYS:HD2	1.95	0.42
1:B:327:LEU:HB2	1:B:704:ARG:CG	2.49	0.41
1:B:600:VAL:O	1:B:604:CYS:HB2	2.21	0.41
1:A:327:LEU:HB2	1:A:704:ARG:HD3	2.02	0.41
1:B:478:ARG:NH2	1:B:715:PRO:HD3	2.35	0.41
1:A:689:SER:HB3	1:A:692:ILE:HD11	2.03	0.41
1:B:613:GLU:H	1:B:613:GLU:HG3	1.64	0.41
1:A:667:MET:HE1	1:A:698:GLN:CD	2.41	0.41
1:A:593:TYR:CD1	1:A:598:ILE:HD11	2.56	0.41
1:B:605:ASP:HB2	1:B:608:ARG:CG	2.51	0.41
1:A:598:ILE:HA	1:A:602:ASP:HB2	2.03	0.40
2:A:801:HEM:HBD1	4:A:803:OSY:C22	2.50	0.40
1:B:691:SER:HA	1:B:696:PHE:CG	2.55	0.40
1:B:467:PHE:HB2	1:B:586:ALA:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	413/421 (98%)	399 (97%)	12 (3%)	2 (0%)	29	18
1	B	407/421 (97%)	397 (98%)	10 (2%)	0	100	100
All	All	820/842 (97%)	796 (97%)	22 (3%)	2 (0%)	47	39

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	352	ASP
1	A	353	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/376 (99%)	361 (97%)	12 (3%)	39	32
1	B	367/376 (98%)	360 (98%)	7 (2%)	57	53
All	All	740/752 (98%)	721 (97%)	19 (3%)	49	39

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	327	LEU
1	A	333	GLU
1	A	351	GLU
1	A	386	LEU
1	A	390	ASN
1	A	394	ASP
1	A	491	LYS
1	A	521	ARG
1	A	528	LEU
1	A	552	ARG
1	A	673[A]	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	673[B]	CYS
1	B	332	THR
1	B	387	GLU
1	B	528	LEU
1	B	552	ARG
1	B	555	LYS
1	B	720	VAL
1	B	722	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	697	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	804	-	5,5,5	0.38	0	5,5,5	0.25	0
2	HEM	B	801	1	27,50,50	2.20	5 (18%)	17,82,82	2.16	6 (35%)
2	HEM	A	801	1	27,50,50	2.19	6 (22%)	17,82,82	1.97	5 (29%)
3	H4B	A	802	-	16,18,18	0.84	0	11,26,26	2.60	6 (54%)
3	H4B	B	802	-	16,18,18	0.92	0	11,26,26	2.58	6 (54%)
4	OSY	B	803	-	22,22,22	1.03	2 (9%)	30,31,31	1.42	4 (13%)
4	OSY	A	803	-	22,22,22	1.03	1 (4%)	30,31,31	1.42	5 (16%)
5	GOL	A	804	-	5,5,5	0.40	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	804	-	-	4/4/4/4	-
2	HEM	B	801	1	-	2/6/54/54	-
2	HEM	A	801	1	-	2/6/54/54	-
3	H4B	A	802	-	-	3/8/17/17	0/2/2/2
3	H4B	B	802	-	-	2/8/17/17	0/2/2/2
4	OSY	B	803	-	-	4/6/6/6	0/3/3/3
4	OSY	A	803	-	-	0/6/6/6	0/3/3/3
5	GOL	A	804	-	-	4/4/4/4	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	HEM	C3D-C2D	5.54	1.54	1.37
2	A	801	HEM	C3D-C2D	5.48	1.53	1.37
2	A	801	HEM	C3B-C2B	-5.20	1.33	1.40
2	B	801	HEM	C3B-C2B	-4.68	1.33	1.40
2	B	801	HEM	C3C-C2C	-3.98	1.34	1.40
2	B	801	HEM	C3B-CAB	3.88	1.55	1.47
2	A	801	HEM	C3C-C2C	-3.86	1.35	1.40
2	A	801	HEM	C3B-CAB	3.53	1.55	1.47
2	B	801	HEM	C3C-CAC	3.50	1.55	1.47
2	A	801	HEM	C3C-CAC	3.23	1.54	1.47
4	A	803	OSY	C05-C10	-2.42	1.38	1.42
2	A	801	HEM	C1D-ND	2.22	1.40	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	803	OSY	C05-C10	-2.12	1.39	1.42
4	B	803	OSY	C04-C05	-2.01	1.38	1.42

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	802	H4B	C4-C4A-C8A	5.48	119.44	114.57
2	B	801	HEM	CBA-CAA-C2A	-5.41	102.51	112.49
2	A	801	HEM	CBA-CAA-C2A	-4.97	103.32	112.49
3	A	802	H4B	C4-C4A-C8A	4.90	118.93	114.57
4	B	803	OSY	C26-C21-C08	-3.65	115.04	121.36
3	A	802	H4B	N3-C2-N1	-3.37	120.14	125.42
4	A	803	OSY	C26-C21-C08	-3.31	115.63	121.36
2	B	801	HEM	CAD-CBD-CGD	-3.24	107.24	112.67
3	A	802	H4B	C4-C4A-N5	3.18	121.79	119.12
3	B	802	H4B	N3-C2-N1	-3.15	120.48	125.42
4	A	803	OSY	C03-C04-C05	3.08	120.81	117.78
4	A	803	OSY	C05-C10-N01	-3.02	119.61	122.81
2	A	801	HEM	CAD-CBD-CGD	-3.02	107.61	112.67
2	A	801	HEM	C1D-C2D-C3D	-3.00	104.91	107.00
3	B	802	H4B	C4-N3-C2	2.97	120.65	115.93
3	A	802	H4B	C4-N3-C2	2.94	120.60	115.93
4	B	803	OSY	C04-C05-C10	2.87	119.56	118.01
2	B	801	HEM	C1D-C2D-C3D	-2.85	105.01	107.00
4	B	803	OSY	C05-C10-N01	-2.82	119.82	122.81
3	A	802	H4B	C2-N1-C8A	2.81	120.84	114.54
3	B	802	H4B	C2-N1-C8A	2.74	120.67	114.54
2	A	801	HEM	CMC-C2C-C3C	2.54	129.43	124.68
2	B	801	HEM	CMA-C3A-C4A	-2.53	124.57	128.46
4	A	803	OSY	C11-C04-C05	-2.38	117.39	121.08
4	B	803	OSY	C22-C21-C08	2.33	125.39	121.36
2	A	801	HEM	CMA-C3A-C4A	-2.25	125.01	128.46
3	B	802	H4B	C4-C4A-N5	2.14	120.92	119.12
3	B	802	H4B	C4A-N5-C6	-2.10	115.44	121.16
4	A	803	OSY	C22-C21-C08	2.08	124.96	121.36
2	B	801	HEM	CMC-C2C-C3C	2.05	128.52	124.68
2	B	801	HEM	CMD-C2D-C3D	2.02	128.75	124.94
3	A	802	H4B	N2-C2-N1	2.01	120.38	117.25

There are no chirality outliers.

All (21) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	804	GOL	O1-C1-C2-C3
5	B	804	GOL	C1-C2-C3-O3
2	B	801	HEM	C2D-C3D-CAD-CBD
2	B	801	HEM	C4D-C3D-CAD-CBD
2	A	801	HEM	C2D-C3D-CAD-CBD
2	A	801	HEM	C4D-C3D-CAD-CBD
5	A	804	GOL	O1-C1-C2-C3
4	B	803	OSY	C07-C08-C21-C26
4	B	803	OSY	C07-C08-C21-C22
4	B	803	OSY	C09-C08-C21-C26
4	B	803	OSY	C09-C08-C21-C22
5	B	804	GOL	O1-C1-C2-O2
5	A	804	GOL	O1-C1-C2-O2
5	B	804	GOL	O2-C2-C3-O3
3	A	802	H4B	C7-C6-C9-O9
5	A	804	GOL	O2-C2-C3-O3
3	A	802	H4B	C7-C6-C9-C10
5	A	804	GOL	C1-C2-C3-O3
3	B	802	H4B	C7-C6-C9-C10
3	A	802	H4B	N5-C6-C9-O9
3	B	802	H4B	N5-C6-C9-O9


There are no ring outliers.


4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	HEM	2	0
2	A	801	HEM	4	0
4	B	803	OSY	1	0
4	A	803	OSY	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	414/421 (98%)	0.09	5 (1%) 79 81	27, 41, 73, 122	0
1	B	409/421 (97%)	0.15	17 (4%) 36 39	28, 47, 89, 116	0
All	All	823/842 (97%)	0.12	22 (2%) 54 59	27, 44, 84, 122	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	302	CYS	7.2
1	B	354	ALA	4.5
1	B	357	ASP	3.8
1	A	353	VAL	3.7
1	B	327	LEU	3.6
1	B	720	VAL	3.6
1	B	326	THR	3.3
1	B	304	ARG	3.2
1	A	352	ASP	3.0
1	B	358	GLN	3.0
1	A	334	TYR	3.0
1	B	383	MET	2.6
1	B	721	TRP	2.6
1	B	600	VAL	2.6
1	B	386	LEU	2.6
1	B	360	PHE	2.5
1	B	624	ARG	2.5
1	B	355	THR	2.3
1	B	377	PHE	2.3
1	B	722	LYS	2.3
1	B	374	ILE	2.2
1	A	620	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	B	804	6/6	0.74	0.33	69,77,86,100	0
5	GOL	A	804	6/6	0.78	0.17	59,67,73,74	0
4	OSY	A	803	20/20	0.92	0.18	28,41,72,79	0
3	H4B	A	802	17/17	0.93	0.16	36,46,51,53	0
4	OSY	B	803	20/20	0.93	0.15	27,42,72,74	0
3	H4B	B	802	17/17	0.94	0.15	32,51,66,67	0
2	HEM	B	801	43/43	0.97	0.12	25,33,50,63	0
2	HEM	A	801	43/43	0.98	0.14	23,30,55,66	0
6	ZN	A	805	1/1	1.00	0.12	37,37,37,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OSY A 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OSY B 803:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.