



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 07:25 pm BST

PDB ID : 6PNH
Title : Structure of human neuronal nitric oxide synthase R354A/G357D mutant heme domain in complex with 7-(3-(Aminomethyl)-4-isopropoxyphenyl)-4-methylquinolin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2019-07-02
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>6% ..</div> </div> </div>
1	B	421	<div> <div>6%</div> <div> <div></div> <div>89%</div> <div>7% .</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7423 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	412	Total	C	N	O	S	0	2	0
			3365	2156	573	615	21			
1	B	409	Total	C	N	O	S	0	2	0
			3343	2142	570	610	21			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	354	ALA	ARG	engineered mutation	UNP P29475
A	357	ASP	GLY	engineered mutation	UNP P29475
B	354	ALA	ARG	engineered mutation	UNP P29475
B	357	ASP	GLY	engineered mutation	UNP P29475

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	B	1	Total	C	Fe	N	O	
			43	34	1	4	4	

- Molecule 3 is 7-{3-(aminomethyl)-4-[(propan-2-yl)oxy]phenyl}-4-methylquinolin-2-amine (three-letter code: OSD) (formula: C₂₀H₂₃N₃O) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O		
			24	20	3	1		
3	A	1	Total	C	N	O		
			24	20	3	1		
3	B	1	Total	C	N	O		
			24	20	3	1		
3	B	1	Total	C	N	O		
			24	20	3	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn		
			1	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	287	Total 287	O 287	0	0
5	B	245	Total 245	O 245	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.18Å 125.17Å 165.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.75 – 1.85 49.75 – 1.85	Depositor EDS
% Data completeness (in resolution range)	95.8 (49.75-1.85) 96.6 (49.75-1.85)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 1.86Å)	Xtriage
Refinement program	PHENIX (1.11.1-2575_1496: ???)	Depositor
R, R_{free}	0.202 , 0.246 0.203 , 0.248	Depositor DCC
R_{free} test set	4450 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	24.3	Xtriage
Anisotropy	1.153	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7423	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, OSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/3467	0.53	1/4703 (0.0%)
1	B	0.34	0/3444	0.49	0/4671
All	All	0.35	0/6911	0.51	1/9374 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	327	LEU	CA-CB-CG	5.15	127.14	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3365	0	3278	16	0
1	B	3343	0	3257	16	0
2	A	43	0	30	2	0
2	B	43	0	30	3	0
3	A	48	0	0	2	0
3	B	48	0	0	3	0
4	B	1	0	0	0	0
5	A	287	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	245	0	0	2	0
All	All	7423	0	6595	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:528:LEU:HD22	1:A:536:PRO:HB2	1.60	0.84
2:B:803:HEM:HBB2	2:B:803:HEM:HHC	1.68	0.76
1:A:601:ARG:NH2	1:A:602:ASP:OD1	2.29	0.66
1:A:602:ASP:OD2	5:A:901:HOH:O	2.12	0.66
1:A:430:GLN:OE1	5:A:902:HOH:O	2.15	0.65
2:A:801:HEM:HMC2	2:A:801:HEM:HBC2	1.80	0.62
1:B:528:LEU:HD22	1:B:536:PRO:HB2	1.85	0.58
1:B:606:ASN:HD21	1:B:609:TYR:HB2	1.69	0.57
2:A:801:HEM:HHC	2:A:801:HEM:HBB2	1.86	0.57
1:B:483:GLN:HB2	1:B:486:ARG:HG3	1.87	0.56
1:B:480:TRP:HB2	1:B:528:LEU:HB3	1.89	0.53
1:B:341:MET:HE2	1:B:683:TRP:HZ2	1.77	0.50
1:A:483:GLN:HB2	1:A:486:ARG:HG3	1.96	0.47
1:A:414:TRP:CE3	1:A:426:TRP:HA	2.50	0.47
1:A:697:HIS:ND1	5:A:907:HOH:O	2.35	0.46
3:B:804:OSD:O29	3:B:804:OSD:N28	2.49	0.46
1:B:691:SER:HA	1:B:696:PHE:CG	2.50	0.46
2:B:803:HEM:HBC2	2:B:803:HEM:HMC2	1.98	0.46
1:B:601:ARG:NH2	1:B:602:ASP:OD1	2.49	0.45
1:B:552:ARG:HB2	1:B:560:LYS:HE2	1.99	0.45
1:B:341:MET:HG2	1:B:342:HIS:CE1	2.52	0.45
1:A:623:MET:HA	1:A:630:TRP:CD1	2.53	0.43
1:A:469:GLN:HB3	1:A:584:PHE:CE2	2.54	0.43
1:B:325:SER:HB2	1:B:704:ARG:O	2.19	0.43
1:B:480:TRP:CE2	1:B:715:PRO:HB2	2.52	0.43
1:A:625:LYS:HE3	1:A:625:LYS:HB2	1.79	0.43
1:A:689:SER:HB3	1:A:692:ILE:HD11	2.00	0.42
1:B:414:TRP:CE3	1:B:426:TRP:HA	2.54	0.42
1:A:555:LYS:HD3	1:A:555:LYS:HA	1.87	0.42
1:B:572:VAL:HG21	3:B:804:OSD:C07	2.50	0.42
2:B:803:HEM:HHC	2:B:803:HEM:CBB	2.45	0.42
1:A:408:TYR:CE1	1:A:412:HIS:CE1	3.08	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:441:HIS:NE2	5:B:903:HOH:O	2.36	0.41
1:A:480:TRP:HB2	1:A:528:LEU:HB3	2.02	0.41
3:B:801:OSD:N28	5:B:904:HOH:O	2.37	0.41
3:A:803:OSD:C07	1:B:341:MET:HE3	2.51	0.41
1:A:572:VAL:HG21	3:A:802:OSD:C07	2.52	0.40
1:B:360:PHE:N	1:B:361:PRO:HD2	2.37	0.40
1:A:540:GLN:HG3	5:A:1113:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/421 (97%)	403 (98%)	7 (2%)	0	100	100
1	B	407/421 (97%)	396 (97%)	10 (2%)	1 (0%)	47	33
All	All	817/842 (97%)	799 (98%)	17 (2%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	495	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/376 (98%)	361 (98%)	9 (2%)	49	33
1	B	367/376 (98%)	358 (98%)	9 (2%)	47	31
All	All	737/752 (98%)	719 (98%)	18 (2%)	47	33

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	327	LEU
1	A	342	HIS
1	A	394	ASP
1	A	528	LEU
1	A	552	ARG
1	A	606	ASN
1	A	625	LYS
1	A	673	CYS
1	A	722	LYS
1	B	365	GLU
1	B	491	LYS
1	B	528	LEU
1	B	552	ARG
1	B	555	LYS
1	B	620	ASN
1	B	668	GLU
1	B	718	THR
1	B	720	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OSD	A	802	-	26,26,26	0.95	0	36,37,37	1.05	3 (8%)
3	OSD	B	804	-	26,26,26	0.96	0	36,37,37	1.25	3 (8%)
3	OSD	A	803	-	26,26,26	0.94	1 (3%)	36,37,37	1.35	5 (13%)
2	HEM	B	803	1	27,50,50	1.97	5 (18%)	17,82,82	2.15	7 (41%)
2	HEM	A	801	1	27,50,50	1.89	4 (14%)	17,82,82	1.88	4 (23%)
3	OSD	B	801	-	26,26,26	1.04	2 (7%)	36,37,37	1.76	7 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OSD	A	802	-	-	3/10/10/10	0/3/3/3
3	OSD	B	804	-	-	1/10/10/10	0/3/3/3
3	OSD	A	803	-	-	2/10/10/10	0/3/3/3
2	HEM	B	803	1	-	0/6/54/54	-
2	HEM	A	801	1	-	0/6/54/54	-
3	OSD	B	801	-	-	0/10/10/10	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	803	HEM	C3B-C2B	-5.16	1.33	1.40
2	A	801	HEM	C3B-C2B	-4.91	1.33	1.40
2	B	803	HEM	C3B-CAB	3.93	1.56	1.47
2	B	803	HEM	C3C-C2C	-3.87	1.35	1.40
2	A	801	HEM	C3B-CAB	3.63	1.55	1.47
2	A	801	HEM	C3C-C2C	-3.37	1.35	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEM	C3C-CAC	3.24	1.54	1.47
2	B	803	HEM	C3C-CAC	3.18	1.54	1.47
3	B	801	OSD	C02-N01	3.01	1.37	1.33
2	B	803	HEM	CAA-C2A	2.17	1.55	1.52
3	A	803	OSD	C02-N01	2.08	1.36	1.33
3	B	801	OSD	C04-C05	-2.07	1.38	1.42

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	803	HEM	CMA-C3A-C4A	-4.84	121.02	128.46
3	B	801	OSD	C24-O29-C30	4.59	128.38	119.53
3	B	801	OSD	C04-C05-C10	3.70	120.02	118.01
2	A	801	HEM	CBA-CAA-C2A	-3.69	105.67	112.49
3	B	804	OSD	C04-C05-C10	3.59	119.95	118.01
2	B	803	HEM	CAD-CBD-CGD	-3.59	106.66	112.67
3	B	801	OSD	N02-C02-N01	3.53	121.18	118.26
2	B	803	HEM	CBA-CAA-C2A	-3.48	106.06	112.49
2	A	801	HEM	CMA-C3A-C4A	-3.41	123.22	128.46
3	A	803	OSD	C04-C05-C10	3.40	119.85	118.01
3	A	803	OSD	C24-O29-C30	3.37	126.02	119.53
3	A	802	OSD	C05-C10-N01	-3.01	119.62	122.81
2	B	803	HEM	CMA-C3A-C2A	2.94	130.49	124.94
3	B	801	OSD	C06-C05-C10	2.92	121.60	118.33
3	B	804	OSD	C05-C10-N01	-2.78	119.86	122.81
3	B	801	OSD	C06-C05-C04	-2.76	118.38	123.66
3	A	803	OSD	C26-C25-C24	2.64	121.03	118.26
2	B	803	HEM	CMC-C2C-C3C	2.57	129.49	124.68
3	A	803	OSD	O29-C24-C25	2.43	120.31	116.04
3	B	801	OSD	C26-C25-C24	2.33	120.71	118.26
2	A	801	HEM	C4C-C3C-C2C	2.32	108.52	106.90
3	A	802	OSD	C04-C05-C10	2.24	119.22	118.01
2	A	801	HEM	CMC-C2C-C3C	2.13	128.67	124.68
2	B	803	HEM	CMD-C2D-C1D	-2.13	125.19	128.46
2	B	803	HEM	C4A-C3A-C2A	2.11	108.46	107.00
3	B	804	OSD	C26-C21-C08	-2.06	117.45	120.86
3	B	801	OSD	C08-C09-C10	-2.05	119.64	121.44
3	A	802	OSD	C03-C04-C05	2.03	119.77	117.78
3	A	803	OSD	C05-C10-N01	-2.02	120.67	122.81

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	803	OSD	C31-C30-O29-C24
3	A	802	OSD	C31-C30-O29-C24
3	A	802	OSD	C24-C25-C27-N28
3	A	802	OSD	C23-C24-O29-C30
3	B	804	OSD	C24-C25-C27-N28
3	A	803	OSD	C23-C24-O29-C30


There are no ring outliers.

6 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	OSD	1	0
3	B	804	OSD	2	0
3	A	803	OSD	1	0
2	B	803	HEM	3	0
2	A	801	HEM	2	0
3	B	801	OSD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand OSD A 802





Ligand OSD B 804











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	412/421 (97%)	0.27	15 (3%) 42 40	24, 37, 66, 121	0
1	B	409/421 (97%)	0.34	25 (6%) 21 20	27, 42, 81, 113	0
All	All	821/842 (97%)	0.31	40 (4%) 29 28	24, 39, 76, 121	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	354	ALA	6.3
1	B	608	ARG	5.5
1	A	609	TYR	5.0
1	A	608	ARG	5.0
1	A	604	CYS	4.9
1	A	607	SER	4.9
1	A	606	ASN	4.4
1	A	600	VAL	4.2
1	A	353	VAL	3.9
1	B	607	SER	3.8
1	B	304	ARG	3.7
1	B	327	LEU	3.5
1	A	605	ASP	3.5
1	B	604	CYS	3.4
1	A	603	TYR	3.4
1	B	326	THR	3.2
1	A	305	PHE	3.0
1	B	360	PHE	2.9
1	B	358	GLN	2.9
1	B	386	LEU	2.9
1	B	609	TYR	2.7
1	A	327	LEU	2.7
1	B	606	ASN	2.6
1	A	352	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	357	ASP	2.6
1	A	303	PRO	2.6
1	B	342	HIS	2.5
1	B	603	TYR	2.5
1	B	495	GLY	2.5
1	B	624	ARG	2.4
1	B	375	LYS	2.4
1	B	493	PRO	2.3
1	B	600	VAL	2.3
1	B	355	THR	2.3
1	B	359	LEU	2.2
1	A	334	TYR	2.1
1	B	522	PHE	2.1
1	A	341	MET	2.1
1	B	356	LYS	2.0
1	B	721	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OSD	B	801	24/24	0.87	0.16	27,39,61,64	0
3	OSD	A	803	24/24	0.90	0.15	29,46,69,71	0
3	OSD	B	804	24/24	0.91	0.16	23,35,51,54	0
3	OSD	A	802	24/24	0.93	0.15	20,40,68,74	0
2	HEM	A	801	43/43	0.96	0.13	21,30,41,49	0
2	HEM	B	803	43/43	0.97	0.10	23,31,42,48	0
4	ZN	B	802	1/1	0.99	0.09	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around OSD A 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OSD B 804:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OSD A 802:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around HEM A 801:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.