



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 10:04 pm BST

PDB ID : 1POW  
Title : THE REFINED STRUCTURES OF A STABILIZED MUTANT AND OF WILD-TYPE PYRUVATE OXIDASE FROM LACTOBACILLUS PLANTARUM  
Authors : Muller, Y.A.; Schulz, G.E.  
Deposited on : 1993-11-09  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	 91% 9%
1	B	585	 91% 9%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9322 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PYRUVATE OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	585	Total	C	N	O	S	0	0	0
			4521	2866	779	862	14			
1	B	585	Total	C	N	O	S	0	0	0
			4521	2866	779	862	14			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: C<sub>12</sub>H<sub>19</sub>N<sub>4</sub>O<sub>7</sub>P<sub>2</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
3	B	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
4	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	60	Total	O	0	0
			60	60		
5	B	60	Total	O	0	0
			60	60		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PYRUVATE OXIDASE



- Molecule 1: PYRUVATE OXIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.60 Å   155.40 Å   167.10 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	10.00 – 2.50 95.77 – 2.25	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.50) 88.2 (95.77-2.25)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.198 , (Not available) 0.177 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.2	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 40.5	EDS
L-test for twinning <sup>1</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9322	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FAD, TPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.65	0/4616	1.24	30/6284 (0.5%)
1	B	0.64	0/4616	1.25	29/6284 (0.5%)
All	All	0.64	0/9232	1.24	59/12568 (0.5%)

There are no bond length outliers.

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	515	ARG	NE-CZ-NH1	8.93	124.77	120.30
1	A	346	TRP	CD1-CG-CD2	8.57	113.15	106.30
1	A	515	ARG	NE-CZ-NH1	8.39	124.50	120.30
1	B	345	TRP	CD1-CG-CD2	8.18	112.85	106.30
1	A	356	TRP	CD1-CG-CD2	8.14	112.82	106.30
1	A	182	TRP	CD1-CG-CD2	8.14	112.81	106.30
1	B	356	TRP	CD1-CG-CD2	8.07	112.75	106.30
1	B	24	TRP	CD1-CG-CD2	7.88	112.60	106.30
1	B	182	TRP	CD1-CG-CD2	7.86	112.59	106.30
1	A	356	TRP	CE2-CD2-CG	-7.72	101.12	107.30
1	A	346	TRP	CE2-CD2-CG	-7.71	101.13	107.30
1	B	356	TRP	CE2-CD2-CG	-7.69	101.15	107.30
1	B	345	TRP	CE2-CD2-CG	-7.66	101.18	107.30
1	B	515	ARG	NE-CZ-NH2	-7.64	116.48	120.30
1	B	346	TRP	CD1-CG-CD2	7.59	112.38	106.30
1	A	345	TRP	CD1-CG-CD2	7.52	112.32	106.30
1	B	346	TRP	CE2-CD2-CG	-7.47	101.32	107.30
1	B	182	TRP	CE2-CD2-CG	-7.34	101.43	107.30
1	A	345	TRP	CE2-CD2-CG	-7.34	101.43	107.30
1	B	174	TRP	CD1-CG-CD2	7.28	112.13	106.30
1	A	174	TRP	CD1-CG-CD2	7.28	112.13	106.30
1	B	439	ARG	NE-CZ-NH1	7.25	123.93	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	447	ASP	CB-CG-OD1	7.16	124.74	118.30
1	A	24	TRP	CD1-CG-CD2	7.11	111.98	106.30
1	A	174	TRP	CE2-CD2-CG	-7.03	101.68	107.30
1	B	24	TRP	CE2-CD2-CG	-6.99	101.70	107.30
1	B	174	TRP	CE2-CD2-CG	-6.99	101.71	107.30
1	A	447	ASP	CB-CG-OD1	6.95	124.56	118.30
1	A	182	TRP	CE2-CD2-CG	-6.94	101.75	107.30
1	A	515	ARG	NE-CZ-NH2	-6.89	116.85	120.30
1	A	24	TRP	CE2-CD2-CG	-6.76	101.89	107.30
1	A	287	TYR	CB-CG-CD2	-6.57	117.06	121.00
1	B	346	TRP	CG-CD2-CE3	6.42	139.68	133.90
1	B	378	ARG	NE-CZ-NH1	6.28	123.44	120.30
1	B	356	TRP	CG-CD2-CE3	6.23	139.50	133.90
1	B	346	TRP	CB-CG-CD1	-6.13	119.03	127.00
1	B	356	TRP	CB-CG-CD1	-6.09	119.08	127.00
1	A	218	TYR	CB-CG-CD2	-6.04	117.37	121.00
1	B	287	TYR	CB-CG-CD2	-5.95	117.43	121.00
1	A	346	TRP	CG-CD2-CE3	5.90	139.21	133.90
1	B	345	TRP	CG-CD2-CE3	5.82	139.14	133.90
1	B	345	TRP	CB-CG-CD1	-5.82	119.44	127.00
1	B	218	TYR	CB-CG-CD2	-5.81	117.51	121.00
1	A	345	TRP	CG-CD2-CE3	5.77	139.09	133.90
1	B	556	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	A	345	TRP	CB-CG-CD1	-5.55	119.79	127.00
1	A	346	TRP	CB-CG-CD1	-5.51	119.84	127.00
1	A	356	TRP	CG-CD2-CE3	5.45	138.80	133.90
1	A	346	TRP	CG-CD1-NE1	-5.44	104.66	110.10
1	A	182	TRP	CG-CD1-NE1	-5.41	104.69	110.10
1	A	265	VAL	N-CA-C	-5.38	96.48	111.00
1	B	439	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	356	TRP	CB-CG-CD1	-5.27	120.15	127.00
1	B	300	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	B	345	TRP	CG-CD1-NE1	-5.16	104.94	110.10
1	A	315	ARG	NE-CZ-NH1	5.15	122.87	120.30
1	A	174	TRP	CG-CD2-CE3	5.12	138.51	133.90
1	A	356	TRP	CG-CD1-NE1	-5.10	105.00	110.10
1	A	345	TRP	CG-CD1-NE1	-5.01	105.09	110.10

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4521	0	4467	19	0
1	B	4521	0	4467	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	26	0	16	0	0
3	B	26	0	16	0	0
4	A	53	0	31	0	0
4	B	53	0	31	0	0
5	A	60	0	0	0	0
5	B	60	0	0	0	0
All	All	9322	0	9028	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:205:LEU:HD23	1:B:333:ILE:HD11	1.80	0.63
1:A:436:TYR:HB3	1:A:439:ARG:HG3	1.83	0.59
1:A:39:SER:HB2	1:A:173:PRO:HB2	1.86	0.58
1:A:243:SER:HB2	1:A:247:ALA:HB3	1.86	0.57
1:A:195:PRO:HD2	1:A:307:ILE:HD11	1.90	0.54
1:B:237:LEU:HD21	1:B:337:VAL:HB	1.91	0.53
1:A:12:LEU:HD23	1:A:176:GLN:HG2	1.90	0.52
1:A:216:LEU:HD13	1:A:242:MET:SD	2.49	0.51
1:B:533:ALA:HA	1:B:536:GLU:O	2.11	0.50
1:A:464:HIS:O	1:A:466:PRO:HD3	2.12	0.50
1:B:242:MET:HE3	1:B:270:ALA:HB1	1.94	0.50
1:B:59:GLU:HB2	1:B:89:HIS:HB3	1.93	0.49
1:B:279:VAL:HG22	1:B:301:TYR:HB2	1.94	0.49
1:A:279:VAL:HG22	1:A:301:TYR:HB2	1.95	0.47
1:B:476:GLN:HG2	1:B:496:GLU:HG2	1.96	0.47
1:B:61:VAL:HG13	1:B:427:PRO:HB3	1.96	0.47
1:B:39:SER:HB2	1:B:173:PRO:HB2	1.96	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:GLU:HB2	1:A:89:HIS:HB3	1.96	0.47
1:B:243:SER:HB2	1:B:247:ALA:HB3	1.97	0.46
1:A:564:ALA:O	1:A:568:GLU:HG2	2.16	0.46
1:B:433:LYS:HE2	1:B:466:PRO:HD2	1.99	0.45
1:A:28:HIS:HA	1:A:52:HIS:O	2.17	0.44
1:B:144:ALA:HA	1:B:177:ILE:HG21	1.98	0.44
1:B:268:LYS:HB3	1:B:269:PRO:HD3	1.99	0.44
1:A:571:LYS:HD3	1:A:577:GLN:HA	2.00	0.44
1:A:533:ALA:HA	1:A:536:GLU:O	2.19	0.43
1:A:287:TYR:HA	1:A:288:PRO:HD2	1.95	0.42
1:A:172:LEU:HB2	1:A:173:PRO:HD3	2.01	0.42
1:A:201:ALA:HA	1:A:204:ARG:HH11	1.83	0.42
1:A:247:ALA:HB1	1:A:250:ILE:HD12	2.01	0.42
1:B:226:ALA:HB1	1:B:330:LEU:HD12	2.01	0.42
1:A:479:PHE:O	1:A:483:GLU:HG3	2.20	0.42
1:B:263:ASN:HB2	1:B:287:TYR:OH	2.20	0.41
1:B:394:VAL:HA	1:B:398:ASN:OD1	2.20	0.41
1:A:287:TYR:CZ	1:A:289:PHE:HB2	2.56	0.40
1:B:217:ILE:HG13	1:B:239:ILE:HG21	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	583/585 (100%)	568 (97%)	13 (2%)	2 (0%)	41	61
1	B	583/585 (100%)	565 (97%)	17 (3%)	1 (0%)	47	68
All	All	1166/1170 (100%)	1133 (97%)	30 (3%)	3 (0%)	41	61

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	263	ASN
1	A	263	ASN
1	A	517	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	470/470 (100%)	463 (98%)	7 (2%)	65	85
1	B	470/470 (100%)	462 (98%)	8 (2%)	60	82
All	All	940/940 (100%)	925 (98%)	15 (2%)	62	84

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LEU
1	A	188	SER
1	A	225	LYS
1	A	232	GLN
1	A	233	LEU
1	A	261	SER
1	A	517	ASN
1	B	89	HIS
1	B	114	THR
1	B	176	GLN
1	B	225	LYS
1	B	410	SER
1	B	438	GLU
1	B	498	ASN
1	B	517	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	148	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	476	GLN
1	B	148	HIS
1	B	176	GLN
1	B	367	GLN
1	B	517	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	TPP	B	611	2	22,27,27	1.97	5 (22%)	29,40,40	1.61	5 (17%)
4	FAD	A	612	-	51,58,58	1.41	5 (9%)	60,89,89	1.73	7 (11%)
4	FAD	B	612	-	51,58,58	1.42	9 (17%)	60,89,89	1.78	9 (15%)
3	TPP	A	611	2	22,27,27	2.16	5 (22%)	29,40,40	1.61	7 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TPP	B	611	2	-	2/16/17/17	0/2/2/2
4	FAD	A	612	-	-	3/30/50/50	0/6/6/6
4	FAD	B	612	-	-	2/30/50/50	0/6/6/6
3	TPP	A	611	2	-	4/16/17/17	0/2/2/2

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	611	TPP	C4-N3	-7.76	1.33	1.39
3	B	611	TPP	C4-N3	-6.47	1.34	1.39
4	A	612	FAD	C4X-C10	5.00	1.43	1.38
4	B	612	FAD	C4X-C10	4.88	1.43	1.38
4	A	612	FAD	C4-N3	3.97	1.39	1.33
4	B	612	FAD	C4-N3	3.38	1.38	1.33
4	A	612	FAD	C2-N1	-2.86	1.32	1.38
3	A	611	TPP	PB-O2B	-2.64	1.44	1.54
4	B	612	FAD	C2B-C1B	-2.56	1.49	1.53
3	B	611	TPP	PB-O3B	-2.50	1.45	1.54
3	B	611	TPP	PB-O2B	-2.48	1.45	1.54
4	B	612	FAD	C8A-N7A	-2.46	1.30	1.34
4	B	612	FAD	C4-C4X	2.44	1.45	1.41
4	B	612	FAD	C10-N1	2.44	1.36	1.33
4	A	612	FAD	C4-C4X	2.29	1.45	1.41
3	A	611	TPP	PB-O3B	-2.27	1.46	1.54
4	B	612	FAD	C2-N1	-2.26	1.33	1.38
3	A	611	TPP	C2-N3	-2.23	1.31	1.36
4	A	612	FAD	C8A-N7A	-2.19	1.30	1.34
3	A	611	TPP	PA-O2A	-2.18	1.45	1.55
4	B	612	FAD	P-O2P	-2.09	1.45	1.55
3	B	611	TPP	C7'-C5'	2.04	1.55	1.51
3	B	611	TPP	C2-N3	-2.03	1.31	1.36
4	B	612	FAD	PA-O5B	-2.01	1.51	1.59

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	612	FAD	C4-N3-C2	7.50	121.48	115.14
4	B	612	FAD	C4-N3-C2	7.06	121.10	115.14
4	A	612	FAD	C4X-C4-N3	-5.89	115.38	123.43
4	B	612	FAD	C4X-C4-N3	-5.35	116.12	123.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	612	FAD	O4B-C1B-C2B	-4.46	100.40	106.93
4	B	612	FAD	C1'-N10-C9A	4.31	121.69	118.29
4	B	612	FAD	O4B-C1B-C2B	-4.03	101.03	106.93
4	A	612	FAD	C1'-N10-C9A	3.80	121.28	118.29
3	A	611	TPP	C6'-N1'-C2'	3.49	121.89	115.96
3	B	611	TPP	C6'-N1'-C2'	3.48	121.89	115.96
3	A	611	TPP	C5-C4-N3	3.28	114.13	107.57
3	B	611	TPP	C5-C4-N3	3.20	113.97	107.57
3	A	611	TPP	C7'-N3-C2	-3.16	119.64	125.35
3	B	611	TPP	C7'-N3-C2	-3.09	119.76	125.35
4	B	612	FAD	C6-C5X-N5	-2.99	115.76	119.05
3	B	611	TPP	C5'-C6'-N1'	-2.91	118.97	123.82
3	A	611	TPP	N1'-C2'-N3'	-2.89	120.57	125.54
4	B	612	FAD	C9A-N10-C10	-2.84	118.19	121.91
4	B	612	FAD	N3A-C2A-N1A	2.82	133.08	128.68
4	A	612	FAD	C9A-N10-C10	-2.65	118.44	121.91
4	B	612	FAD	C1'-C2'-C3'	2.58	116.99	109.79
3	A	611	TPP	C5'-C6'-N1'	-2.57	119.53	123.82
3	B	611	TPP	N1'-C2'-N3'	-2.33	121.53	125.54
3	A	611	TPP	CM2-C2'-N1'	2.24	119.60	117.14
4	B	612	FAD	N6A-C6A-N1A	2.09	122.91	118.57
3	A	611	TPP	C2'-N3'-C4'	2.03	121.24	118.08
4	A	612	FAD	N3A-C2A-N1A	2.01	131.82	128.68
4	A	612	FAD	C1'-N10-C10	2.01	120.21	118.41

There are no chirality outliers.


All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	611	TPP	C4-C5-C6-C7
3	B	611	TPP	PB-O3A-PA-O7
4	A	612	FAD	C5'-O5'-P-O1P
3	A	611	TPP	C4-C5-C6-C7
3	A	611	TPP	C5-C6-C7-O7
3	A	611	TPP	PA-O3A-PB-O2B
3	A	611	TPP	PA-O3A-PB-O3B
4	B	612	FAD	O4B-C4B-C5B-O5B
4	A	612	FAD	C5'-O5'-P-O3P
4	A	612	FAD	O4B-C4B-C5B-O5B
4	B	612	FAD	C5'-O5'-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## Ligand FAD B 612



Bond lengths



Bond angles




Torsions




Rings


## Ligand TPP A 611




Bond lengths



Bond angles



Torsions



Rings

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	585/585 (100%)	-0.83	0 100 100	2, 10, 32, 48	0
1	B	585/585 (100%)	-0.83	0 100 100	2, 8, 28, 49	0
All	All	1170/1170 (100%)	-0.83	0 100 100	2, 9, 30, 49	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	A	610	1/1	0.91	0.12	2,2,2,2	0
2	MG	B	610	1/1	0.92	0.13	2,2,2,2	0
4	FAD	B	612	53/53	0.97	0.10	2,8,14,16	0
4	FAD	A	612	53/53	0.97	0.11	3,8,13,14	0
3	TPP	B	611	26/26	0.98	0.10	2,3,6,10	0
3	TPP	A	611	26/26	0.98	0.10	2,6,10,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around FAD B 612:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around FAD A 612:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around TPP B 611:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.