



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 04:55 am BST

PDB ID : 4PP0  
Title : Structure of the PBP NocT-M117N in complex with pyronopaline  
Authors : Morera, S.; Vigouroux, A.  
Deposited on : 2014-02-26  
Resolution : 1.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5534 (1.60-1.56)
Clashscore	141614	5861 (1.60-1.56)
Ramachandran outliers	138981	5708 (1.60-1.56)
Sidechain outliers	138945	5703 (1.60-1.56)
RSRZ outliers	127900	5431 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	<div> <div>89%</div> <div>6% . .</div> </div>
1	B	265	<div> <div>8%</div> <div>90%</div> <div>6% .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	310	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4546 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nopaline-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	1	0
			1955	1251	320	373	11			
1	B	255	Total	C	N	O	S	0	0	0
			1958	1253	321	373	11			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	INITIATING METHIONINE	UNP P35120
A	117	ASN	MET	ENGINEERED MUTATION	UNP P35120
A	284	HIS	-	EXPRESSION TAG	UNP P35120
A	285	HIS	-	EXPRESSION TAG	UNP P35120
A	286	HIS	-	EXPRESSION TAG	UNP P35120
A	287	HIS	-	EXPRESSION TAG	UNP P35120
A	288	HIS	-	EXPRESSION TAG	UNP P35120
A	289	HIS	-	EXPRESSION TAG	UNP P35120
B	25	MET	-	INITIATING METHIONINE	UNP P35120
B	117	ASN	MET	ENGINEERED MUTATION	UNP P35120
B	284	HIS	-	EXPRESSION TAG	UNP P35120
B	285	HIS	-	EXPRESSION TAG	UNP P35120
B	286	HIS	-	EXPRESSION TAG	UNP P35120
B	287	HIS	-	EXPRESSION TAG	UNP P35120
B	288	HIS	-	EXPRESSION TAG	UNP P35120
B	289	HIS	-	EXPRESSION TAG	UNP P35120

- Molecule 2 is 1-[(1S)-4-carbamimidamido-1-carboxybutyl]-5-oxo-D-proline (three-letter code: OP1) (formula: C<sub>11</sub>H<sub>18</sub>N<sub>4</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			20	11	4	5		
2	B	1	Total	C	N	O	0	0
			20	11	4	5		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	343	Total 343	O 343	0	0
5	B	189	Total 189	O 189	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nopaline-binding periplasmic protein

Chain A: 



- Molecule 1: Nopaline-binding periplasmic protein

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.56 Å   114.56 Å   37.91 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	22.99 – 1.57 49.61 – 1.57	Depositor EDS
% Data completeness (in resolution range)	100.0 (22.99-1.57) 99.8 (49.61-1.57)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 1.57 Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, $R_{free}$	0.168   ,   0.182 0.165   ,   0.180	Depositor DCC
$R_{free}$ test set	3885 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.2	Xtriage
Anisotropy	0.002	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36   ,   52.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l 0.023 for h,-h-k,-l 0.015 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4546	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, OP1, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/1996	0.61	0/2696
1	B	0.44	0/1996	0.58	0/2695
All	All	0.49	0/3992	0.60	0/5391

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1955	0	1971	18	0
1	B	1958	0	1976	9	0
2	A	20	0	15	0	0
2	B	20	0	15	0	0
3	A	28	0	42	5	0
3	B	12	0	18	2	0
4	A	14	0	20	2	0
4	B	7	0	10	1	0
5	A	343	0	0	0	0
5	B	189	0	0	1	0
All	All	4546	0	4067	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:ARG:HG3	1:A:110:ARG:HH11	1.35	0.90
1:B:175:LYS:HE3	1:B:183:ILE:HD11	1.81	0.62
1:A:175:LYS:HZ3	3:A:310:EDO:H11	1.67	0.58
1:A:183:ILE:HG23	3:A:310:EDO:H22	1.86	0.58
3:B:305:EDO:H12	5:B:461:HOH:O	2.02	0.58
1:A:110:ARG:HG3	1:A:110:ARG:NH1	2.15	0.57
1:A:175:LYS:NZ	3:A:310:EDO:H11	2.20	0.56
1:A:185:THR:HG23	3:A:310:EDO:H21	1.90	0.54
1:B:160:VAL:HG13	1:B:202:ASP:HB2	1.90	0.54
1:B:41:PRO:HB3	3:B:305:EDO:H11	1.90	0.54
1:A:99:GLN:NE2	1:A:101:ALA:HB3	2.23	0.53
1:B:99:GLN:NE2	1:B:101:ALA:HB3	2.26	0.50
1:A:218:PRO:O	1:A:221:LYS:HG2	2.12	0.48
1:A:175:LYS:HD3	4:A:307:PEG:H21	1.96	0.48
1:A:99:GLN:NE2	1:A:102:ARG:HG3	2.29	0.47
1:A:110:ARG:HH11	1:A:110:ARG:CG	2.16	0.46
1:A:194:MET:HG2	1:B:47:ALA:O	2.15	0.45
1:A:273:GLN:HE22	1:B:221:LYS:HZ3	1.65	0.45
1:A:67:ASN:HB2	3:A:309:EDO:H11	1.99	0.44
1:B:139:LEU:O	4:B:302:PEG:H31	2.18	0.44
1:A:273:GLN:HE22	1:B:221:LYS:NZ	2.16	0.43
1:A:130:THR:CG2	1:A:156:ILE:HD13	2.49	0.43
1:A:110:ARG:NE	1:A:259:ASP:OD1	2.53	0.42
1:A:175:LYS:NZ	4:A:307:PEG:H32	2.34	0.42
1:B:63:CYS:HA	1:B:66:MET:HE3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/265 (96%)	248 (98%)	5 (2%)	0	100	100
1	B	253/265 (96%)	248 (98%)	5 (2%)	0	100	100
All	All	506/530 (96%)	496 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/219 (95%)	203 (97%)	6 (3%)	42	16
1	B	209/219 (95%)	203 (97%)	6 (3%)	42	16
All	All	418/438 (95%)	406 (97%)	12 (3%)	42	16

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ARG
1	A	129	LYS
1	A	158	GLU
1	A	219	ASP
1	A	221	LYS
1	A	273	GLN
1	B	88	ARG
1	B	110	ARG
1	B	160	VAL
1	B	197	LYS
1	B	200	ARG
1	B	219	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	GLN
1	A	99	GLN
1	A	273	GLN
1	B	99	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	PEG	A	306	-	6,6,6	0.16	0	5,5,5	0.08	0
4	PEG	A	307	-	6,6,6	0.07	0	5,5,5	0.07	0
3	EDO	A	303	-	3,3,3	0.47	0	2,2,2	0.48	0
3	EDO	A	308	-	3,3,3	0.61	0	2,2,2	0.14	0
3	EDO	A	302	-	3,3,3	0.53	0	2,2,2	0.27	0
4	PEG	B	302	-	6,6,6	0.12	0	5,5,5	0.06	0
3	EDO	A	310	-	3,3,3	0.45	0	2,2,2	0.18	0
3	EDO	A	305	-	3,3,3	0.55	0	2,2,2	0.28	0
3	EDO	B	303	-	3,3,3	0.43	0	2,2,2	0.37	0
2	OP1	B	301	-	14,20,20	2.94	5 (35%)	15,27,27	4.46	5 (33%)
3	EDO	B	304	-	3,3,3	0.54	0	2,2,2	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	A	304	-	3,3,3	0.58	0	2,2,2	0.29	0
2	OP1	A	301	-	14,20,20	2.46	4 (28%)	15,27,27	4.07	7 (46%)
3	EDO	B	305	-	3,3,3	0.63	0	2,2,2	0.51	0
3	EDO	A	309	-	3,3,3	0.59	0	2,2,2	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	306	-	-	2/4/4/4	-
4	PEG	A	307	-	-	3/4/4/4	-
3	EDO	A	303	-	-	1/1/1/1	-
3	EDO	A	308	-	-	1/1/1/1	-
3	EDO	A	302	-	-	0/1/1/1	-
4	PEG	B	302	-	-	0/4/4/4	-
3	EDO	A	310	-	-	1/1/1/1	-
3	EDO	A	305	-	-	1/1/1/1	-
3	EDO	B	303	-	-	1/1/1/1	-
2	OP1	B	301	-	-	1/11/32/32	0/1/1/1
3	EDO	B	304	-	-	0/1/1/1	-
3	EDO	A	304	-	-	1/1/1/1	-
2	OP1	A	301	-	-	2/11/32/32	0/1/1/1
3	EDO	B	305	-	-	1/1/1/1	-
3	EDO	A	309	-	-	0/1/1/1	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	OP1	OAM-CAQ	5.99	1.36	1.23
2	B	301	OP1	CAQ-N	5.93	1.45	1.35
2	B	301	OP1	CZ-NE	5.81	1.44	1.33
2	A	301	OP1	CZ-NE	5.56	1.44	1.33
2	A	301	OP1	OAM-CAQ	5.18	1.34	1.23
2	A	301	OP1	CAQ-N	3.99	1.42	1.35
2	B	301	OP1	CAM-CAU	-2.67	1.47	1.54
2	A	301	OP1	CAM-CAU	-2.08	1.48	1.54
2	B	301	OP1	CAM-CAK	-2.06	1.48	1.53

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	OP1	OAM-CAQ-N	-10.34	113.27	125.22
2	B	301	OP1	CAU-N-CAQ	-9.36	104.77	113.36
2	A	301	OP1	CAU-N-CAQ	-9.17	104.94	113.36
2	A	301	OP1	CAM-CAU-N	7.36	106.87	101.97
2	A	301	OP1	CAK-CAQ-N	6.75	114.59	108.00
2	B	301	OP1	OAM-CAQ-CAK	-5.81	112.23	127.08
2	B	301	OP1	CAM-CAU-N	5.80	105.83	101.97
2	B	301	OP1	CA-N-CAU	5.45	137.91	123.44
2	A	301	OP1	CA-N-CAU	5.31	137.52	123.44
2	A	301	OP1	OAM-CAQ-N	-3.87	120.75	125.22
2	A	301	OP1	CAM-CAK-CAQ	-2.95	101.02	104.90
2	A	301	OP1	CAK-CAM-CAU	2.28	108.69	105.02

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	301	OP1	CB-CA-N-CAU
2	A	301	OP1	CB-CA-N-CAU
3	B	303	EDO	O1-C1-C2-O2
3	A	304	EDO	O1-C1-C2-O2
3	A	305	EDO	O1-C1-C2-O2
4	A	307	PEG	C1-C2-O2-C3
4	A	306	PEG	C1-C2-O2-C3
4	A	307	PEG	C4-C3-O2-C2
3	A	310	EDO	O1-C1-C2-O2
4	A	306	PEG	C4-C3-O2-C2
4	A	307	PEG	O1-C1-C2-O2
3	A	308	EDO	O1-C1-C2-O2
2	A	301	OP1	CB-CA-N-CAQ
3	B	305	EDO	O1-C1-C2-O2
3	A	303	EDO	O1-C1-C2-O2


There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	307	PEG	2	0
4	B	302	PEG	1	0
3	A	310	EDO	4	0
3	B	305	EDO	2	0
3	A	309	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	254/265 (95%)	-0.12	1 (0%) 92 93	15, 21, 38, 60	0
1	B	255/265 (96%)	0.25	20 (7%) 13 13	24, 35, 59, 80	0
All	All	509/530 (96%)	0.06	21 (4%) 37 37	15, 29, 50, 80	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	159	GLY	6.1
1	B	68	ILE	5.2
1	B	67	ASN	4.3
1	B	28	TYR	4.0
1	B	31	ILE	3.4
1	B	30	SER	3.4
1	B	29	LYS	3.3
1	A	28	TYR	2.9
1	B	70	CYS	2.9
1	B	89	TYR	2.8
1	B	253	LEU	2.8
1	B	129	LYS	2.5
1	B	282	LYS	2.4
1	B	246	GLU	2.4
1	B	155	LYS	2.2
1	B	71	LYS	2.2
1	B	87	GLY	2.2
1	B	69	GLU	2.1
1	B	78	ASP	2.1
1	B	105	VAL	2.1
1	B	88	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PEG	B	302	7/7	0.50	0.35	53,60,64,64	0
4	PEG	A	307	7/7	0.52	0.24	50,57,62,64	0
3	EDO	A	309	4/4	0.73	0.19	72,72,72,72	0
3	EDO	A	304	4/4	0.76	0.13	58,59,60,60	0
3	EDO	A	302	4/4	0.78	0.21	36,37,39,42	0
3	EDO	B	304	4/4	0.81	0.11	57,57,58,58	0
3	EDO	B	303	4/4	0.83	0.17	45,46,48,48	0
4	PEG	A	306	7/7	0.83	0.12	43,55,63,64	0
3	EDO	A	308	4/4	0.84	0.12	43,45,48,50	0
3	EDO	B	305	4/4	0.87	0.17	30,35,36,42	0
3	EDO	A	310	4/4	0.88	0.27	40,42,42,44	0
3	EDO	A	305	4/4	0.90	0.16	34,43,47,49	0
3	EDO	A	303	4/4	0.90	0.24	44,47,47,47	0
2	OP1	B	301	20/20	0.91	0.09	24,26,29,30	0
2	OP1	A	301	20/20	0.97	0.08	15,16,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around OP1 B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around OP1 A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.