



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 10:09 pm BST

PDB ID : 6PTV  
Title : Crystal structure of a DnaN sliding clamp (DNA polymerase III subunit beta) from *Rickettsia rickettsii* bound to griselimycin  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2019-07-16  
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 89%, yellow 9%, grey 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>89%</span> <span>9%</span> <span>.</span> </div> </div>
1	B	387	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 89%, yellow 9%, grey 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>89%</span> <span>8%</span> <span>.</span> </div> </div>
1	C	387	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 88%, yellow 9%, grey 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>88%</span> <span>9%</span> <span>.</span> </div> </div>
1	D	387	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 90%, yellow 8%, grey 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>90%</span> <span>8%</span> <span>.</span> </div> </div>
2	X	11	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 55%, yellow 27%, orange 18%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>55%</span> <span>27%</span> <span>18%</span> </div> </div>
2	Y	11	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 45%, yellow 55%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>45%</span> <span>55%</span> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13188 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta sliding clamp.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	380	Total	C	N	O	S	0	10	0
			2959	1888	475	585	11			
1	B	376	Total	C	N	O	S	0	7	0
			2917	1864	469	573	11			
1	C	374	Total	C	N	O	S	0	9	0
			2895	1854	460	570	11			
1	D	381	Total	C	N	O	S	0	7	0
			2942	1878	472	581	11			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP A0A0H3AWV3
A	-6	ALA	-	expression tag	UNP A0A0H3AWV3
A	-5	HIS	-	expression tag	UNP A0A0H3AWV3
A	-4	HIS	-	expression tag	UNP A0A0H3AWV3
A	-3	HIS	-	expression tag	UNP A0A0H3AWV3
A	-2	HIS	-	expression tag	UNP A0A0H3AWV3
A	-1	HIS	-	expression tag	UNP A0A0H3AWV3
A	0	HIS	-	expression tag	UNP A0A0H3AWV3
A	373	ILE	VAL	conflict	UNP A0A0H3AWV3
A	375	LEU	MET	conflict	UNP A0A0H3AWV3
B	-7	MET	-	initiating methionine	UNP A0A0H3AWV3
B	-6	ALA	-	expression tag	UNP A0A0H3AWV3
B	-5	HIS	-	expression tag	UNP A0A0H3AWV3
B	-4	HIS	-	expression tag	UNP A0A0H3AWV3
B	-3	HIS	-	expression tag	UNP A0A0H3AWV3
B	-2	HIS	-	expression tag	UNP A0A0H3AWV3
B	-1	HIS	-	expression tag	UNP A0A0H3AWV3
B	0	HIS	-	expression tag	UNP A0A0H3AWV3
B	373	ILE	VAL	conflict	UNP A0A0H3AWV3
B	375	LEU	MET	conflict	UNP A0A0H3AWV3
C	-7	MET	-	initiating methionine	UNP A0A0H3AWV3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-6	ALA	-	expression tag	UNP A0A0H3AWV3
C	-5	HIS	-	expression tag	UNP A0A0H3AWV3
C	-4	HIS	-	expression tag	UNP A0A0H3AWV3
C	-3	HIS	-	expression tag	UNP A0A0H3AWV3
C	-2	HIS	-	expression tag	UNP A0A0H3AWV3
C	-1	HIS	-	expression tag	UNP A0A0H3AWV3
C	0	HIS	-	expression tag	UNP A0A0H3AWV3
C	373	ILE	VAL	conflict	UNP A0A0H3AWV3
C	375	LEU	MET	conflict	UNP A0A0H3AWV3
D	-7	MET	-	initiating methionine	UNP A0A0H3AWV3
D	-6	ALA	-	expression tag	UNP A0A0H3AWV3
D	-5	HIS	-	expression tag	UNP A0A0H3AWV3
D	-4	HIS	-	expression tag	UNP A0A0H3AWV3
D	-3	HIS	-	expression tag	UNP A0A0H3AWV3
D	-2	HIS	-	expression tag	UNP A0A0H3AWV3
D	-1	HIS	-	expression tag	UNP A0A0H3AWV3
D	0	HIS	-	expression tag	UNP A0A0H3AWV3
D	373	ILE	VAL	conflict	UNP A0A0H3AWV3
D	375	LEU	MET	conflict	UNP A0A0H3AWV3

- Molecule 2 is a protein called ACE-MVA-MP8-NZC-LEU-MP8-LEU-MVA-PRO-MLU-GLY.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	X	11	Total	C	N	O	0	0	0
			79	57	10	12			
2	Y	11	Total	C	N	O	0	0	0
			79	57	10	12			

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		
3	D	1	Total	C	O	0	0
			8	6	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	364	Total	O	0	0
			364	364		
4	B	312	Total	O	0	0
			312	312		
4	X	1	Total	O	0	0
			1	1		
4	C	273	Total	O	0	0
			273	273		
4	Y	6	Total	O	0	0
			6	6		
4	D	345	Total	O	0	0
			345	345		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Beta sliding clamp




#### • Molecule 1: Beta sliding clamp



#### • Molecule 1: Beta sliding clamp



#### • Molecule 1: Beta sliding clamp





- Molecule 2: ACE-MVA-MP8-NZC-LEU-MP8-LEU-MVA-PRO-MLU-GLY



- Molecule 2: ACE-MVA-MP8-NZC-LEU-MP8-LEU-MVA-PRO-MLU-GLY



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.21Å 82.79Å 82.86Å 109.38° 89.94° 106.53°	Depositor
Resolution (Å)	45.90 – 1.85 46.36 – 1.85	Depositor EDS
% Data completeness (in resolution range)	97.4 (45.90-1.85) 97.4 (46.36-1.85)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.21 (at 1.86Å)	Xtriage
Refinement program	PHENIX (1.16_3546: ???)	Depositor
R, $R_{free}$	0.169 , 0.206 0.169 , 0.206	Depositor DCC
$R_{free}$ test set	1969 reflections (1.28%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.9	Xtriage
Anisotropy	0.451	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 51.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	13188	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, ACE, NZC, MVA, MLU, MP8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/3031	0.56	0/4111
1	B	0.36	0/2982	0.54	0/4041
1	C	0.34	0/2965	0.55	0/4021
1	D	0.35	0/3007	0.54	0/4077
2	X	0.31	0/24	0.82	0/26
2	Y	0.28	0/24	0.92	0/26
All	All	0.35	0/12033	0.55	0/16302

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	X	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	X	8	MVA	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2959	0	2965	21	0
1	B	2917	0	2941	21	0
1	C	2895	0	2923	19	0
1	D	2942	0	2956	18	0
2	X	79	0	96	3	0
2	Y	79	0	96	3	0
3	A	8	0	14	0	0
3	D	8	0	14	1	0
4	A	364	0	0	1	0
4	B	312	0	0	3	0
4	C	273	0	0	1	0
4	D	345	0	0	4	0
4	X	1	0	0	0	0
4	Y	6	0	0	0	0
All	All	13188	0	12005	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:134:ASP:HB3	1:C:182[B]:VAL:HG11	1.73	0.70
1:A:253:PRO:HD2	1:A:373:ILE:HD12	1.74	0.68
1:A:359:PRO:HB3	1:A:375:LEU:HD23	1.76	0.66
1:B:80[A]:ARG:NH2	4:B:403:HOH:O	2.29	0.65
1:A:42[B]:ASN:OD1	4:A:501:HOH:O	2.13	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/387 (100%)	376 (97%)	12 (3%)	0	100	100
1	B	379/387 (98%)	367 (97%)	12 (3%)	0	100	100
1	C	379/387 (98%)	367 (97%)	12 (3%)	0	100	100
1	D	386/387 (100%)	373 (97%)	13 (3%)	0	100	100
2	X	3/11 (27%)	2 (67%)	0	1 (33%)	0	0
2	Y	3/11 (27%)	3 (100%)	0	0	100	100
All	All	1538/1570 (98%)	1488 (97%)	49 (3%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	X	9	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/355 (95%)	333 (98%)	5 (2%)	65	53
1	B	335/355 (94%)	333 (99%)	2 (1%)	86	83
1	C	333/355 (94%)	329 (99%)	4 (1%)	71	62
1	D	336/355 (95%)	334 (99%)	2 (1%)	86	83
2	X	3/3 (100%)	3 (100%)	0	100	100
2	Y	3/3 (100%)	3 (100%)	0	100	100
All	All	1348/1426 (94%)	1335 (99%)	13 (1%)	78	69

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	137	LYS
1	B	176	ARG
1	C	250	THR
1	A	365	PRO

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	176	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MP8	X	6	2	5,8,9	0.72	0	3,10,12	0.84	0
2	MLU	Y	10	2	7,8,9	0.24	0	6,9,11	0.83	0
2	MP8	Y	3	2	5,8,9	0.81	0	3,10,12	0.93	0
2	MP8	X	3	2	5,8,9	0.82	0	3,10,12	0.97	0
2	NZC	X	4	2	6,7,8	0.40	0	6,8,10	0.62	0
2	MVA	Y	8	2	6,7,8	0.30	0	7,8,10	0.87	0
2	NZC	Y	4	2	6,7,8	0.35	0	6,8,10	0.60	0
2	MVA	X	8	2	6,7,8	0.25	0	7,8,10	0.67	0
2	MLU	X	10	2	7,8,9	0.28	0	6,9,11	0.78	0
2	MVA	X	2	2	6,7,8	0.21	0	7,8,10	0.73	0
2	MP8	Y	6	2	5,8,9	0.74	0	3,10,12	0.72	0
2	MVA	Y	2	2	6,7,8	0.15	0	7,8,10	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MP8	X	6	2	-	0/0/11/13	0/1/1/1
2	MLU	Y	10	2	-	1/5/8/10	-
2	MP8	Y	3	2	-	0/0/11/13	0/1/1/1
2	MP8	X	3	2	-	0/0/11/13	0/1/1/1
2	NZC	X	4	2	-	1/5/8/10	-
2	MVA	Y	8	2	-	2/6/8/10	-
2	NZC	Y	4	2	-	1/5/8/10	-
2	MVA	X	8	2	-	1/6/8/10	-
2	MLU	X	10	2	-	0/5/8/10	-
2	MVA	X	2	2	-	1/6/8/10	-
2	MP8	Y	6	2	-	0/0/11/13	0/1/1/1
2	MVA	Y	2	2	-	1/6/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Y	10	MLU	O-C-CA-CB
2	X	2	MVA	CB-CA-N-CN
2	X	4	NZC	CB-CA-N-C40
2	Y	4	NZC	CB-CA-N-C40
2	Y	8	MVA	CB-CA-N-CN

There are no ring outliers.

7 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Y	10	MLU	1	0
2	Y	3	MP8	1	0
2	X	3	MP8	1	0
2	X	4	NZC	1	0
2	Y	8	MVA	1	0
2	Y	4	NZC	1	0
2	X	8	MVA	2	0

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	MPD	D	401	-	7,7,7	0.24	0	9,10,10	0.23	0
3	MPD	A	401	-	7,7,7	0.23	0	9,10,10	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	D	401	-	-	4/5/5/5	-
3	MPD	A	401	-	-	1/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	401	MPD	O2-C2-C3-C4
3	D	401	MPD	C1-C2-C3-C4
3	D	401	MPD	CM-C2-C3-C4
3	A	401	MPD	C2-C3-C4-C5
3	D	401	MPD	C2-C3-C4-O4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	401	MPD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	380/387 (98%)	-0.45	4 (1%) 80 81	20, 31, 60, 85	0
1	B	376/387 (97%)	-0.49	1 (0%) 94 93	19, 36, 64, 92	0
1	C	374/387 (96%)	-0.49	1 (0%) 94 93	18, 38, 65, 92	0
1	D	381/387 (98%)	-0.47	3 (0%) 86 86	22, 33, 64, 85	0
2	X	4/11 (36%)	0.11	0 100 100	45, 52, 58, 63	0
2	Y	4/11 (36%)	-0.08	0 100 100	40, 47, 51, 52	0
All	All	1519/1570 (96%)	-0.47	9 (0%) 89 89	18, 35, 64, 92	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	31	TYR	4.1
1	D	113	VAL	3.9
1	A	27	VAL	3.8
1	D	118	ALA	3.8
1	A	118	ALA	3.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MLU	Y	10	9/10	0.67	0.29	54,59,65,69	0
2	MLU	X	10	9/10	0.74	0.29	51,57,67,80	0
2	MVA	X	8	8/9	0.78	0.26	73,78,94,95	0
2	MVA	Y	8	8/9	0.81	0.15	48,55,58,60	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NZC	Y	4	8/9	0.92	0.09	30,36,39,40	0
2	MP8	X	3	8/9	0.93	0.10	25,27,32,39	0
2	NZC	X	4	8/9	0.93	0.10	29,32,37,45	0
2	MP8	Y	3	8/9	0.93	0.09	23,28,34,41	0
2	MP8	X	6	8/9	0.95	0.12	43,47,53,65	0
2	MVA	Y	2	8/9	0.97	0.08	25,27,30,33	0
2	MP8	Y	6	8/9	0.98	0.09	41,44,50,59	0
2	MVA	X	2	8/9	0.98	0.07	24,29,32,33	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MPD	A	401	8/8	0.88	0.18	40,55,62,65	0
3	MPD	D	401	8/8	0.90	0.13	33,48,54,56	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.