



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 05:47 am BST

PDB ID : 1PUP  
Title : CRYSTAL STRUCTURE OF A PEPTIDE NUCLEIC ACID (PNA) DU-  
PLEX AT 1.7 ANGSTROMS RESOLUTION  
Authors : Rasmussen, H.; Kastrup, J.S.  
Deposited on : 1996-11-01  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4695 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	7	 86% 14%
1	B	7	 57% 43%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called PNA (H-P(\*CPN\*GPN\*TPN\*APN\*CPN\*GPN)-NH<sub>2</sub>).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	7	Total	C	N	O	0	2	1
			128	70	38	20			
1	B	7	Total	C	N	O	0	2	1
			128	70	38	20			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	33	Total	O	0	0
			33	33		
2	B	49	Total	O	0	0
			49	49		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: PNA (H-P(\*CPN\*GPN\*TPN\*APN\*CPN\*GPN)-NH<sub>2</sub>)

Chain A: 



- Molecule 1: PNA (H-P(\*CPN\*GPN\*TPN\*APN\*CPN\*GPN)-NH<sub>2</sub>)

Chain B: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P -1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	17.97Å 26.92Å 33.78Å 88.20° 79.40° 82.50°	Depositor
Resolution (Å)	6.00 – 1.70	Depositor
% Data completeness (in resolution range)	77.0 (6.00-1.70)	Depositor
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.205 , 0.250	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	338	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPN, GPN, APN, CPN, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	128	0	82	1	0
1	B	128	0	81	2	0
2	A	33	0	0	1	0
2	B	49	0	0	1	0
All	All	338	0	163	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3[B]:TPN:O1'	2:A:65:HOH:O	1.92	0.87
1:B:10[A]:TPN:O1'	2:B:33:HOH:O	2.15	0.63
1:B:8:CPN:O7'	1:B:9:GPN:H8'1	2.17	0.45

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

16 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CPN	B	12	1	15,18,19	1.30	1 (6%)	16,23,25	1.72	4 (25%)
1	TPN	B	10[A]	1	16,19,20	1.31	3 (18%)	15,25,27	3.36	4 (26%)
1	CPN	B	8	1	15,18,19	0.81	0	16,23,25	2.05	4 (25%)
1	APN	B	11[B]	1	18,21,22	0.92	0	15,28,30	1.38	2 (13%)
1	APN	A	4[A]	1	18,21,22	0.95	1 (5%)	15,28,30	1.80	3 (20%)
1	APN	B	11[A]	1	18,21,22	0.92	0	15,28,30	1.27	0
1	TPN	A	3[A]	1	16,19,20	1.29	3 (18%)	15,25,27	3.40	3 (20%)
1	GPN	B	9	1	18,22,23	1.31	2 (11%)	17,30,32	3.03	6 (35%)
1	APN	A	4[B]	1	18,21,22	0.97	1 (5%)	15,28,30	1.94	4 (26%)
1	CPN	A	5	1	15,18,19	0.95	1 (6%)	16,23,25	1.82	4 (25%)
1	TPN	A	3[B]	1	16,19,20	1.29	3 (18%)	15,25,27	3.60	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CPN	A	1	1	15,18,19	0.86	0	16,23,25	1.97	6 (37%)
1	GPN	A	6	1	18,22,23	1.50	3 (16%)	17,30,32	2.87	5 (29%)
1	GPN	A	2	1	18,22,23	1.19	1 (5%)	17,30,32	3.01	8 (47%)
1	GPN	B	13	1	18,22,23	1.62	3 (16%)	17,30,32	2.91	7 (41%)
1	TPN	B	10[B]	1	16,19,20	1.33	3 (18%)	15,25,27	3.42	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CPN	B	12	1	-	0/13/14/15	0/1/1/1
1	TPN	B	10[A]	1	-	0/13/14/15	0/1/1/1
1	CPN	B	8	1	-	1/13/14/15	0/1/1/1
1	APN	B	11[B]	1	-	0/13/14/15	0/2/2/2
1	APN	A	4[A]	1	-	0/13/14/15	0/2/2/2
1	APN	B	11[A]	1	-	0/13/14/15	0/2/2/2
1	TPN	A	3[A]	1	-	1/13/14/15	0/1/1/1
1	GPN	B	9	1	-	0/13/14/15	0/2/2/2
1	APN	A	4[B]	1	-	0/13/14/15	0/2/2/2
1	CPN	A	5	1	-	0/13/14/15	0/1/1/1
1	TPN	A	3[B]	1	-	0/13/14/15	0/1/1/1
1	CPN	A	1	1	-	2/13/14/15	0/1/1/1
1	GPN	A	6	1	-	0/13/14/15	0/2/2/2
1	GPN	A	2	1	-	0/13/14/15	0/2/2/2
1	GPN	B	13	1	-	1/13/14/15	0/2/2/2
1	TPN	B	10[B]	1	-	1/13/14/15	0/1/1/1

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	13	GPN	C6-N1	4.25	1.40	1.33
1	A	6	GPN	C6-N1	4.01	1.40	1.33
1	B	9	GPN	C6-N1	3.39	1.38	1.33
1	B	12	CPN	C8'-N1	3.28	1.50	1.47
1	A	2	GPN	C6-N1	3.06	1.38	1.33
1	B	10[A]	TPN	C6-C5	-2.75	1.32	1.40
1	B	10[B]	TPN	C6-C5	-2.75	1.32	1.40
1	A	3[A]	TPN	C6-C5	-2.50	1.33	1.40
1	A	3[B]	TPN	C6-C5	-2.50	1.33	1.40

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	13	GPN	O1'-C'	2.48	1.33	1.19
1	B	10[A]	TPN	C4-C5	2.47	1.46	1.41
1	B	10[B]	TPN	C4-C5	2.47	1.46	1.41
1	A	3[A]	TPN	C4-N3	2.46	1.37	1.33
1	A	3[B]	TPN	C4-N3	2.46	1.37	1.33
1	A	6	GPN	C8-N7	-2.39	1.30	1.34
1	A	3[A]	TPN	C4-C5	2.37	1.46	1.41
1	A	3[B]	TPN	C4-C5	2.37	1.46	1.41
1	B	9	GPN	C6-C5	2.18	1.45	1.41
1	B	10[A]	TPN	C4-N3	2.17	1.36	1.33
1	B	10[B]	TPN	C4-N3	2.17	1.36	1.33
1	B	13	GPN	C8-N7	-2.12	1.30	1.34
1	A	6	GPN	C2-N1	2.08	1.39	1.35
1	A	5	CPN	C5-C4	2.04	1.46	1.41
1	A	4[B]	APN	C8-N7	-2.02	1.31	1.34
1	A	4[A]	APN	C8-N7	-2.02	1.31	1.34

All (68) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3[A]	TPN	C4-N3-C2	12.23	125.47	115.14
1	A	3[B]	TPN	C4-N3-C2	12.23	125.47	115.14
1	B	10[A]	TPN	C4-N3-C2	11.70	125.02	115.14
1	B	10[B]	TPN	C4-N3-C2	11.70	125.02	115.14
1	B	9	GPN	C5-C6-N1	-8.32	112.06	123.43
1	A	2	GPN	C5-C6-N1	-8.02	112.46	123.43
1	A	6	GPN	C5-C6-N1	-7.79	112.77	123.43
1	B	13	GPN	C5-C6-N1	-6.92	113.96	123.43
1	B	9	GPN	C6-N1-C2	6.10	125.62	115.93
1	A	6	GPN	C6-N1-C2	5.80	125.14	115.93
1	A	2	GPN	C6-N1-C2	5.73	125.04	115.93
1	B	8	CPN	C2-N3-C4	5.44	121.85	116.34
1	B	13	GPN	O1'-C'-C5'	-5.30	110.40	126.39
1	B	13	GPN	C6-N1-C2	4.81	123.58	115.93
1	A	1	CPN	C2-N3-C4	4.44	120.84	116.34
1	A	3[B]	TPN	C2'-C3'-N4'	-4.33	104.92	113.11
1	A	4[B]	APN	C8'-N9-C8	4.14	130.54	125.66
1	A	4[A]	APN	C8'-N9-C8	4.14	130.54	125.66
1	A	5	CPN	C2-N3-C4	4.09	120.48	116.34
1	B	9	GPN	N3-C2-N1	-3.99	121.91	127.22
1	A	4[B]	APN	C8'-C7'-N4'	3.85	121.89	117.07
1	A	4[A]	APN	C8'-C7'-N4'	3.85	121.89	117.07

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	6	GPN	N3-C2-N1	-3.81	122.14	127.22
1	B	10[A]	TPN	C8'-N1-C2	-3.80	113.53	117.86
1	B	10[B]	TPN	C8'-N1-C2	-3.80	113.53	117.86
1	B	12	CPN	C2-N3-C4	3.62	120.01	116.34
1	B	12	CPN	C8'-N1-C6	3.59	125.03	118.63
1	A	5	CPN	C8'-N1-C6	3.52	124.91	118.63
1	A	2	GPN	N3-C2-N1	-3.51	122.54	127.22
1	B	13	GPN	N3-C2-N1	-3.36	122.74	127.22
1	B	8	CPN	O1'-C'-C5'	-3.27	116.51	126.39
1	B	13	GPN	C5'-N4'-C3'	-3.25	112.44	117.06
1	A	1	CPN	O1'-C'-C5'	-3.16	116.86	126.39
1	B	10[B]	TPN	C5'-N4'-C3'	-3.11	112.64	117.06
1	A	4[B]	APN	C2'-C3'-N4'	-3.07	107.30	113.11
1	A	2	GPN	O1'-C'-C5'	-3.05	117.18	126.39
1	A	1	CPN	C8'-N1-C6	2.90	123.81	118.63
1	B	9	GPN	O1'-C'-C5'	-2.88	117.69	126.39
1	A	5	CPN	C8'-N1-C2	-2.84	114.63	117.86
1	A	3[A]	TPN	C8'-N1-C2	-2.74	114.74	117.86
1	A	3[B]	TPN	C8'-N1-C2	-2.74	114.74	117.86
1	A	3[B]	TPN	C5'-N4'-C3'	-2.73	113.18	117.06
1	A	2	GPN	C5'-N4'-C3'	-2.66	113.28	117.06
1	A	1	CPN	C5'-N4'-C3'	-2.65	113.30	117.06
1	A	2	GPN	C8'-N9-C8	2.55	128.68	125.66
1	A	2	GPN	C7'-C8'-N9	-2.54	107.74	110.76
1	B	12	CPN	C8'-N1-C2	-2.52	114.99	117.86
1	B	13	GPN	C7'-C8'-N9	-2.46	107.83	110.76
1	B	8	CPN	C2'-C3'-N4'	-2.45	108.47	113.11
1	B	9	GPN	C8'-N9-C8	2.45	128.56	125.66
1	B	13	GPN	C6-C5-C4	-2.45	118.46	120.80
1	A	6	GPN	O1'-C'-C5'	-2.38	119.21	126.39
1	B	11[B]	APN	C2'-C3'-N4'	-2.36	108.65	113.11
1	A	5	CPN	O1'-C'-C5'	-2.33	119.37	126.39
1	B	8	CPN	C8'-N1-C6	2.31	122.75	118.63
1	B	10[A]	TPN	C8'-N1-C6	2.30	124.88	119.72
1	B	10[B]	TPN	C8'-N1-C6	2.30	124.88	119.72
1	B	9	GPN	C8'-C7'-N4'	2.27	119.91	117.07
1	A	4[A]	APN	O1'-C'-C5'	-2.26	119.56	126.39
1	B	12	CPN	O1'-C'-C5'	-2.26	119.58	126.39
1	A	6	GPN	C7'-C8'-N9	-2.23	108.11	110.76
1	B	10[A]	TPN	O1'-C'-C5'	-2.22	119.69	126.39
1	A	1	CPN	C8'-N1-C2	-2.22	115.33	117.86
1	A	1	CPN	C2'-C3'-N4'	-2.19	108.97	113.11

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3[A]	TPN	C5'-N4'-C3'	-2.19	113.95	117.06
1	A	2	GPN	C2-N3-C4	-2.04	113.02	115.36
1	A	4[B]	APN	O1'-C'-C5'	-2.03	120.25	126.39
1	B	11[B]	APN	O1'-C'-C5'	-2.02	120.29	126.39

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	10[B]	TPN	O7'-C7'-N4'-C5'
1	A	1	CPN	C8'-C7'-N4'-C3'
1	A	3[A]	TPN	C'-C5'-N4'-C7'
1	B	13	GPN	C'-C5'-N4'-C7'
1	B	8	CPN	C'-C5'-N4'-C3'
1	A	1	CPN	C'-C5'-N4'-C3'

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	10[A]	TPN	1	0
1	B	8	CPN	1	0
1	B	9	GPN	1	0
1	A	3[B]	TPN	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.