



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2020 – 10:21 PM BST

PDB ID : 6PUK
Title : Structure of human MAIT A-F7 TCR in complex with human MR1-JYM72
Authors : Awad, W.; Keller, A.N.; Rossjohn, J.
Deposited on : 2019-07-18
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>••</div> </div> </div>
1	C	271	<div> <div>89%</div> <div>9%</div> <div>•</div> </div>
2	B	204	<div> <div>94%</div> <div>5%</div> <div>•</div> </div>
2	D	204	<div> <div>11%</div> <div>88%</div> <div>8%</div> <div>•</div> </div>
3	E	246	<div> <div>4%</div> <div>89%</div> <div>10%</div> <div>•</div> </div>
3	G	246	<div> <div>92%</div> <div>7%</div> <div>••</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	100	<div><div></div><div>95%</div><div></div></div>
4	H	100	<div>%<div><div></div><div>94%</div><div></div></div></div>

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 14369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	8	0
			2216	1421	380	402	13			
1	C	266	Total	C	N	O	S	0	8	0
			2221	1423	384	403	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q95460
A	261	SER	CYS	conflict	UNP Q95460
C	0	MET	-	initiating methionine	UNP Q95460
C	261	SER	CYS	conflict	UNP Q95460

- Molecule 2 is a protein called TRA@ protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	201	Total	C	N	O	S	0	10	0
			1626	1029	260	327	10			
2	D	196	Total	C	N	O	S	0	3	0
			1514	960	240	304	10			

- Molecule 3 is a protein called Human TCR beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	243	Total	C	N	O	S	0	8	0
			1914	1208	324	370	12			
3	G	244	Total	C	N	O	S	0	10	0
			1964	1237	344	370	13			


- Molecule 4 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	99	Total	C	N	O	S	0	2	0
			822	527	138	154	3			
4	H	98	Total	C	N	O	S	0	0	0
			782	501	133	145	3			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
F	0	MET	-	initiating methionine	UNP P61769
H	0	MET	-	initiating methionine	UNP P61769

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 1,2-dideoxy-1-{2,6-dioxo-5-[(1E)-3-oxobut-1-en-1-yl]-1,2,3,6-tetrahydropyrimidin-4-yl}-D-ribo-hexitol (three-letter code: OYV) (formula: $C_{14}H_{20}N_2O_7$) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			22	14	2	6		
6	C	1	Total	C	N	O	0	0
			22	14	2	6		

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	H	1	Total Na 1 1	0	0
8	G	1	Total Na 1 1	0	0
8	F	1	Total Na 1 1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	209	Total O 209 209	0	0
9	B	183	Total O 183 183	0	0
9	C	237	Total O 237 237	0	0
9	D	109	Total O 109 109	0	0
9	E	106	Total O 106 106	0	0
9	F	99	Total O 99 99	0	0
9	G	242	Total O 242 242	0	0
9	H	56	Total O 56 56	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 2: TRA@ protein




- Molecule 2: TRA@ protein



- Molecule 3: Human TCR beta chain





- Molecule 3: Human TCR beta chain

Chain G: 92% 7% ..




- Molecule 4: Beta-2-microglobulin

Chain F: 95% ..



- Molecule 4: Beta-2-microglobulin

Chain H: 94% ..



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	217.06Å 69.69Å 143.32Å 90.00° 104.35° 90.00°	Depositor
Resolution (Å)	52.57 – 2.08 66.15 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.8 (52.57-2.08) 99.9 (66.15-2.08)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.08Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.175 , 0.202 0.176 , 0.202	Depositor DCC
R_{free} test set	6344 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	43.1	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	14369	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, OYV, NA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.24	0/2307	0.43	0/3136
1	C	0.24	0/2312	0.44	0/3142
2	B	0.25	0/1692	0.46	0/2292
2	D	0.25	0/1553	0.45	0/2109
3	E	0.25	0/1989	0.45	0/2713
3	G	0.27	0/2037	0.46	0/2768
4	F	0.25	0/848	0.47	0/1151
4	H	0.24	0/805	0.43	0/1098
All	All	0.25	0/13543	0.45	0/18409

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2216	0	2105	14	0
1	C	2221	0	2110	12	0
2	B	1626	0	1563	7	0
2	D	1514	0	1403	9	0
3	E	1914	0	1786	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	1964	0	1855	9	0
4	F	822	0	776	1	0
4	H	782	0	717	2	0
5	A	6	0	8	1	0
5	C	6	0	8	3	0
5	F	6	0	8	2	0
6	A	22	0	0	0	0
6	C	22	0	0	0	0
7	F	4	0	3	0	0
8	F	1	0	0	0	0
8	G	1	0	0	0	0
8	H	1	0	0	0	0
9	A	209	0	0	1	0
9	B	183	0	0	1	0
9	C	237	0	0	1	0
9	D	109	0	0	1	0
9	E	106	0	0	2	0
9	F	99	0	0	0	0
9	G	242	0	0	1	0
9	H	56	0	0	0	0
All	All	14369	0	12342	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:157:CYS:SG	3:E:172[B]:CYS:HB2	2.36	0.66
3:E:155:HIS:HB3	3:E:216:TYR:HB2	1.79	0.65
1:C:226[A]:ASP:OD1	1:C:243:SER:OG	2.14	0.65
3:G:1:ASN:N	9:G:402:HOH:O	2.28	0.64
3:G:174:ASP:OD1	3:G:194:ARG:NH2	2.31	0.61
1:A:77:LEU:HD13	1:A:92:TYR:HB2	1.84	0.59
2:B:184[A]:ASN:ND2	9:B:301:HOH:O	2.35	0.58
1:A:221:ILE:HB	1:A:224:GLU:HG3	1.86	0.58
3:E:154[B]:ASP:OD2	9:E:301:HOH:O	2.17	0.57
1:C:28:VAL:HG23	1:C:33:ILE:HD13	1.87	0.57
1:C:151:LEU:HD22	2:D:51:LEU:HD12	1.91	0.53
1:A:113:ALA:HB2	4:H:60:TRP:CE2	2.44	0.52
2:B:150:ASP:HB2	2:B:177:LYS:HD2	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:VAL:HG23	1:A:33:ILE:HD13	1.90	0.52
3:G:128:VAL:HG23	3:G:238:ALA:HB3	1.90	0.52
3:E:13:LEU:HD11	3:E:19[B]:MET:HG3	1.92	0.51
3:E:13:LEU:HD21	3:E:19[B]:MET:HG3	1.93	0.50
1:A:6[B]:ARG:NE	9:A:407:HOH:O	2.44	0.50
3:E:95:VAL:HG12	3:E:96:TRP:CD1	2.46	0.50
1:C:3:HIS:CD2	1:C:169:LEU:HD21	2.47	0.50
2:D:28:GLY:HA3	2:D:93[A]:SER:HB3	1.94	0.49
5:C:401:GOL:HO1	5:F:102:GOL:HO2	1.56	0.49
1:A:198:LEU:HD13	1:A:269:VAL:HG21	1.94	0.49
2:B:148:ASP:OD2	2:B:177:LYS:NZ	2.46	0.48
1:C:215:MET:HG3	1:C:257:HIS:CD2	2.49	0.48
1:C:241:TRP:HE1	5:C:401:GOL:C3	2.28	0.47
1:C:225:ILE:HG12	1:C:244:ILE:HG22	1.95	0.47
1:C:9:ARG:NH1	9:C:510:HOH:O	2.48	0.47
2:B:28:GLY:HA3	2:B:93[A]:SER:HB3	1.97	0.47
1:C:169:LEU:HD12	1:C:176:LEU:HD13	1.96	0.46
2:B:122[B]:ARG:H	2:B:122[B]:ARG:HG2	1.52	0.46
3:G:99[B]:GLU:HG2	3:G:100:GLY:N	2.31	0.46
1:C:113:ALA:HB2	4:F:60:TRP:CE2	2.50	0.46
3:G:123:PRO:HD3	3:G:231:PRO:HB3	1.98	0.46
2:B:157:CYS:SG	3:G:172[B]:CYS:HB2	2.55	0.46
2:D:36:GLN:HB2	2:D:46:LEU:HD11	1.98	0.45
1:A:54:LEU:HD13	1:A:58:HIS:CD2	2.51	0.45
2:D:144:SER:N	2:D:189:SER:OG	2.46	0.45
3:G:95:VAL:HG12	3:G:96:TRP:CD1	2.51	0.45
1:C:61:ARG:HG2	2:D:94:ASN:HB3	1.99	0.45
3:E:128:VAL:HG23	3:E:238:ALA:HB3	1.97	0.45
3:G:8:PRO:HD2	3:G:21:LEU:HD22	1.98	0.44
2:D:3:ASN:ND2	2:D:5:ASP:OD2	2.40	0.44
5:C:401:GOL:O1	5:F:102:GOL:O2	2.30	0.44
1:A:246:LEU:HD12	1:A:246:LEU:HA	1.82	0.43
1:C:77:LEU:HD13	1:C:92:TYR:HB2	2.00	0.43
1:A:6[B]:ARG:NH2	1:A:29:ASP:O	2.51	0.43
3:E:174:ASP:OD1	3:E:194:ARG:NH2	2.49	0.43
1:A:4:SER:HB3	1:A:99:GLU:HG2	2.01	0.43
1:A:233:SER:HB3	4:H:12:ARG:HG3	2.01	0.43
2:D:91:LYS:HG2	2:D:95:TYR:HA	2.01	0.42
1:A:64:GLN:HE22	5:A:301:GOL:H2	1.85	0.42
3:E:97:THR:O	9:E:302:HOH:O	2.22	0.42
1:A:215:MET:HE3	1:A:218:GLY:HA2	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:86:THR:HG23	3:G:113:THR:HA	2.01	0.42
3:E:83:PRO:HA	3:E:114:VAL:HB	2.02	0.42
2:B:28:GLY:HA3	2:B:93[B]:SER:OG	2.19	0.41
3:E:119:LYS:O	3:E:228:ARG:NH2	2.53	0.41
3:E:131:PRO:HG2	3:E:142:ALA:HB1	2.02	0.40
2:D:108:LYS:NZ	9:D:301:HOH:O	2.45	0.40
1:A:3:HIS:CD2	1:A:169[B]:LEU:HD21	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/271 (100%)	266 (98%)	4 (2%)	0	100	100
1	C	270/271 (100%)	265 (98%)	5 (2%)	0	100	100
2	B	209/204 (102%)	206 (99%)	3 (1%)	0	100	100
2	D	195/204 (96%)	193 (99%)	2 (1%)	0	100	100
3	E	249/246 (101%)	246 (99%)	3 (1%)	0	100	100
3	G	252/246 (102%)	251 (100%)	1 (0%)	0	100	100
4	F	99/100 (99%)	99 (100%)	0	0	100	100
4	H	96/100 (96%)	96 (100%)	0	0	100	100
All	All	1640/1642 (100%)	1622 (99%)	18 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/241 (98%)	230 (97%)	7 (3%)	41	43
1	C	236/241 (98%)	228 (97%)	8 (3%)	37	37
2	B	187/181 (103%)	185 (99%)	2 (1%)	73	78
2	D	164/181 (91%)	160 (98%)	4 (2%)	49	52
3	E	206/212 (97%)	201 (98%)	5 (2%)	49	52
3	G	210/212 (99%)	204 (97%)	6 (3%)	42	44
4	F	91/95 (96%)	87 (96%)	4 (4%)	28	27
4	H	83/95 (87%)	81 (98%)	2 (2%)	49	52
All	All	1414/1458 (97%)	1376 (97%)	38 (3%)	49	47

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	PHE
1	A	83	HIS
1	A	152	TYR
1	A	169[A]	LEU
1	A	169[B]	LEU
1	A	220	GLU
1	A	221	ILE
2	B	112	GLN
2	B	162	ARG
1	C	17	HIS
1	C	73	PHE
1	C	149[A]	GLU
1	C	149[B]	GLU
1	C	152	TYR
1	C	247	ASP
1	C	259	GLU
1	C	261	SER
2	D	138	ASP
2	D	150	ASP
2	D	157	CYS
2	D	163	SER
3	E	22	GLN
3	E	77	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	79	GLU
3	E	111	ARG
3	E	221	ASN
4	F	64[A]	LEU
4	F	64[B]	LEU
4	F	70	PHE
4	F	98	ASP
3	G	99[A]	GLU
3	G	99[B]	GLU
3	G	185	ASN
3	G	194	ARG
3	G	206[A]	ARG
3	G	206[B]	ARG
4	H	45	ARG
4	H	70	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	OYV	C	402	1	21,22,23	0.96	2 (9%)	18,29,31	4.70	2 (11%)
6	OYV	A	302	1	21,22,23	0.93	2 (9%)	18,29,31	4.73	2 (11%)
7	ACT	F	101	-	1,3,3	7.06	1 (100%)	0,3,3	0.00	-
5	GOL	C	401	-	5,5,5	0.86	0	5,5,5	1.01	0
5	GOL	A	301	-	5,5,5	0.91	0	5,5,5	0.98	0
5	GOL	F	102	-	5,5,5	0.92	0	5,5,5	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	C	401	-	-	4/4/4/4	-
5	GOL	A	301	-	-	2/4/4/4	-
6	OYV	A	302	1	-	3/19/19/20	0/1/1/1
6	OYV	C	402	1	-	2/19/19/20	0/1/1/1
5	GOL	F	102	-	-	0/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	F	101	ACT	CH3-C	7.06	1.57	1.48
6	C	402	OYV	C13-N1	3.21	1.38	1.33
6	A	302	OYV	C13-N1	2.94	1.38	1.33
6	C	402	OYV	C5-N	2.18	1.38	1.34
6	A	302	OYV	C5-N	2.05	1.37	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	302	OYV	C4-C13-N1	-14.61	114.22	124.40
6	C	402	OYV	C4-C13-N1	-14.52	114.28	124.40
6	A	302	OYV	C13-N1-C12	13.53	126.56	115.14
6	C	402	OYV	C13-N1-C12	13.51	126.55	115.14

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	C	402	OYV	C2-C3-C4-C5
5	C	401	GOL	O1-C1-C2-C3
5	A	301	GOL	O1-C1-C2-C3
5	C	401	GOL	O2-C2-C3-O3
6	C	402	OYV	C5-C6-C7-C8
6	A	302	OYV	C5-C6-C7-C8
5	C	401	GOL	C1-C2-C3-O3
5	C	401	GOL	O1-C1-C2-O2
5	A	301	GOL	O1-C1-C2-O2
6	A	302	OYV	O3-C10-C11-O4
6	A	302	OYV	C2-C3-C4-C5


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	401	GOL	3	0
5	A	301	GOL	1	0
5	F	102	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	266/271 (98%)	-0.05	5 (1%) 66 70	33, 48, 85, 97	5 (1%)
1	C	266/271 (98%)	-0.00	1 (0%) 92 93	34, 44, 69, 91	7 (2%)
2	B	201/204 (98%)	-0.12	0 100 100	32, 42, 65, 93	7 (3%)
2	D	196/204 (96%)	0.49	22 (11%) 5 6	37, 58, 95, 109	4 (2%)
3	E	243/246 (98%)	0.15	9 (3%) 41 46	41, 59, 91, 111	9 (3%)
3	G	244/246 (99%)	-0.17	0 100 100	34, 45, 60, 81	6 (2%)
4	F	99/100 (99%)	-0.21	0 100 100	36, 48, 66, 75	5 (5%)
4	H	98/100 (98%)	-0.02	1 (1%) 82 84	42, 67, 94, 99	1 (1%)
All	All	1613/1642 (98%)	0.02	38 (2%) 59 63	32, 48, 86, 111	44 (2%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	147	LYS	4.2
1	A	253	LEU	3.4
2	D	197	PHE	3.4
2	D	180	PHE	3.1
3	E	136	ILE	3.1
2	D	194	ASP	3.0
2	D	164	MET	3.0
2	D	138	ASP	2.9
2	D	146	SER	2.8
2	D	119	TYR	2.8
1	A	222	VAL	2.7
2	D	133	LEU	2.7
2	D	117	ALA	2.6
2	D	188	ASN	2.6
1	A	221	ILE	2.6
2	D	149	SER	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	121	LEU	2.6
2	D	1	GLY	2.5
3	E	202	TRP	2.5
3	E	167	VAL	2.5
2	D	196	PHE	2.4
2	D	198	PRO	2.4
3	E	197	VAL	2.4
1	A	220	GLU	2.4
3	E	201	PHE	2.4
3	E	137	SER	2.3
2	D	143	VAL	2.3
2	D	163	SER	2.2
2	D	124	SER	2.1
3	E	240	ALA	2.1
3	E	244	ALA	2.1
2	D	112	GLN	2.1
2	D	131	VAL	2.1
1	A	223	GLN	2.1
1	C	219	GLU	2.1
2	D	199	SER	2.1
3	E	209	PHE	2.1
4	H	79	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	C	401	6/6	0.78	0.21	45,50,51,52	6


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	NA	H	101	1/1	0.85	0.14	84,84,84,84	0
5	GOL	A	301	6/6	0.85	0.23	43,50,51,51	6
7	ACT	F	101	4/4	0.89	0.10	40,50,52,54	0
5	GOL	F	102	6/6	0.89	0.16	45,45,46,48	6
6	OYV	A	302	22/23	0.94	0.10	35,38,42,43	0
6	OYV	C	402	22/23	0.96	0.14	38,41,44,45	0
8	NA	F	103	1/1	0.96	0.08	54,54,54,54	0
8	NA	G	301	1/1	0.99	0.12	51,51,51,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.