



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 12:42 am BST

PDB ID : 1PVL
Title : STRUCTURE OF THE PANTON-VALENTINE LEUCOCIDIN F COMPONENT FROM STAPHYLOCOCCUS AUREUS
Authors : Pedelacq, J.D.; Mourey, L.; Maveyraud, L.; Prevost, G.; Samama, J.P.
Deposited on : 1999-01-12
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	301	<div>82% 14% . .</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MES	A	401	-	X	X	-
2	MES	A	402	-	X	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2593 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called LEUCOCIDIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	2401	1506	413	477	5	0	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	247	ARG	SER	CONFLICT	UNP O50604
A	257	ASN	ILE	CONFLICT	UNP O50604
A	258	GLN	ASN	CONFLICT	UNP O50604
A	259	LEU	PHE	CONFLICT	UNP O50604
A	260	HIS	ASN	CONFLICT	UNP O50604
A	269	GLU	HIS	CONFLICT	UNP O50604
A	270	ASN	ILE	CONFLICT	UNP O50604

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	168	Total	O	0	0
			168	168		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LEUCOCIDIN

Chain A:  82% 14% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.74Å 72.38Å 98.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.40 – 2.00 23.44 – 2.01	Depositor EDS
% Data completeness (in resolution range)	85.0 (24.40-2.00) 84.6 (23.44-2.01)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.75 (at 2.01Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.201 , 0.243 0.258 , 0.291	Depositor DCC
R_{free} test set	1080 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	15.3	Xtriage
Anisotropy	0.391	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 53.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	2593	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2456	1.21	10/3320 (0.3%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	251	ARG	NE-CZ-NH1	-9.68	115.46	120.30
1	A	130	GLY	C-N-CA	8.94	141.07	122.30
1	A	14	ASP	CB-CG-OD1	8.02	125.52	118.30
1	A	251	ARG	NE-CZ-NH2	7.34	123.97	120.30
1	A	129	SER	N-CA-CB	5.82	119.23	110.50
1	A	271	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	A	77	TYR	CB-CG-CD1	5.78	124.47	121.00
1	A	131	GLY	N-CA-C	-5.73	98.77	113.10
1	A	77	TYR	CB-CG-CD2	-5.35	117.79	121.00
1	A	291	ASP	N-CA-CB	5.23	120.02	110.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2401	0	2278	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	24	0	20	6	0
3	A	168	0	0	1	0
All	All	2593	0	2298	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:ASN:ND2	1:A:107:GLU:H	1.84	0.75
2:A:401:MES:O1S	2:A:401:MES:H52	1.89	0.73
2:A:401:MES:H32	3:A:541:HOH:O	1.93	0.69
2:A:401:MES:C5	2:A:401:MES:O1S	2.42	0.67
1:A:11:LYS:HD2	1:A:11:LYS:N	2.11	0.66
1:A:75:GLN:NE2	1:A:251:ARG:HD3	2.16	0.61
1:A:270:ASN:ND2	1:A:298:ASN:H	2.00	0.59
1:A:274:HIS:HD2	1:A:294:SER:OG	1.88	0.57
1:A:105:ASN:HD22	1:A:107:GLU:H	1.54	0.56
1:A:15:LYS:HG3	1:A:118:TYR:HE2	1.72	0.55
1:A:269:GLU:HA	1:A:269:GLU:OE1	2.07	0.55
1:A:159:PHE:H	1:A:159:PHE:HD1	1.55	0.54
1:A:156:ARG:HG3	1:A:156:ARG:HH11	1.72	0.54
1:A:161:LYS:HG2	1:A:162:ILE:N	2.22	0.54
1:A:258:GLN:HE21	1:A:258:GLN:HA	1.73	0.53
1:A:97:VAL:HG23	1:A:228:VAL:CG2	2.39	0.52
1:A:257:ASN:O	1:A:258:GLN:HB2	2.12	0.49
1:A:105:ASN:HD21	1:A:107:GLU:HB2	1.78	0.49
1:A:5:THR:O	1:A:21:THR:HG23	2.13	0.48
1:A:173:ASN:HD22	1:A:173:ASN:C	2.17	0.47
1:A:100:ALA:HA	1:A:101:PRO:C	2.35	0.47
1:A:97:VAL:HG23	1:A:228:VAL:HG23	1.98	0.46
1:A:176:TRP:CD1	2:A:401:MES:H62	2.50	0.46
1:A:105:ASN:HD22	1:A:105:ASN:C	2.19	0.45
1:A:15:LYS:HG3	1:A:118:TYR:CE2	2.50	0.45
1:A:53:LEU:HD13	1:A:53:LEU:C	2.37	0.45
2:A:401:MES:O1S	2:A:401:MES:H51	2.17	0.44
1:A:173:ASN:ND2	1:A:208:LEU:HD13	2.33	0.43
1:A:129:SER:O	1:A:130:GLY:C	2.56	0.43
1:A:146:GLN:NE2	1:A:151:THR:OG1	2.52	0.42
1:A:154:ASP:OD1	1:A:156:ARG:NH1	2.52	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:GLN:OE1	1:A:289:LEU:HD23	2.20	0.42
2:A:401:MES:S	2:A:401:MES:C5	3.07	0.41
1:A:90:SER:O	1:A:91:ASN:HB2	2.20	0.41
1:A:256:TRP:HE1	1:A:258:GLN:HE21	1.68	0.40
1:A:196:SER:HB3	1:A:199:SER:HB2	2.02	0.40
1:A:64:THR:O	1:A:76:PHE:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/301 (98%)	280 (95%)	12 (4%)	2 (1%)	22	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	130	GLY
1	A	91	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/270 (98%)	255 (97%)	9 (3%)	37	36

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	LEU
1	A	65	LYS
1	A	105	ASN
1	A	159	PHE
1	A	173	ASN
1	A	194	LEU
1	A	228	VAL
1	A	234	ASN
1	A	258	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	75	GLN
1	A	83	ASN
1	A	105	ASN
1	A	111	GLN
1	A	112	GLN
1	A	123	ASN
1	A	146	GLN
1	A	169	HIS
1	A	173	ASN
1	A	200	ASN
1	A	233	GLN
1	A	258	GLN
1	A	264	ASN
1	A	270	ASN
1	A	274	HIS
1	A	284	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MES	A	402	-	12,12,12	10.12	10 (83%)	14,16,16	2.96	7 (50%)
2	MES	A	401	-	12,12,12	10.03	10 (83%)	14,16,16	3.53	9 (64%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	A	402	-	-	2/6/14/14	0/1/1/1
2	MES	A	401	-	-	1/6/14/14	0/1/1/1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	MES	C8-S	-23.96	1.43	1.77
2	A	401	MES	C8-S	-23.62	1.43	1.77
2	A	402	MES	C7-N4	-15.44	1.11	1.47
2	A	401	MES	C7-N4	-15.23	1.12	1.47
2	A	401	MES	O1S-S	11.95	1.80	1.45
2	A	402	MES	O2S-S	11.94	1.80	1.45
2	A	402	MES	O1S-S	11.51	1.79	1.45
2	A	401	MES	O2S-S	10.90	1.77	1.45
2	A	401	MES	O3S-S	8.86	1.78	1.47
2	A	402	MES	O3S-S	8.62	1.78	1.47
2	A	402	MES	C7-C8	-5.40	1.38	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	MES	C7-C8	-5.14	1.38	1.52
2	A	401	MES	C3-C2	-3.85	1.35	1.50
2	A	401	MES	C5-N4	-3.73	1.36	1.46
2	A	402	MES	C3-C2	-3.42	1.37	1.50
2	A	401	MES	C5-C6	-3.39	1.37	1.50
2	A	401	MES	C3-N4	-3.18	1.38	1.46
2	A	402	MES	C5-C6	-2.98	1.38	1.50
2	A	402	MES	C5-N4	-2.86	1.39	1.46
2	A	402	MES	C3-N4	-2.77	1.39	1.46

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	MES	C7-N4-C3	7.54	130.51	111.23
2	A	402	MES	O3S-S-C8	6.46	116.21	105.77
2	A	401	MES	O3S-S-C8	6.39	116.10	105.77
2	A	402	MES	C7-N4-C3	6.07	126.77	111.23
2	A	401	MES	O1S-S-C8	4.70	112.57	106.92
2	A	401	MES	C2-C3-N4	4.30	116.62	110.10
2	A	402	MES	O1S-S-C8	3.65	111.31	106.92
2	A	401	MES	C6-C5-N4	3.23	115.00	110.10
2	A	402	MES	O3S-S-O2S	-2.85	104.31	111.27
2	A	402	MES	O2S-S-C8	2.56	110.00	106.92
2	A	401	MES	O3S-S-O1S	-2.53	105.08	111.27
2	A	401	MES	O2S-S-C8	2.42	109.83	106.92
2	A	401	MES	O2S-S-O1S	-2.19	106.36	113.95
2	A	402	MES	O2S-S-O1S	-2.17	106.44	113.95
2	A	402	MES	O1-C2-C3	-2.16	107.03	111.80
2	A	401	MES	O3S-S-O2S	-2.05	106.25	111.27

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	402	MES	C8-C7-N4-C3
2	A	402	MES	C8-C7-N4-C5
2	A	401	MES	C8-C7-N4-C5

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	MES	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.