



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 07:51 pm BST

PDB ID : 3PY8  
Title : Crystal structure of a mutant of the large fragment of DNA polymerase I from thermus aquaticus in a closed ternary complex with DNA and ddCTP  
Authors : Schnur, A.; Marx, A.; Welte, W.; Diederichs, K.  
Deposited on : 2010-12-12  
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	<div> <div>8%</div> <div> <div></div> <div>91%</div> <div>9%</div> </div> </div>
2	B	12	<div> <div>33%</div> <div>67%</div> </div>
3	C	16	<div> <div>6%</div> <div> <div>44%</div> <div>25%</div> <div>19%</div> <div>13%</div> </div> </div>

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5383 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	539	Total	C	N	O	S	0	1	0
			4287	2724	773	778	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	614	LYS	ILE	ENGINEERED MUTATION	UNP P19821
A	747	LYS	MET	ENGINEERED MUTATION	UNP P19821


- Molecule 2 is a DNA chain called DNA (5'-D(\*GP\*AP\*CP\*CP\*AP\*CP\*GP\*GP\*CP\*GP\*CP\*(DOC))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	P	0	0	0
			240	114	48	67	11			

- Molecule 3 is a DNA chain called DNA (5'-D(\*AP\*A\*AP\*GP\*GP\*GP\*CP\*GP\*CP\*CP\*GP\*TP\*GP\*GP\*TP\*C)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	14	Total	C	N	O	P	0	0	0
			288	136	56	83	13			

- Molecule 4 is 2',3'-DIDEOXYCYTIDINE 5'-TRIPHOSPHATE (three-letter code: DCT) (formula: C<sub>9</sub>H<sub>16</sub>N<sub>3</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	9	3	12	3		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total 1	Mn 1	0	0
6	A	3	Total 3	Mn 3	0	0
6	C	1	Total 1	Mn 1	0	0

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total 1	Mg 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	430	Total 430	O 430	0	0
8	B	37	Total 37	O 37	0	0
8	C	56	Total 56	O 56	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase I



- Molecule 2: DNA (5'-D(\*GP\*AP\*CP\*CP\*AP\*CP\*GP\*GP\*CP\*GP\*CP\*(DOC))-3')



- Molecule 3: DNA (5'-D(\*AP\*A\*AP\*GP\*GP\*GP\*CP\*GP\*CP\*CP\*GP\*TP\*GP\*GP\*TP\*C)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	108.61 Å   108.61 Å   90.57 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	41.74 – 1.74 46.57 – 1.74	Depositor EDS
% Data completeness (in resolution range)	99.9 (41.74-1.74) 99.9 (46.57-1.74)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.90 (at 1.74 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_572)	Depositor
R, $R_{free}$	0.169 , 0.199 0.172 , 0.204	Depositor DCC
$R_{free}$ test set	3183 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.6	Xtriage
Anisotropy	0.214	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5383	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, MN, DCT, DOC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/4380	0.47	0/5931
2	B	0.70	0/249	1.29	3/382 (0.8%)
3	C	0.73	0/323	1.49	6/498 (1.2%)
All	All	0.37	0/4952	0.67	9/6811 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	C	214	DG	O4'-C4'-C3'	-9.79	100.12	106.00
3	C	214	DG	C1'-O4'-C4'	-7.59	102.51	110.10
3	C	203	DA	O4'-C4'-C3'	-7.41	101.53	104.50
2	B	106	DC	O4'-C1'-N1	-6.94	103.14	108.00
3	C	214	DG	O4'-C1'-N9	6.85	112.80	108.00
3	C	212	DT	O4'-C1'-N1	-6.59	103.39	108.00
2	B	108	DG	C1'-O4'-C4'	-6.14	103.96	110.10
2	B	108	DG	O4'-C1'-N9	6.10	112.27	108.00
3	C	210	DC	O4'-C1'-C2'	5.14	110.02	105.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4287	0	4343	35	0
2	B	240	0	134	8	0
3	C	288	0	158	6	0
4	A	27	0	12	2	0
5	A	12	0	16	2	0
6	A	3	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
7	A	1	0	0	0	0
8	A	430	0	0	4	0
8	B	37	0	0	0	0
8	C	56	0	0	1	0
All	All	5383	0	4663	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:631:VAL:HG11	1:A:638:ILE:HD11	1.59	0.82
3:C:204:DG:H5''	8:C:471:HOH:O	1.93	0.68
1:A:376:LEU:HD22	1:A:420:LEU:HD12	1.78	0.65
3:C:214:DG:H2'	3:C:215:DT:H71	1.81	0.63
3:C:210:DC:H2'	3:C:211:DG:C8	2.34	0.63
2:B:111:DC:H2'	2:B:112:DOC:H6	1.83	0.61
2:B:101:DG:H2'	2:B:102:DA:C8	2.38	0.59
1:A:674:SER:HB3	3:C:203:DA:C8	2.38	0.58
1:A:642:THR:O	1:A:646:MET:HG2	2.06	0.56
1:A:299:TRP:CG	1:A:300:PRO:HA	2.41	0.55
1:A:627:ASN:O	1:A:631:VAL:HG23	2.08	0.54
1:A:471:GLU:HG2	1:A:475:PHE:CE2	2.44	0.53
1:A:694:GLU:O	1:A:698:GLN:HG3	2.09	0.53
1:A:631:VAL:HG13	1:A:636:ARG:HB2	1.90	0.53
1:A:432:GLU:OE2	5:A:834:GOL:H31	2.09	0.52
1:A:615:GLU:CD	4:A:113:DCT:H2''	2.30	0.51
2:B:104:DC:H2'	2:B:105:DA:C8	2.45	0.50
1:A:638:ILE:HG13	8:A:921:HOH:O	2.12	0.50
1:A:435:ARG:HB3	1:A:436:PRO:HD3	1.94	0.50
1:A:638:ILE:N	1:A:638:ILE:HD12	2.27	0.50
1:A:691:ALA:HA	1:A:694:GLU:HG2	1.95	0.49
2:B:111:DC:H2'	2:B:112:DOC:C6	2.41	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:625:ASP:HA	1:A:702:LYS:HB2	1.95	0.48
1:A:678:LEU:HD12	1:A:682:LEU:HD13	1.95	0.47
3:C:214:DG:H2'	3:C:215:DT:C6	2.50	0.47
2:B:104:DC:H2''	2:B:105:DA:H5'	1.96	0.46
1:A:644:SER:HB2	1:A:649:VAL:O	2.16	0.46
1:A:453:VAL:HG13	1:A:550:PRO:HB3	1.98	0.46
1:A:713:GLU:OE2	1:A:717:ARG:HG3	2.16	0.45
1:A:306:PHE:O	1:A:326:ALA:HA	2.17	0.44
4:A:113:DCT:O4'	2:B:112:DOC:H2'	2.18	0.44
1:A:691:ALA:O	1:A:694:GLU:HG2	2.17	0.43
1:A:685:PRO:HB2	1:A:688:GLU:HG3	2.00	0.43
1:A:625:ASP:O	1:A:629:ILE:HG13	2.19	0.43
1:A:647:PHE:HB2	1:A:658:MET:HE3	2.01	0.42
1:A:677:ARG:HD2	1:A:681:GLU:OE2	2.19	0.42
1:A:647:PHE:CB	1:A:658:MET:HE3	2.49	0.42
1:A:655:ASP:HB2	1:A:656:PRO:HD2	2.01	0.42
1:A:432:GLU:OE1	5:A:834:GOL:O2	2.22	0.42
1:A:633:GLN:HA	8:A:961:HOH:O	2.20	0.42
2:B:101:DG:H2''	2:B:102:DA:H5'	2.01	0.41
1:A:638:ILE:HA	8:A:921:HOH:O	2.19	0.41
1:A:668:GLY:HA2	3:C:204:DG:C4	2.55	0.41
1:A:583:ASN:HD22	1:A:583:ASN:H	1.68	0.41
1:A:462:GLU:HB3	8:A:939:HOH:O	2.21	0.41
2:B:104:DC:H2''	2:B:105:DA:C5'	2.51	0.41
1:A:770:PRO:O	1:A:774:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	538/540 (100%)	523 (97%)	14 (3%)	1 (0%)	47	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	586	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	441/441 (100%)	438 (99%)	3 (1%)	84	75

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	487	ARG
1	A	583	ASN
1	A	695	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	583	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	DOC	B	112	3,2	14,19,20	2.10	4 (28%)	13,26,29	1.39	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DOC	B	112	3,2	-	0/4/18/19	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	112	DOC	C6-N1	5.16	1.42	1.35
2	B	112	DOC	O4'-C4'	-3.16	1.38	1.44
2	B	112	DOC	C4-N4	3.02	1.44	1.35
2	B	112	DOC	O5'-C5'	-2.47	1.38	1.44

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	112	DOC	C2-N3-C4	2.75	119.13	116.34
2	B	112	DOC	C2'-C3'-C4'	-2.01	98.95	102.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	112	DOC	3	0

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	DCT	A	113	7,6	22,28,28	1.51	3 (13%)	26,43,43	1.69	7 (26%)
5	GOL	A	1	-	5,5,5	0.31	0	5,5,5	0.20	0
5	GOL	A	834	-	5,5,5	0.41	0	5,5,5	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DCT	A	113	7,6	-	3/19/31/31	0/2/2/2
5	GOL	A	1	-	-	0/4/4/4	-
5	GOL	A	834	-	-	0/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	113	DCT	C6-N1	3.67	1.40	1.35
4	A	113	DCT	O4'-C4'	-2.81	1.39	1.44
4	A	113	DCT	C4-N4	2.54	1.42	1.35

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	113	DCT	C2-N3-C4	3.16	119.55	116.34
4	A	113	DCT	C2'-C1'-N1	-2.92	106.98	112.48
4	A	113	DCT	PB-O3A-PA	-2.64	123.78	132.83
4	A	113	DCT	O2G-PG-O3B	2.53	113.11	104.64
4	A	113	DCT	O3G-PG-O3B	2.44	112.82	104.64
4	A	113	DCT	O2A-PA-O1A	-2.30	100.88	112.24
4	A	113	DCT	O4'-C4'-C5'	2.19	113.12	109.52

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	113	DCT	PB-O3A-PA-O2A
4	A	113	DCT	PG-O3B-PB-O2B
4	A	113	DCT	PB-O3A-PA-O1A

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	113	DCT	2	0
5	A	834	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	539/540 (99%)	0.17	44 (8%) <span>11</span> <span>15</span>	16, 32, 80, 107	0
2	B	11/12 (91%)	-0.32	0 <span>100</span> <span>100</span>	20, 23, 44, 45	0
3	C	14/16 (87%)	-0.11	1 (7%) <span>16</span> <span>20</span>	18, 24, 48, 134	0
All	All	564/568 (99%)	0.15	45 (7%) <span>12</span> <span>16</span>	16, 32, 79, 134	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	692	PHE	8.7
1	A	647	PHE	6.9
1	A	653	ALA	5.4
1	A	654	VAL	5.1
1	A	651	ARG	5.0
1	A	645	TRP	5.0
1	A	697	PHE	4.8
1	A	684	ILE	4.6
1	A	693	ILE	4.6
1	A	648	GLY	4.5
1	A	689	ALA	4.4
1	A	691	ALA	4.2
1	A	650	PRO	4.2
1	A	686	TYR	4.1
1	A	657	LEU	3.9
1	A	682	LEU	3.9
1	A	683	ALA	3.8
1	A	655	ASP	3.6
1	A	649	VAL	3.6
1	A	700	PHE	3.5
1	A	687	GLU	3.5
1	A	680	GLN	3.4
1	A	696	TYR	3.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	695	ARG	3.3
3	C	203	DA	3.3
1	A	699	SER	3.1
1	A	652	GLU	3.0
1	A	685	PRO	2.9
1	A	669	VAL	2.9
1	A	676	HIS	2.9
1	A	644	SER	2.8
1	A	677	ARG	2.8
1	A	643	ALA	2.8
1	A	736	ARG	2.7
1	A	638	ILE	2.7
1	A	688	GLU	2.7
1	A	667	PHE	2.6
1	A	642	THR	2.3
1	A	299	TRP	2.3
1	A	678	LEU	2.2
1	A	673	MET	2.2
1	A	646	MET	2.2
1	A	637	ASP	2.1
1	A	639	HIS	2.1
1	A	607	VAL	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DOC	B	112	18/19	0.97	0.10	16,21,24,25	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	A	834	6/6	0.88	0.13	31,35,43,43	6
6	MN	B	3	1/1	0.90	0.15	38,38,38,38	1
5	GOL	A	1	6/6	0.92	0.13	27,29,31,31	0
4	DCT	A	113	27/27	0.95	0.10	23,28,34,36	0
7	MG	A	2	1/1	0.96	0.07	29,29,29,29	0
6	MN	C	4	1/1	0.97	0.05	53,53,53,53	0
6	MN	A	6	1/1	0.98	0.13	46,46,46,46	0
6	MN	A	5	1/1	0.99	0.07	38,38,38,38	0
6	MN	A	833	1/1	1.00	0.07	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers ⓘ

There are no such residues in this entry.