



wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 07:25 am BST

PDB ID : 1PZU
Title : An asymmetric NFAT1-RHR homodimer on a pseudo-palindromic, Kappa-B site
Authors : Jin, L.; Sliz, P.; Chen, L.; Macian, F.; Rao, A.; Hogan, P.G.; Harrison, S.C.
Deposited on : 2003-07-14
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	T	14	
1	V	14	
1	X	14	
2	W	14	
2	Y	14	
2	Z	14	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	B	301	<div><div></div><div>7%</div><div>44%</div><div>42%</div><div>6%</div><div>8%</div></div>
3	D	301	<div><div></div><div>12%</div><div>44%</div><div>41%</div><div>7%</div><div>8%</div></div>
3	H	301	<div><div></div><div>8%</div><div>44%</div><div>41%</div><div>6%</div><div>8%</div></div>
3	I	301	<div><div></div><div>%</div><div>45%</div><div>41%</div><div>6%</div><div>8%</div></div>
3	L	301	<div><div></div><div>4%</div><div>44%</div><div>41%</div><div>7%</div><div>8%</div></div>
3	M	301	<div><div></div><div>2%</div><div>46%</div><div>38%</div><div>7%</div><div>8%</div></div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 14952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*TP*TP*GP*AP*GP*GP*AP*AP*TP*TP*TP*CP*CP*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	14	Total	C	N	O	P	0	0	0
			285	138	51	83	13			
1	V	14	Total	C	N	O	P	0	0	0
			285	138	51	83	13			
1	T	14	Total	C	N	O	P	37	0	0
			285	138	51	83	13			

- Molecule 2 is a DNA chain called 5'-D(*AP*AP*TP*GP*GP*AP*AP*AP*TP*TP*CP*CP*TP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Y	14	Total	C	N	O	P	0	0	0
			283	137	52	81	13			
2	Z	14	Total	C	N	O	P	0	0	0
			283	137	52	81	13			
2	W	14	Total	C	N	O	P	0	0	0
			283	137	52	81	13			

- Molecule 3 is a protein called Nuclear factor of activated T-cells, cytoplasmic 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	276	Total	C	N	O	S	0	0	0
			2208	1386	401	412	9			
3	D	276	Total	C	N	O	S	0	0	0
			2208	1386	401	412	9			
3	H	276	Total	C	N	O	S	0	0	0
			2208	1386	401	412	9			
3	I	276	Total	C	N	O	S	0	0	0
			2208	1386	401	412	9			
3	L	276	Total	C	N	O	S	0	0	0
			2208	1386	401	412	9			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	276	Total	C	N	O	S	0	0	0
			2208	1386	401	412	9			

There are 108 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	378	MET	-	CLONING ARTIFACT	UNP Q13469
B	379	ARG	-	CLONING ARTIFACT	UNP Q13469
B	380	GLY	-	CLONING ARTIFACT	UNP Q13469
B	381	SER	-	CLONING ARTIFACT	UNP Q13469
B	382	HIS	-	EXPRESSION TAG	UNP Q13469
B	383	HIS	-	EXPRESSION TAG	UNP Q13469
B	384	HIS	-	EXPRESSION TAG	UNP Q13469
B	385	HIS	-	EXPRESSION TAG	UNP Q13469
B	386	HIS	-	EXPRESSION TAG	UNP Q13469
B	387	HIS	-	EXPRESSION TAG	UNP Q13469
B	388	THR	-	CLONING ARTIFACT	UNP Q13469
B	389	ASP	-	CLONING ARTIFACT	UNP Q13469
B	390	PRO	-	CLONING ARTIFACT	UNP Q13469
B	391	HIS	-	CLONING ARTIFACT	UNP Q13469
B	392	ALA	-	CLONING ARTIFACT	UNP Q13469
B	393	SER	-	CLONING ARTIFACT	UNP Q13469
B	394	SER	-	CLONING ARTIFACT	UNP Q13469
B	395	VAL	-	CLONING ARTIFACT	UNP Q13469
D	378	MET	-	CLONING ARTIFACT	UNP Q13469
D	379	ARG	-	CLONING ARTIFACT	UNP Q13469
D	380	GLY	-	CLONING ARTIFACT	UNP Q13469
D	381	SER	-	CLONING ARTIFACT	UNP Q13469
D	382	HIS	-	EXPRESSION TAG	UNP Q13469
D	383	HIS	-	EXPRESSION TAG	UNP Q13469
D	384	HIS	-	EXPRESSION TAG	UNP Q13469
D	385	HIS	-	EXPRESSION TAG	UNP Q13469
D	386	HIS	-	EXPRESSION TAG	UNP Q13469
D	387	HIS	-	EXPRESSION TAG	UNP Q13469
D	388	THR	-	CLONING ARTIFACT	UNP Q13469
D	389	ASP	-	CLONING ARTIFACT	UNP Q13469
D	390	PRO	-	CLONING ARTIFACT	UNP Q13469
D	391	HIS	-	CLONING ARTIFACT	UNP Q13469
D	392	ALA	-	CLONING ARTIFACT	UNP Q13469
D	393	SER	-	CLONING ARTIFACT	UNP Q13469
D	394	SER	-	CLONING ARTIFACT	UNP Q13469
D	395	VAL	-	CLONING ARTIFACT	UNP Q13469

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	378	MET	-	CLONING ARTIFACT	UNP Q13469
H	379	ARG	-	CLONING ARTIFACT	UNP Q13469
H	380	GLY	-	CLONING ARTIFACT	UNP Q13469
H	381	SER	-	CLONING ARTIFACT	UNP Q13469
H	382	HIS	-	EXPRESSION TAG	UNP Q13469
H	383	HIS	-	EXPRESSION TAG	UNP Q13469
H	384	HIS	-	EXPRESSION TAG	UNP Q13469
H	385	HIS	-	EXPRESSION TAG	UNP Q13469
H	386	HIS	-	EXPRESSION TAG	UNP Q13469
H	387	HIS	-	EXPRESSION TAG	UNP Q13469
H	388	THR	-	CLONING ARTIFACT	UNP Q13469
H	389	ASP	-	CLONING ARTIFACT	UNP Q13469
H	390	PRO	-	CLONING ARTIFACT	UNP Q13469
H	391	HIS	-	CLONING ARTIFACT	UNP Q13469
H	392	ALA	-	CLONING ARTIFACT	UNP Q13469
H	393	SER	-	CLONING ARTIFACT	UNP Q13469
H	394	SER	-	CLONING ARTIFACT	UNP Q13469
H	395	VAL	-	CLONING ARTIFACT	UNP Q13469
I	378	MET	-	CLONING ARTIFACT	UNP Q13469
I	379	ARG	-	CLONING ARTIFACT	UNP Q13469
I	380	GLY	-	CLONING ARTIFACT	UNP Q13469
I	381	SER	-	CLONING ARTIFACT	UNP Q13469
I	382	HIS	-	EXPRESSION TAG	UNP Q13469
I	383	HIS	-	EXPRESSION TAG	UNP Q13469
I	384	HIS	-	EXPRESSION TAG	UNP Q13469
I	385	HIS	-	EXPRESSION TAG	UNP Q13469
I	386	HIS	-	EXPRESSION TAG	UNP Q13469
I	387	HIS	-	EXPRESSION TAG	UNP Q13469
I	388	THR	-	CLONING ARTIFACT	UNP Q13469
I	389	ASP	-	CLONING ARTIFACT	UNP Q13469
I	390	PRO	-	CLONING ARTIFACT	UNP Q13469
I	391	HIS	-	CLONING ARTIFACT	UNP Q13469
I	392	ALA	-	CLONING ARTIFACT	UNP Q13469
I	393	SER	-	CLONING ARTIFACT	UNP Q13469
I	394	SER	-	CLONING ARTIFACT	UNP Q13469
I	395	VAL	-	CLONING ARTIFACT	UNP Q13469
L	378	MET	-	CLONING ARTIFACT	UNP Q13469
L	379	ARG	-	CLONING ARTIFACT	UNP Q13469
L	380	GLY	-	CLONING ARTIFACT	UNP Q13469
L	381	SER	-	CLONING ARTIFACT	UNP Q13469
L	382	HIS	-	EXPRESSION TAG	UNP Q13469
L	383	HIS	-	EXPRESSION TAG	UNP Q13469

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	384	HIS	-	EXPRESSION TAG	UNP Q13469
L	385	HIS	-	EXPRESSION TAG	UNP Q13469
L	386	HIS	-	EXPRESSION TAG	UNP Q13469
L	387	HIS	-	EXPRESSION TAG	UNP Q13469
L	388	THR	-	CLONING ARTIFACT	UNP Q13469
L	389	ASP	-	CLONING ARTIFACT	UNP Q13469
L	390	PRO	-	CLONING ARTIFACT	UNP Q13469
L	391	HIS	-	CLONING ARTIFACT	UNP Q13469
L	392	ALA	-	CLONING ARTIFACT	UNP Q13469
L	393	SER	-	CLONING ARTIFACT	UNP Q13469
L	394	SER	-	CLONING ARTIFACT	UNP Q13469
L	395	VAL	-	CLONING ARTIFACT	UNP Q13469
M	378	MET	-	CLONING ARTIFACT	UNP Q13469
M	379	ARG	-	CLONING ARTIFACT	UNP Q13469
M	380	GLY	-	CLONING ARTIFACT	UNP Q13469
M	381	SER	-	CLONING ARTIFACT	UNP Q13469
M	382	HIS	-	EXPRESSION TAG	UNP Q13469
M	383	HIS	-	EXPRESSION TAG	UNP Q13469
M	384	HIS	-	EXPRESSION TAG	UNP Q13469
M	385	HIS	-	EXPRESSION TAG	UNP Q13469
M	386	HIS	-	EXPRESSION TAG	UNP Q13469
M	387	HIS	-	EXPRESSION TAG	UNP Q13469
M	388	THR	-	CLONING ARTIFACT	UNP Q13469
M	389	ASP	-	CLONING ARTIFACT	UNP Q13469
M	390	PRO	-	CLONING ARTIFACT	UNP Q13469
M	391	HIS	-	CLONING ARTIFACT	UNP Q13469
M	392	ALA	-	CLONING ARTIFACT	UNP Q13469
M	393	SER	-	CLONING ARTIFACT	UNP Q13469
M	394	SER	-	CLONING ARTIFACT	UNP Q13469
M	395	VAL	-	CLONING ARTIFACT	UNP Q13469

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*TP*TP*GP*AP*GP*GP*AP*AP*TP*TP*TP*CP*CP*A)-3'

Chain X: 



- Molecule 1: 5'-D(*TP*TP*GP*AP*GP*GP*AP*AP*TP*TP*TP*CP*CP*A)-3'

Chain V: 



- Molecule 1: 5'-D(*TP*TP*GP*AP*GP*GP*AP*AP*TP*TP*TP*CP*CP*A)-3'

Chain T: 





- Molecule 2: 5'-D(*AP*AP*TP*GP*GP*AP*AP*AP*TP*TP*CP*CP*TP*C)-3'

Chain Y: 





- Molecule 2: 5'-D(*AP*AP*TP*GP*GP*AP*AP*AP*TP*TP*CP*CP*TP*C)-3'


Chain Z: 




- Molecule 2: 5'-D(*AP*AP*TP*GP*GP*AP*AP*AP*TP*TP*CP*CP*TP*C)-3'

Chain W: 






• Molecule 3: Nuclear factor of activated T-cells, cytoplasmic 2




• Molecule 3: Nuclear factor of activated T-cells, cytoplasmic 2





● Molecule 3: Nuclear factor of activated T-cells, cytoplasmic 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.70Å 122.90Å 241.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.78 – 3.10 49.11 – 3.10	Depositor EDS
% Data completeness (in resolution range)	82.4 (29.78-3.10) 82.3 (49.11-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 3.12Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.296 , 0.319 0.284 , 0.301	Depositor DCC
R_{free} test set	1857 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	65.9	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 65.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	14952	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	T	0.90	0/319	0.96	2/491 (0.4%)
1	V	0.84	1/319 (0.3%)	0.86	0/491
1	X	0.85	0/319	0.98	2/491 (0.4%)
2	W	0.90	0/317	1.02	1/487 (0.2%)
2	Y	0.84	0/317	0.95	1/487 (0.2%)
2	Z	0.91	0/317	0.96	1/487 (0.2%)
3	B	0.45	0/2255	0.66	0/3048
3	D	0.43	0/2255	0.66	0/3048
3	H	0.42	0/2255	0.66	0/3048
3	I	0.42	0/2255	0.66	0/3048
3	L	0.41	0/2255	0.66	0/3048
3	M	0.42	0/2255	0.66	0/3048
All	All	0.50	1/15438 (0.0%)	0.71	7/21222 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	T	0	1
1	X	0	1
2	W	0	4
2	Y	0	1
2	Z	0	1
All	All	0	8

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	V	5	DG	C5-C6	-5.68	1.36	1.42

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	T	3	DG	N9-C1'-C2'	5.75	123.52	112.60
1	X	4	DA	N9-C1'-C2'	5.61	123.26	112.60
2	Y	4	DG	C5'-C4'-C3'	-5.44	104.30	114.10
2	W	5	DG	O5'-P-OP1	-5.35	100.88	105.70
1	X	9	DT	OP1-P-O3'	5.16	116.56	105.20

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	T	4	DA	Sidechain
2	W	1	DA	Sidechain
1	X	4	DA	Sidechain
2	Y	4	DG	Sidechain
2	Z	14	DC	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	T	285	0	161	18	0
1	V	285	0	161	32	0
1	X	285	0	161	33	0
2	W	283	0	160	28	0
2	Y	283	0	160	20	0
2	Z	283	0	160	25	0
3	B	2208	0	2214	168	1
3	D	2208	0	2214	174	0
3	H	2208	0	2214	163	0
3	I	2208	0	2214	167	1
3	L	2208	0	2214	170	0
3	M	2208	0	2214	163	0
All	All	14952	0	14247	1106	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 1106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:W:2:DA:H2''	2:W:3:DT:H5''	1.18	1.15
1:X:5:DG:H2''	1:X:6:DG:H5'	1.25	1.12
2:Y:4:DG:H2''	2:Y:5:DG:H5'	1.28	1.11
1:V:3:DG:C2'	1:V:4:DA:H5''	1.82	1.10
2:Y:2:DA:H2''	2:Y:3:DT:H5''	1.29	1.08

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:649:HIS:NE2	3:I:602:ASN:ND2[4_556]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	B	272/301 (90%)	213 (78%)	45 (16%)	14 (5%)	2	13
3	D	272/301 (90%)	212 (78%)	48 (18%)	12 (4%)	2	15
3	H	272/301 (90%)	215 (79%)	43 (16%)	14 (5%)	2	13
3	I	272/301 (90%)	212 (78%)	47 (17%)	13 (5%)	2	14
3	L	272/301 (90%)	211 (78%)	47 (17%)	14 (5%)	2	13
3	M	272/301 (90%)	214 (79%)	45 (16%)	13 (5%)	2	14
All	All	1632/1806 (90%)	1277 (78%)	275 (17%)	80 (5%)	2	14

5 of 80 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	406	GLY
3	B	467	ILE
3	B	590	VAL
3	B	628	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	D	406	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	B	247/269 (92%)	232 (94%)	15 (6%)	18	49
3	D	247/269 (92%)	233 (94%)	14 (6%)	20	52
3	H	247/269 (92%)	233 (94%)	14 (6%)	20	52
3	I	247/269 (92%)	233 (94%)	14 (6%)	20	52
3	L	247/269 (92%)	233 (94%)	14 (6%)	20	52
3	M	247/269 (92%)	232 (94%)	15 (6%)	18	49
All	All	1482/1614 (92%)	1396 (94%)	86 (6%)	20	51

5 of 86 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	577	LEU
3	I	485	THR
3	M	508	ASN
3	H	621	ILE
3	I	426	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 84 such sidechains are listed below:

Mol	Chain	Res	Type
3	H	583	GLN
3	I	495	ASN
3	M	583	GLN
3	H	595	GLN
3	H	647	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	T	12/14 (85%)	-0.25	0	100	100	18, 23, 30, 31	0
1	V	14/14 (100%)	-0.18	0	100	100	25, 30, 38, 38	0
1	X	14/14 (100%)	-0.18	0	100	100	21, 28, 34, 47	0
2	W	14/14 (100%)	-0.23	0	100	100	19, 26, 33, 38	0
2	Y	14/14 (100%)	-0.07	0	100	100	21, 27, 42, 48	0
2	Z	14/14 (100%)	-0.11	0	100	100	22, 31, 41, 41	0
3	B	276/301 (91%)	0.47	20 (7%)	15	6	8, 60, 142, 160	0
3	D	276/301 (91%)	0.68	36 (13%)	3	1	14, 68, 153, 172	0
3	H	276/301 (91%)	0.57	24 (8%)	10	4	45, 72, 106, 114	0
3	I	276/301 (91%)	0.30	4 (1%)	75	56	21, 70, 90, 107	0
3	L	276/301 (91%)	0.39	13 (4%)	31	15	33, 79, 98, 112	0
3	M	276/301 (91%)	0.28	6 (2%)	62	41	14, 66, 87, 106	0
All	All	1738/1890 (91%)	0.42	103 (5%)	22	10	8, 69, 141, 172	0

The worst 5 of 103 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	678	VAL	8.5
3	D	647	ASN	8.1
3	B	596	MET	5.5
3	D	656	VAL	5.3
3	D	649	HIS	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.