



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 07:10 AM BST

PDB ID : 6PZE
Title : Crystal structure of human NA-45 Fab in complex with neuraminidase Y169aH mutant from A/Shanghai/2/2013 (H7N9)
Authors : Zhu, X.; Wilson, I.A.
Deposited on : 2019-07-31
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	<div> <div>92%</div> <div>6%</div> </div>
2	L	217	<div>8%</div> <div>82%</div> <div>15%</div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	388	Total	C	N	O	S	0	0	0
			3053	1899	543	588	23			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	77	GLY	-	expression tag	UNP V9NZ28
A	78	SER	-	expression tag	UNP V9NZ28
A	79	PRO	-	expression tag	UNP V9NZ28
A	80	SER	-	expression tag	UNP V9NZ28
A	81	ARG	-	expression tag	UNP V9NZ28
A	169A	HIS	TYR	engineered mutation	UNP V9NZ28


- Molecule 2 is a protein called NA-45 FAB LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	211	Total	C	N	O	S	0	0	0
			1565	982	259	320	4			

- Molecule 3 is a protein called NA-45 FAB HEAVY CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	218	Total	C	N	O	S	0	0	0
			1642	1050	264	324	4			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	B	10	Total	C	N	O	0	0	0
			116	64	2	50			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	366	Total	O	0	0
			366	366		
7	L	100	Total	O	0	0
			100	100		
7	H	124	Total	O	0	0
			124	124		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Neuraminidase



- Molecule 2: NA-45 FAB LIGHT CHAIN



- Molecule 3: NA-45 FAB HEAVY CHAIN



- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:

100%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	161.27 Å 161.27 Å 87.38 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.79 – 2.30 45.79 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.2 (45.79-2.30) 98.2 (45.79-2.29)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.29 Å)	Xtriage
Refinement program	PHENIX (1.12_2829)	Depositor
R, R_{free}	0.156 , 0.197 0.157 , 0.197	Depositor DCC
R_{free} test set	2564 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6995	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/3135	0.61	1/4268 (0.0%)
2	L	0.33	0/1604	0.56	1/2195 (0.0%)
3	H	0.35	0/1688	0.59	0/2314
All	All	0.37	0/6427	0.59	2/8777 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	224	ARG	NE-CZ-NH1	-5.75	117.43	120.30
2	L	100	THR	C-N-CA	-5.07	111.66	122.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3053	0	2876	13	0
2	L	1565	0	1524	18	0
3	H	1642	0	1607	25	0
4	B	116	0	97	0	0
5	C	28	0	25	1	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	366	0	0	1	0
7	H	124	0	0	0	0
7	L	100	0	0	0	0
All	All	6995	0	6129	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:210:ARG:HD2	3:H:212:GLU:OE2	1.72	0.89
3:H:43:LYS:HD2	3:H:44:GLY:H	1.40	0.87
3:H:210:ARG:NH1	3:H:212:GLU:HB2	2.02	0.74
3:H:211:VAL:O	3:H:212:GLU:HG3	1.91	0.70
1:A:212:VAL:HG21	1:A:260:GLY:HA3	1.78	0.63
3:H:59:TYR:HE1	3:H:69:ILE:HG12	1.65	0.62
3:H:191:THR:HG23	3:H:192:GLN:H	1.64	0.61
2:L:194:GLN:NE2	2:L:203:GLU:OE1	2.34	0.61
1:A:97:TRP:H	1:A:395:GLN:HE22	1.47	0.60
3:H:160:THR:O	3:H:163:VAL:HG12	2.01	0.59
3:H:195:ILE:HD12	3:H:209:LYS:C	2.24	0.58
3:H:43:LYS:CD	3:H:44:GLY:H	2.13	0.58
2:L:39:LEU:HB2	2:L:42:THR:HG21	1.84	0.58
3:H:123:PRO:HB2	3:H:211:VAL:HG13	1.87	0.56
1:A:121:TYR:CG	1:A:228:SER:HA	2.42	0.55
3:H:210:ARG:HH12	3:H:212:GLU:HB2	1.70	0.55
2:L:110:LYS:HD3	2:L:141:PRO:HD3	1.89	0.55
2:L:37:GLN:HB2	2:L:47:LEU:HD11	1.88	0.55
3:H:100(H):TRP:CZ3	3:H:101:ASP:HB3	2.42	0.54
3:H:163:VAL:HA	3:H:182:VAL:HG12	1.90	0.54
3:H:184:VAL:HG11	3:H:194:TYR:OH	2.08	0.53
2:L:189:ARG:HG2	2:L:189:ARG:HH11	1.75	0.52
1:A:283:GLU:OE2	1:A:285:THR:OG1	2.28	0.51
1:A:437:TRP:CE3	2:L:53:LYS:HE2	2.48	0.49
3:H:162:GLY:O	3:H:182:VAL:HA	2.14	0.48
2:L:39:LEU:HB2	2:L:42:THR:CG2	2.43	0.48
1:A:378:LYS:HE2	1:A:392:ILE:HD11	1.96	0.48
2:L:80:THR:OG1	2:L:170:ASN:ND2	2.47	0.47
3:H:138:LEU:HD12	3:H:211:VAL:HB	1.96	0.47
1:A:465:GLU:CD	1:A:465:GLU:H	2.18	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:18:LYS:HG3	2:L:76:THR:HG22	1.97	0.46
3:H:195:ILE:HD12	3:H:210:ARG:N	2.31	0.46
1:A:367:SER:HB2	1:A:400:ASN:HD21	1.81	0.46
2:L:55:PRO:HD2	2:L:58:ILE:HG13	1.97	0.45
2:L:120:PRO:HD3	2:L:132:LEU:CD2	2.47	0.45
2:L:19:VAL:HG22	2:L:75:ILE:HB	1.99	0.44
2:L:61:ARG:NH2	2:L:82:ASP:OD2	2.51	0.44
3:H:36:TRP:HD1	3:H:69:ILE:HD12	1.82	0.44
5:C:1:NAG:O4	5:C:2:NAG:O7	2.35	0.44
3:H:36:TRP:CD1	3:H:69:ILE:HD12	2.52	0.44
3:H:100:TRP:CG	3:H:100(A):GLU:N	2.86	0.43
3:H:105:GLN:HG2	3:H:106:GLY:N	2.33	0.43
3:H:210:ARG:HH11	3:H:212:GLU:CD	2.22	0.43
2:L:27(B):ASN:OD1	2:L:28:ILE:N	2.44	0.43
2:L:189:ARG:NH1	2:L:189:ARG:HG2	2.33	0.43
2:L:196:THR:HG22	2:L:201:THR:HG23	2.01	0.43
2:L:4:LEU:HA	2:L:4:LEU:HD23	1.77	0.43
1:A:172:ARG:NE	1:A:209:ARG:NH2	2.68	0.42
2:L:3:VAL:O	2:L:25:GLY:HA2	2.19	0.42
3:H:13:LYS:HB2	3:H:16:GLU:OE2	2.20	0.42
1:A:152:ARG:HE	1:A:198:ASN:HD21	1.67	0.41
1:A:188:SER:HB2	1:A:207:TYR:CZ	2.55	0.41
1:A:82:ARG:N	7:A:619:HOH:O	2.54	0.41
1:A:127:ASP:OD1	1:A:127:ASP:N	2.54	0.41
3:H:43:LYS:HD2	3:H:44:GLY:N	2.21	0.40
3:H:66:ARG:HH22	3:H:86:ASP:CG	2.25	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/393 (98%)	367 (95%)	18 (5%)	1 (0%)	41	50
2	L	209/217 (96%)	199 (95%)	9 (4%)	1 (0%)	29	35
3	H	214/237 (90%)	205 (96%)	9 (4%)	0	100	100
All	All	809/847 (96%)	771 (95%)	36 (4%)	2 (0%)	47	58

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	40	PRO
1	A	222	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/343 (99%)	338 (100%)	1 (0%)	92	97
2	L	177/183 (97%)	176 (99%)	1 (1%)	86	94
3	H	190/207 (92%)	187 (98%)	3 (2%)	62	78
All	All	706/733 (96%)	701 (99%)	5 (1%)	84	92

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	456	TRP
2	L	27(A)	SER
3	H	28	SER
3	H	164	HIS
3	H	210	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	198	ASN
1	A	345	ASN
1	A	395	GLN
1	A	400	ASN
1	A	455	GLN
2	L	170	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	B	1	1,4	14,14,15	1.47	3 (21%)	17,19,21	1.18	2 (11%)
4	MAN	B	10	4	11,11,12	0.86	0	15,15,17	1.17	1 (6%)
4	NAG	B	2	4	14,14,15	0.50	0	17,19,21	0.93	1 (5%)
4	BMA	B	3	4	11,11,12	0.80	0	15,15,17	0.87	0
4	MAN	B	4	4	11,11,12	0.92	1 (9%)	15,15,17	1.55	4 (26%)
4	MAN	B	5	4	11,11,12	0.93	0	15,15,17	1.06	1 (6%)
4	MAN	B	6	4	11,11,12	0.78	0	15,15,17	1.44	1 (6%)
4	MAN	B	7	4	11,11,12	0.77	0	15,15,17	0.99	2 (13%)
4	MAN	B	8	4	11,11,12	0.87	0	15,15,17	1.18	2 (13%)
4	MAN	B	9	4	11,11,12	0.97	0	15,15,17	1.05	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	C	1	1,5	14,14,15	0.72	1 (7%)	17,19,21	0.81	1 (5%)
5	NAG	C	2	5	14,14,15	1.13	1 (7%)	17,19,21	1.27	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1	1,4	-	2/6/23/26	0/1/1/1
4	MAN	B	10	4	-	0/2/19/22	0/1/1/1
4	NAG	B	2	4	-	0/6/23/26	0/1/1/1
4	BMA	B	3	4	-	0/2/19/22	0/1/1/1
4	MAN	B	4	4	-	0/2/19/22	0/1/1/1
4	MAN	B	5	4	-	0/2/19/22	0/1/1/1
4	MAN	B	6	4	-	0/2/19/22	0/1/1/1
4	MAN	B	7	4	-	0/2/19/22	0/1/1/1
4	MAN	B	8	4	-	0/2/19/22	0/1/1/1
4	MAN	B	9	4	-	0/2/19/22	0/1/1/1
5	NAG	C	1	1,5	-	4/6/23/26	0/1/1/1
5	NAG	C	2	5	-	1/6/23/26	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	2	NAG	C1-C2	4.07	1.58	1.52
4	B	1	NAG	O5-C1	-3.35	1.38	1.43
4	B	1	NAG	C3-C2	3.02	1.58	1.52
5	C	1	NAG	C1-C2	2.46	1.56	1.52
4	B	4	MAN	O5-C5	2.12	1.47	1.43
4	B	1	NAG	O7-C7	2.03	1.27	1.23

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	6	MAN	C1-O5-C5	4.33	118.06	112.19
5	C	2	NAG	C2-N2-C7	4.20	128.89	122.90
4	B	10	MAN	C1-O5-C5	3.67	117.17	112.19
4	B	8	MAN	O2-C2-C3	-3.39	103.35	110.14
4	B	4	MAN	O3-C3-C2	3.11	115.95	109.99
4	B	5	MAN	C1-O5-C5	2.83	116.02	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	9	MAN	C1-O5-C5	2.63	115.76	112.19
4	B	1	NAG	C1-O5-C5	2.61	115.73	112.19
4	B	4	MAN	C1-O5-C5	2.61	115.72	112.19
4	B	1	NAG	C3-C4-C5	-2.52	105.75	110.24
4	B	4	MAN	C3-C4-C5	-2.44	105.88	110.24
4	B	9	MAN	O2-C2-C3	-2.41	105.32	110.14
5	C	1	NAG	C1-O5-C5	2.38	115.41	112.19
4	B	7	MAN	C1-O5-C5	2.36	115.40	112.19
4	B	4	MAN	C1-C2-C3	-2.25	106.90	109.67
4	B	8	MAN	C1-O5-C5	2.13	115.08	112.19
4	B	2	NAG	C1-O5-C5	2.08	115.01	112.19
4	B	7	MAN	O2-C2-C3	-2.06	106.02	110.14

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	2	NAG	C1-C2-N2-C7
5	C	1	NAG	C8-C7-N2-C2
5	C	1	NAG	O7-C7-N2-C2
4	B	1	NAG	C4-C5-C6-O6
5	C	1	NAG	C4-C5-C6-O6
4	B	1	NAG	O5-C5-C6-O6
5	C	1	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	2	NAG	1	0
5	C	1	NAG	1	0

5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/393 (98%)	-0.38	0 100 100	13, 20, 34, 50	0
2	L	211/217 (97%)	0.21	18 (8%) 10 14	18, 40, 77, 97	0
3	H	218/237 (91%)	0.22	26 (11%) 4 6	20, 35, 101, 143	0
All	All	817/847 (96%)	-0.07	44 (5%) 25 32	13, 26, 76, 143	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	189	LEU	8.7
3	H	158	ALA	5.6
3	H	190	GLY	5.4
3	H	187	SER	5.1
2	L	207	ALA	5.0
3	H	191	THR	4.6
3	H	195	ILE	4.5
3	H	193	THR	4.5
3	H	212	GLU	3.8
3	H	194	TYR	3.8
3	H	210	ARG	3.7
3	H	185	PRO	3.7
3	H	192	GLN	3.6
2	L	189	ARG	3.2
2	L	187	SER	3.2
3	H	186	SER	3.2
2	L	150	ALA	3.1
3	H	184	VAL	3.1
2	L	186	LYS	3.0
3	H	188	SER	2.9
3	H	138	LEU	2.7
2	L	181	THR	2.7
3	H	211	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	H	183	THR	2.5
2	L	153	SER	2.5
2	L	156	LYS	2.4
3	H	159	LEU	2.4
2	L	40	PRO	2.3
3	H	135	THR	2.2
2	L	183	GLU	2.2
2	L	122	SER	2.2
3	H	125	ALA	2.2
2	L	2	SER	2.2
3	H	197	ASN	2.2
2	L	119	PRO	2.2
2	L	151	ASP	2.2
3	H	136	ALA	2.1
2	L	190	SER	2.1
2	L	191	TYR	2.1
3	H	181	VAL	2.1
3	H	153	SER	2.1
2	L	125	LEU	2.1
2	L	149	LYS	2.0
3	H	157	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	C	2	14/15	0.69	0.33	81,91,101,108	0
5	NAG	C	1	14/15	0.79	0.25	58,71,81,84	0
4	MAN	B	9	11/12	0.85	0.25	48,58,63,63	0
4	MAN	B	10	11/12	0.90	0.21	50,58,63,64	0
4	MAN	B	8	11/12	0.94	0.14	38,43,52,56	0
4	MAN	B	6	11/12	0.95	0.09	22,26,30,32	0
4	MAN	B	5	11/12	0.95	0.16	28,30,33,33	0

Continued on next page...

Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	BMA	B	3	11/12	0.96	0.12	20,26,32,36	0
4	MAN	B	7	11/12	0.96	0.16	29,34,41,42	0
4	NAG	B	2	14/15	0.96	0.11	23,28,37,53	0
4	NAG	B	1	14/15	0.96	0.11	24,28,34,39	0
4	MAN	B	4	11/12	0.97	0.12	22,27,39,48	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	CA	A	513	1/1	0.96	0.09	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.