



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 02:48 am BST

PDB ID : 3Q07  
Title : CTX-M-9 S70G in complex with piperacillin  
Authors : Delmas, J.; Leyssne, D.; Robin, F.; Coignoux, A.; Bonnet, R.  
Deposited on : 2010-12-15  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>%</span> <span>91%</span> <span>8%</span> </div> </div>
1	B	263	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 2%, yellow 1%, green 95%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>2%</span> <span>90%</span> <span>9%</span> </div> </div>



In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total 2033	C 1266	N 362	O 399	S 6	0	15	0
1	B	260	Total 2025	C 1269	N 356	O 393	S 7	0	18	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	70	GLY	SER	ENGINEERED MUTATION	UNP Q9L5C8
B	70	GLY	SER	ENGINEERED MUTATION	UNP Q9L5C8

- Molecule 2 is Piperacillin (three-letter code: WPP) (formula:  $\text{C}_{23}\text{H}_{27}\text{N}_5\text{O}_7\text{S}$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	1
			36	23	5	7	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	S	0	1
			36	23	5	7	1		

- Molecule 3 is Hydrolyzed piperacillin (three-letter code: YPP) (formula:  $C_{23}H_{29}N_5O_8S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	1
			37	23	5	8	1		
3	B	1	Total	C	N	O	S	0	1
			37	23	5	8	1		
3	B	1	Total	C	N	O	S	0	0
			37	23	5	8	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	376	Total O 378 378	0	2
4	B	411	Total O 412 412	0	1



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Beta-lactamase



#### • Molecule 1: Beta-lactamase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.13 Å 106.34 Å 47.61 Å 90.00° 101.90° 90.00°	Depositor
Resolution (Å)	19.65 – 1.50 19.47 – 1.50	Depositor EDS
% Data completeness (in resolution range)	94.5 (19.65-1.50) 94.5 (19.47-1.50)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	9.22 (at 1.50 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.146 , 0.175 0.145 , 0.175	Depositor DCC
$R_{free}$ test set	3328 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	5.9	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 50.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5031	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	9.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: YPP, WPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	0/2110	0.59	0/2868
1	B	0.48	0/2108	0.58	0/2865
All	All	0.49	0/4218	0.58	0/5733

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2033	0	2092	31	0
1	B	2025	0	2104	33	0
2	A	36	0	26	1	0
2	B	36	0	26	2	0
3	A	37	0	27	2	0
3	B	74	0	54	2	0
4	A	378	0	0	12	0
4	B	412	0	0	10	0
All	All	5031	0	4329	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:96[B]:GLU:OE1	1:B:98:LYS:HE3	1.33	1.23
1:A:27:THR:CG2	1:A:32:GLN:HE21	1.54	1.21
1:B:186[B]:MET:HE3	1:B:189[B]:THR:HG22	1.28	1.09
1:B:186[B]:MET:CE	1:B:189[B]:THR:CG2	2.37	1.02
1:B:186[B]:MET:HE3	1:B:189[B]:THR:CG2	1.89	1.02
1:B:186[B]:MET:HE1	1:B:189[B]:THR:HG21	1.46	0.98
1:B:50:ILE:HB	1:B:258[B]:VAL:HG12	1.52	0.92
1:B:269:ASN:HB3	4:B:419:HOH:O	1.70	0.92
1:B:186[B]:MET:CE	1:B:189[B]:THR:HG21	2.03	0.88
1:A:27:THR:HG22	1:A:32:GLN:HE21	1.37	0.88
1:A:27:THR:CG2	1:A:32:GLN:NE2	2.37	0.86
1:A:27:THR:HG23	1:A:32:GLN:HG3	1.62	0.80
1:B:186[B]:MET:HE1	1:B:189[B]:THR:CG2	2.06	0.78
1:B:96[B]:GLU:CD	1:B:98:LYS:HE3	2.05	0.78
1:B:267[B]:GLN:HG2	1:B:270:ALA:HB2	1.71	0.71
1:B:50:ILE:HB	1:B:258[B]:VAL:CG1	2.19	0.71
1:A:42[A]:GLY:HA2	4:A:518:HOH:O	1.89	0.70
1:A:27:THR:HG21	1:A:32:GLN:HE21	1.55	0.70
1:A:57[B]:GLN:NE2	4:A:499:HOH:O	2.26	0.69
1:B:267[B]:GLN:HG3	4:B:562:HOH:O	1.94	0.68
1:A:87[A]:GLN:NE2	1:A:93:GLN:HE22	1.93	0.66
1:A:42[A]:GLY:CA	4:A:518:HOH:O	2.43	0.66
3:B:401[B]:YPP:CAW	3:B:401[B]:YPP:HANB	2.27	0.64
1:A:27:THR:CG2	1:A:27:THR:O	2.46	0.63
1:A:42[A]:GLY:N	4:A:518:HOH:O	2.31	0.62
1:B:111:LYS:HE3	4:B:730:HOH:O	2.00	0.61
1:B:267[B]:GLN:CG	4:B:562:HOH:O	2.48	0.61
1:B:111:LYS:CE	4:B:730:HOH:O	2.50	0.58
1:A:27:THR:O	1:A:27:THR:HG22	2.04	0.57
1:B:269:ASN:CB	4:B:419:HOH:O	2.42	0.57
3:B:401[B]:YPP:HANB	3:B:401[B]:YPP:CAX	2.34	0.57
1:A:27:THR:HG23	1:A:32:GLN:CG	2.35	0.56
1:B:29[B]:ALA:N	4:B:364:HOH:O	2.39	0.55
1:A:212[B]:LYS:NZ	4:A:491:HOH:O	2.20	0.54
1:B:48:ALA:HB3	1:B:260[B]:VAL:HG22	1.88	0.54
1:A:27:THR:HG23	1:A:32:GLN:HE21	1.64	0.53
1:B:267[B]:GLN:CD	1:B:267[B]:GLN:H	2.11	0.53
1:B:48:ALA:HB3	1:B:260[B]:VAL:CG2	2.38	0.53
1:B:237[A]:SER:OG	2:B:400[A]:WPP:HAE	2.08	0.52
1:B:96[B]:GLU:OE1	1:B:98:LYS:CE	2.29	0.52

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:GLN:HB3	4:A:411:HOH:O	2.09	0.52
1:B:53:ALA:HB2	1:B:255:ALA:HB3	1.91	0.52
1:A:48:ALA:HB3	1:A:260[B]:VAL:HG22	1.93	0.51
1:A:57[A]:GLN:NE2	4:A:289:HOH:O	2.40	0.51
1:B:111:LYS:NZ	4:B:730:HOH:O	2.29	0.50
1:B:50:ILE:HD12	1:B:258[B]:VAL:HG11	1.93	0.50
2:A:300[A]:WPP:NAY	2:A:300[A]:WPP:OBI	2.38	0.50
1:B:29[A]:ALA:O	1:B:33[A]:LYS:HG3	2.12	0.50
3:A:301[B]:YPP:HAMB	3:A:301[B]:YPP:CAW	2.42	0.49
1:A:27:THR:HG21	1:A:32:GLN:NE2	2.19	0.49
1:A:137:LYS:HE2	4:A:772:HOH:O	2.14	0.48
1:A:27:THR:CG2	1:A:32:GLN:CG	2.91	0.48
3:A:301[B]:YPP:HAMB	3:A:301[B]:YPP:CAX	2.44	0.48
1:B:39[B]:LYS:HD3	4:B:590:HOH:O	2.14	0.47
2:B:400[A]:WPP:NAY	2:B:400[A]:WPP:OBI	2.47	0.47
1:B:96[B]:GLU:CD	1:B:98:LYS:CE	2.81	0.46
1:B:189[B]:THR:CG2	1:B:190:LEU:N	2.79	0.45
1:A:42[B]:GLY:N	4:A:518:HOH:O	2.36	0.45
1:A:87[A]:GLN:HE22	1:A:93:GLN:HE22	1.64	0.44
1:A:102:LEU:HD11	1:A:113[A]:VAL:HG21	1.99	0.43
1:A:222[B]:ARG:NH1	4:A:364:HOH:O	2.44	0.43
1:B:91:LEU:HB3	1:B:120:ALA:HB2	1.99	0.43
1:A:158:GLU:OE2	4:A:338:HOH:O	2.21	0.43
1:B:29[A]:ALA:HA	4:B:364:HOH:O	2.19	0.43
1:A:251:PRO:HB2	1:A:254:ARG:HB2	2.01	0.42
1:A:27:THR:CG2	1:A:32:GLN:HG3	2.39	0.42
1:A:42[B]:GLY:HA2	4:A:518:HOH:O	2.19	0.41
1:A:91:LEU:HB3	1:A:120:ALA:HB2	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/263 (106%)	272 (98%)	5 (2%)	1 (0%)	34	13
1	B	276/263 (105%)	271 (98%)	4 (1%)	1 (0%)	34	13
All	All	554/526 (105%)	543 (98%)	9 (2%)	2 (0%)	34	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	103	VAL
1	B	103	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/204 (108%)	217 (99%)	3 (1%)	67	42
1	B	219/204 (107%)	217 (99%)	2 (1%)	78	61
All	All	439/408 (108%)	434 (99%)	5 (1%)	71	53

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	THR
1	A	106	ASN
1	A	130	SER
1	B	106	ASN
1	B	130	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	141	GLN
1	A	206	GLN
1	A	252	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	31	GLN
1	B	89	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	WPP	A	300[A]	-	36,39,39	1.60	8 (22%)	50,59,59	1.53	11 (22%)
3	YPP	B	401[B]	-	30,39,39	2.13	5 (16%)	37,57,57	1.28	4 (10%)
2	WPP	B	400[A]	-	36,39,39	1.54	10 (27%)	50,59,59	1.61	13 (26%)
3	YPP	A	301[B]	-	30,39,39	2.15	5 (16%)	37,57,57	1.30	4 (10%)
3	YPP	B	402	-	30,39,39	2.02	5 (16%)	37,57,57	1.20	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	WPP	A	300[A]	-	-	0/22/74/74	0/4/4/4
3	YPP	B	401[B]	-	-	0/22/66/66	0/3/3/3
2	WPP	B	400[A]	-	-	0/22/74/74	0/4/4/4
3	YPP	A	301[B]	-	-	0/22/66/66	0/3/3/3
3	YPP	B	402	-	-	0/22/66/66	0/3/3/3

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301[B]	YPP	CBH-CBF	-8.32	1.40	1.53
3	B	401[B]	YPP	CBH-CBF	-8.31	1.40	1.53
3	B	402	YPP	CBH-CBF	-8.09	1.40	1.53
3	B	401[B]	YPP	CBF-NBE	4.36	1.40	1.35
3	A	301[B]	YPP	CBF-NBE	3.99	1.39	1.35
3	A	301[B]	YPP	CAG-NAE	3.93	1.51	1.46
3	A	301[B]	YPP	CAZ-NBB	-3.93	1.34	1.42
3	B	401[B]	YPP	CAG-NAE	3.85	1.51	1.46
2	A	300[A]	WPP	CAP-CAN	-3.83	1.47	1.53
3	A	301[B]	YPP	CAH-SAI	-3.69	1.77	1.85
2	B	400[A]	WPP	CAP-CAN	-3.65	1.47	1.53
3	B	402	YPP	CBF-NBE	3.63	1.39	1.35
2	A	300[A]	WPP	CAF-SBJ	-3.62	1.77	1.85
2	A	300[A]	WPP	CAG-NAZ	-3.60	1.34	1.42
3	B	401[B]	YPP	CAZ-NBB	-3.60	1.34	1.42
3	B	401[B]	YPP	CAH-SAI	-3.58	1.78	1.85
2	A	300[A]	WPP	CB-SBJ	-3.56	1.75	1.81
3	B	402	YPP	CAG-NAE	3.35	1.51	1.46
3	B	402	YPP	CAZ-NBB	-3.34	1.35	1.42
2	B	400[A]	WPP	CAF-SBJ	-3.12	1.79	1.85
2	B	400[A]	WPP	CA-C	-3.09	1.46	1.54
2	B	400[A]	WPP	CAG-NAZ	-2.85	1.36	1.42
2	B	400[A]	WPP	C-NBB	-2.82	1.31	1.37
2	A	300[A]	WPP	CA-C	-2.79	1.47	1.54
2	B	400[A]	WPP	CB-SBJ	-2.78	1.76	1.81
2	A	300[A]	WPP	C-NBB	-2.73	1.31	1.37
2	B	400[A]	WPP	CAS-CAM	-2.46	1.48	1.52
2	A	300[A]	WPP	CAN-NAZ	-2.44	1.35	1.40
3	B	402	YPP	CAH-SAI	-2.44	1.80	1.85
2	B	400[A]	WPP	CAE-NBB	-2.36	1.45	1.47
2	B	400[A]	WPP	CAN-NAZ	-2.22	1.36	1.40
2	B	400[A]	WPP	CAM-CAK	-2.07	1.49	1.54
2	A	300[A]	WPP	CAP-NBA	-2.01	1.32	1.35

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401[B]	YPP	NAY-CAZ-NBB	4.57	120.69	114.04
2	B	400[A]	WPP	NAY-CAG-NAZ	4.55	120.66	114.04
3	A	301[B]	YPP	NAY-CAZ-NBB	4.52	120.61	114.04
2	A	300[A]	WPP	NAY-CAG-NAZ	4.40	120.44	114.04
2	B	400[A]	WPP	C-CA-N	-4.09	103.73	115.38
3	B	402	YPP	NAY-CAZ-NBB	3.96	119.81	114.04
3	B	402	YPP	OBG-CBF-NBE	-3.90	119.60	123.66
3	A	301[B]	YPP	OBG-CBF-NBE	-3.76	119.75	123.66
2	A	300[A]	WPP	C-CA-N	-3.69	104.86	115.38
3	B	401[B]	YPP	OBG-CBF-NBE	-3.57	119.95	123.66
2	B	400[A]	WPP	CAF-CAE-NBB	3.46	111.71	106.49
2	B	400[A]	WPP	O-C-CA	-2.96	128.03	136.31
2	B	400[A]	WPP	OBI-CAN-NAZ	2.92	124.94	119.98
2	A	300[A]	WPP	CAS-CAM-CAK	2.87	114.60	108.16
2	A	300[A]	WPP	CAF-CAE-NBB	2.86	110.81	106.49
2	A	300[A]	WPP	CA-C-NBB	2.84	95.74	91.87
2	B	400[A]	WPP	CA-CB-NBB	-2.74	84.45	87.98
2	A	300[A]	WPP	O-C-CA	-2.72	128.70	136.31
2	B	400[A]	WPP	CAS-CAM-CAK	2.67	114.17	108.16
2	B	400[A]	WPP	CB-CA-N	-2.52	112.77	118.27
2	B	400[A]	WPP	CAQ-NBA-CAP	2.50	122.64	119.64
2	A	300[A]	WPP	CAJ-CAR-NBA	-2.49	105.11	110.44
2	B	400[A]	WPP	CB-NBB-CAE	-2.47	114.19	117.26
3	A	301[B]	YPP	CAH-SAI-CB	2.42	99.17	93.99
2	A	300[A]	WPP	OBI-CAN-NAZ	2.41	124.06	119.98
2	B	400[A]	WPP	CA-C-NBB	2.28	94.97	91.87
2	A	300[A]	WPP	CB-NBB-CAE	-2.28	114.42	117.26
2	B	400[A]	WPP	SBJ-CB-NBB	2.26	108.14	105.10
2	A	300[A]	WPP	CB-CA-N	-2.23	113.40	118.27
2	B	400[A]	WPP	CAJ-CAR-NBA	-2.23	105.67	110.44
3	A	301[B]	YPP	OBA-CAZ-NBB	-2.18	117.33	119.78
3	B	401[B]	YPP	CAH-SAI-CB	2.13	98.56	93.99
3	B	401[B]	YPP	OBA-CAZ-NBB	-2.09	117.43	119.78
2	A	300[A]	WPP	CA-CB-NBB	-2.06	85.33	87.98

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300[A]	WPP	1	0
3	B	401[B]	YPP	2	0
2	B	400[A]	WPP	2	0
3	A	301[B]	YPP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




## Ligand YPP B 401 (B)




Bond lengths




Bond angles



Torsions




Rings







## Ligand YPP A 301 (B)




Bond lengths




Bond angles



Torsions



Rings



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/263 (99%)	-0.31	3 (1%) 80 84	3, 5, 16, 28	1 (0%)
1	B	260/263 (98%)	-0.32	4 (1%) 73 78	3, 6, 16, 26	0
All	All	522/526 (99%)	-0.32	7 (1%) 77 81	3, 6, 16, 28	1 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	88	LYS	3.4
1	B	288	LEU	3.1
1	B	88	LYS	2.7
1	A	27	THR	2.6
1	B	287	GLY	2.2
1	B	29[A]	ALA	2.2
1	A	201	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	YPP	B	402	37/37	0.91	0.15	9,13,17,17	37
3	YPP	A	301[B]	37/37	0.94	0.13	2,9,12,14	37
3	YPP	B	401[B]	37/37	0.94	0.11	2,5,9,10	37
2	WPP	A	300[A]	36/36	0.94	0.11	5,14,22,25	36
2	WPP	B	400[A]	36/36	0.95	0.09	5,10,25,25	36

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around YPP A 301 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around YPP B 401 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around WPP A 300 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around WPP B 400 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.