



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 30, 2020 – 12:07 PM BST

PDB ID : 2Q4W  
Title : Ensemble refinement of the protein crystal structure of cytokinin oxidase/dehydrogenase (CKX) from Arabidopsis thaliana At5g21482  
Authors : Levin, E.J.; Kondrashov, D.A.; Wesenberg, G.E.; Phillips Jr., G.N.; Center for Eukaryotic Structural Genomics (CESG)  
Deposited on : 2007-05-31  
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1-A	524	<div> <div>3%</div> <div>90%</div> <div>6%</div> </div>
1	10-A	524	<div> <div>3%</div> <div>90%</div> <div>6%</div> </div>
1	11-A	524	<div> <div>3%</div> <div>90%</div> <div>6%</div> </div>
1	12-A	524	<div> <div>3%</div> <div>90%</div> <div>6%</div> </div>
1	13-A	524	<div> <div>3%</div> <div>89%</div> <div>6%</div> </div>
1	14-A	524	<div> <div>3%</div> <div>88%</div> <div>5% 6%</div> </div>
1	15-A	524	<div> <div>3%</div> <div>88%</div> <div>5% 6%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	16-A	524	<p>3% 86% 7% 6%</p>
1	2-A	524	<p>3% 91% • 6%</p>
1	3-A	524	<p>3% 91% • 6%</p>
1	4-A	524	<p>3% 89% 5% 6%</p>
1	5-A	524	<p>3% 90% • 6%</p>
1	6-A	524	<p>3% 89% • • 6%</p>
1	7-A	524	<p>3% 90% • 6%</p>
1	8-A	524	<p>3% 91% • 6%</p>
1	9-A	524	<p>3% 90% • 6%</p>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 72624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cytokinin dehydrogenase 7.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	1-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	2-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	3-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	4-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	5-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	6-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	7-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	8-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	9-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	10-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	11-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	12-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	13-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	14-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	15-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			
1	16-A	491	Total	C	N	O	S	Se	0	0	0
			3853	2450	667	720	8	8			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	EXPRESSION TAG	UNP Q9FUJ1
A	57	MSE	MET	MODIFIED RESIDUE	UNP Q9FUJ1
A	103	MSE	MET	MODIFIED RESIDUE	UNP Q9FUJ1
A	112	MSE	MET	MODIFIED RESIDUE	UNP Q9FUJ1
A	241	MSE	MET	MODIFIED RESIDUE	UNP Q9FUJ1
A	412	MSE	MET	MODIFIED RESIDUE	UNP Q9FUJ1
A	458	MSE	MET	MODIFIED RESIDUE	UNP Q9FUJ1
A	506	MSE	MET	MODIFIED RESIDUE	UNP Q9FUJ1
A	510	MSE	MET	MODIFIED RESIDUE	UNP Q9FUJ1

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	1-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	2-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	3-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	4-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	5-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	6-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	7-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	8-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	9-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	10-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	11-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	12-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	13-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	14-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	15-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	16-A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	1-A	633	Total	O	0	0
			633	633		
3	2-A	633	Total	O	0	0
			633	633		
3	3-A	633	Total	O	0	0
			633	633		
3	4-A	633	Total	O	0	0
			633	633		
3	5-A	633	Total	O	0	0
			633	633		
3	6-A	633	Total	O	0	0
			633	633		
3	7-A	633	Total	O	0	0
			633	633		
3	8-A	633	Total	O	0	0
			633	633		
3	9-A	633	Total	O	0	0
			633	633		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	10-A	633	Total 633	O 633	0	0
3	11-A	633	Total 633	O 633	0	0
3	12-A	633	Total 633	O 633	0	0
3	13-A	633	Total 633	O 633	0	0
3	14-A	633	Total 633	O 633	0	0
3	15-A	633	Total 633	O 633	0	0
3	16-A	633	Total 633	O 633	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Cytokinin dehydrogenase 7




#### • Molecule 1: Cytokinin dehydrogenase 7




#### • Molecule 1: Cytokinin dehydrogenase 7




#### • Molecule 1: Cytokinin dehydrogenase 7









• Molecule 1: Cytokinin dehydrogenase 7




• Molecule 1: Cytokinin dehydrogenase 7




• Molecule 1: Cytokinin dehydrogenase 7




• Molecule 1: Cytokinin dehydrogenase 7




- Molecule 1: Cytokinin dehydrogenase 7




- Molecule 1: Cytokinin dehydrogenase 7




- Molecule 1: Cytokinin dehydrogenase 7



- Molecule 1: Cytokinin dehydrogenase 7





- Molecule 1: Cytokinin dehydrogenase 7









- Molecule 1: Cytokinin dehydrogenase 7



- Molecule 1: Cytokinin dehydrogenase 7



- Molecule 1: Cytokinin dehydrogenase 7



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.86Å 114.50Å 190.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.57 – 1.70 42.57 – 1.70	Depositor EDS
% Data completeness (in resolution range)	94.9 (42.57-1.70) 95.0 (42.57-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.92 (at 1.70Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.143 , 0.201 0.153 , 0.213	Depositor DCC
$R_{free}$ test set	2669 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.6	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 64.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	72624	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	1-A	0.84	3/3938 (0.1%)	0.86	1/5335 (0.0%)
1	2-A	0.83	3/3938 (0.1%)	0.88	0/5335
1	3-A	0.81	1/3938 (0.0%)	0.86	0/5335
1	4-A	0.82	2/3938 (0.1%)	0.87	3/5335 (0.1%)
1	5-A	0.82	3/3938 (0.1%)	0.85	1/5335 (0.0%)
1	6-A	0.83	2/3938 (0.1%)	0.90	4/5335 (0.1%)
1	7-A	0.82	0/3938	0.86	2/5335 (0.0%)
1	8-A	0.81	0/3938	0.84	0/5335
1	9-A	0.84	3/3938 (0.1%)	0.89	2/5335 (0.0%)
1	10-A	0.83	2/3938 (0.1%)	0.90	5/5335 (0.1%)
1	11-A	0.83	2/3938 (0.1%)	0.90	4/5335 (0.1%)
1	12-A	0.82	2/3938 (0.1%)	0.88	1/5335 (0.0%)
1	13-A	0.97	3/3938 (0.1%)	0.99	2/5335 (0.0%)
1	14-A	0.99	4/3938 (0.1%)	1.04	8/5335 (0.1%)
1	15-A	0.97	4/3938 (0.1%)	1.02	2/5335 (0.0%)
1	16-A	0.99	6/3938 (0.2%)	1.01	3/5335 (0.1%)
All	All	0.87	40/63008 (0.1%)	0.91	38/85360 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	6-A	0	1
1	13-A	0	1
1	15-A	0	2
1	16-A	0	3
All	All	0	7

The worst 5 of 40 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	2-A	309	CYS	CB-SG	-7.64	1.69	1.82
1	16-A	309	CYS	CB-SG	-7.54	1.69	1.82
1	10-A	309	CYS	CB-SG	-7.47	1.69	1.82
1	3-A	309	CYS	CB-SG	-7.33	1.69	1.82
1	16-A	377	GLU	CG-CD	7.32	1.62	1.51

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	14-A	475	ASP	CB-CG-OD1	9.25	126.62	118.30
1	15-A	230	ARG	NE-CZ-NH2	8.24	124.42	120.30
1	6-A	385	LEU	CA-CB-CG	6.61	130.50	115.30
1	4-A	161	THR	N-CA-C	-6.56	93.28	111.00
1	14-A	92	ARG	NE-CZ-NH1	-6.54	117.03	120.30

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	13-A	322	TYR	Sidechain
1	15-A	153	TYR	Sidechain
1	15-A	415	TYR	Sidechain
1	16-A	273	TYR	Sidechain
1	6-A	124	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-A	3853	0	3764	0	0
1	2-A	3853	0	3764	0	0
1	3-A	3853	0	3764	0	0
1	4-A	3853	0	3764	0	0
1	5-A	3853	0	3764	0	0
1	6-A	3853	0	3764	0	0
1	7-A	3853	0	3764	0	0
1	8-A	3853	0	3764	0	0
1	9-A	3853	0	3764	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	10-A	3853	0	3764	0	0
1	11-A	3853	0	3764	0	0
1	12-A	3853	0	3764	0	0
1	13-A	3853	0	3764	0	0
1	14-A	3853	0	3764	0	0
1	15-A	3853	0	3764	0	0
1	16-A	3853	0	3764	0	0
2	1-A	53	0	31	0	0
2	2-A	53	0	29	0	0
2	3-A	53	0	31	0	0
2	4-A	53	0	31	0	0
2	5-A	53	0	31	0	0
2	6-A	53	0	31	0	0
2	7-A	53	0	31	0	0
2	8-A	53	0	31	0	0
2	9-A	53	0	31	0	0
2	10-A	53	0	30	0	0
2	11-A	53	0	31	0	0
2	12-A	53	0	31	0	0
2	13-A	53	0	31	0	0
2	14-A	53	0	29	0	0
2	15-A	53	0	28	0	0
2	16-A	53	0	30	0	0
3	1-A	633	0	0	0	0
3	2-A	633	0	0	0	0
3	3-A	633	0	0	0	0
3	4-A	633	0	0	0	0
3	5-A	633	0	0	0	0
3	6-A	633	0	0	0	0
3	7-A	633	0	0	0	0
3	8-A	633	0	0	0	0
3	9-A	633	0	0	0	0
3	10-A	633	0	0	0	0
3	11-A	633	0	0	0	0
3	12-A	633	0	0	0	0
3	13-A	633	0	0	0	0
3	14-A	633	0	0	0	0
3	15-A	633	0	0	0	0
3	16-A	633	0	0	0	0
All	All	72624	0	60711	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). Clashscore could not be calculated for this entry.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1-A	489/524 (93%)	439 (90%)	44 (9%)	6 (1%)	13	3
1	2-A	489/524 (93%)	458 (94%)	24 (5%)	7 (1%)	11	2
1	3-A	489/524 (93%)	451 (92%)	35 (7%)	3 (1%)	25	11
1	4-A	489/524 (93%)	442 (90%)	37 (8%)	10 (2%)	7	1
1	5-A	489/524 (93%)	454 (93%)	30 (6%)	5 (1%)	15	4
1	6-A	489/524 (93%)	441 (90%)	39 (8%)	9 (2%)	8	1
1	7-A	489/524 (93%)	434 (89%)	42 (9%)	13 (3%)	5	0
1	8-A	489/524 (93%)	447 (91%)	37 (8%)	5 (1%)	15	4
1	9-A	489/524 (93%)	441 (90%)	42 (9%)	6 (1%)	13	3
1	10-A	489/524 (93%)	442 (90%)	35 (7%)	12 (2%)	5	1
1	11-A	489/524 (93%)	444 (91%)	37 (8%)	8 (2%)	9	1
1	12-A	489/524 (93%)	454 (93%)	32 (6%)	3 (1%)	25	11
1	13-A	489/524 (93%)	444 (91%)	39 (8%)	6 (1%)	13	3
1	14-A	489/524 (93%)	433 (88%)	49 (10%)	7 (1%)	11	2
1	15-A	489/524 (93%)	439 (90%)	45 (9%)	5 (1%)	15	4
1	16-A	489/524 (93%)	450 (92%)	35 (7%)	4 (1%)	19	6
All	All	7824/8384 (93%)	7113 (91%)	602 (8%)	109 (1%)	11	2

5 of 109 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1-A	382	TRP
1	1-A	414	VAL

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	2-A	269	SER
1	4-A	188	THR
1	4-A	341	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1-A	415/430 (96%)	406 (98%)	9 (2%)	52	34
1	2-A	415/430 (96%)	411 (99%)	4 (1%)	76	67
1	3-A	415/430 (96%)	407 (98%)	8 (2%)	57	41
1	4-A	415/430 (96%)	402 (97%)	13 (3%)	40	21
1	5-A	415/430 (96%)	405 (98%)	10 (2%)	49	31
1	6-A	415/430 (96%)	403 (97%)	12 (3%)	42	23
1	7-A	415/430 (96%)	407 (98%)	8 (2%)	57	41
1	8-A	415/430 (96%)	407 (98%)	8 (2%)	57	41
1	9-A	415/430 (96%)	403 (97%)	12 (3%)	42	23
1	10-A	415/430 (96%)	408 (98%)	7 (2%)	60	46
1	11-A	415/430 (96%)	407 (98%)	8 (2%)	57	41
1	12-A	415/430 (96%)	402 (97%)	13 (3%)	40	21
1	13-A	415/430 (96%)	401 (97%)	14 (3%)	37	18
1	14-A	415/430 (96%)	398 (96%)	17 (4%)	30	12
1	15-A	415/430 (96%)	398 (96%)	17 (4%)	30	12
1	16-A	415/430 (96%)	390 (94%)	25 (6%)	19	6
All	All	6640/6880 (96%)	6455 (97%)	185 (3%)	43	25

5 of 185 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	9-A	522	ARG
1	12-A	166	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	16-A	206	CYS
1	10-A	322	TYR
1	11-A	68	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 61 such sidechains are listed below:

Mol	Chain	Res	Type
1	8-A	119	HIS
1	9-A	211	ASN
1	15-A	467	HIS
1	8-A	211	ASN
1	8-A	345	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	10-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	11-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	2-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	13-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	4-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	5-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	16-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	1-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	12-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	7-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	8-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	9-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	3-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	14-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	15-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)
2	FAD	6-A	701	-	51,58,58	2.40	16 (31%)	60,89,89	2.48	17 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	10-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	11-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	2-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	13-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	4-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	5-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	16-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	1-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	12-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	7-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	8-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	9-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	3-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	14-A	701	-	-	5/30/50/50	0/6/6/6

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	15-A	701	-	-	5/30/50/50	0/6/6/6
2	FAD	6-A	701	-	-	5/30/50/50	0/6/6/6

The worst 5 of 256 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	10-A	701	FAD	O5'-C5'	-6.39	1.20	1.44
2	11-A	701	FAD	O5'-C5'	-6.39	1.20	1.44
2	2-A	701	FAD	O5'-C5'	-6.39	1.20	1.44
2	13-A	701	FAD	O5'-C5'	-6.39	1.20	1.44
2	4-A	701	FAD	O5'-C5'	-6.39	1.20	1.44

The worst 5 of 272 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	10-A	701	FAD	C4-N3-C2	11.03	124.45	115.14
2	11-A	701	FAD	C4-N3-C2	11.03	124.45	115.14
2	2-A	701	FAD	C4-N3-C2	11.03	124.45	115.14
2	13-A	701	FAD	C4-N3-C2	11.03	124.45	115.14
2	4-A	701	FAD	C4-N3-C2	11.03	124.45	115.14

There are no chirality outliers.

5 of 80 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	10-A	701	FAD	O4'-C4'-C5'-O5'
2	11-A	701	FAD	O4'-C4'-C5'-O5'
2	2-A	701	FAD	O4'-C4'-C5'-O5'
2	13-A	701	FAD	O4'-C4'-C5'-O5'
2	4-A	701	FAD	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	1-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	2-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	3-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	4-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	5-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	6-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	7-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	8-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	9-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	10-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	11-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	12-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	13-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	14-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	15-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
1	16-A	483/524 (92%)	-0.15	14 (2%)	51	56	8, 18, 37, 55	483 (100%)
All	All	7728/8384 (92%)	-0.15	224 (2%)	49	56	8, 18, 37, 55	7728 (100%)

The worst 5 of 224 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	1-A	449	CYS	5.9
1	2-A	449	CYS	5.9
1	3-A	449	CYS	5.9
1	4-A	449	CYS	5.9
1	5-A	449	CYS	5.9

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ


There are no monosaccharides in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FAD	10-A	701	53/53	0.97	0.11	0,9,14,17	53
2	FAD	11-A	701	53/53	0.97	0.11	0,10,14,15	53
2	FAD	2-A	701	53/53	0.97	0.11	0,9,14,18	53
2	FAD	13-A	701	53/53	0.97	0.11	0,9,14,16	53
2	FAD	4-A	701	53/53	0.97	0.11	0,9,14,15	53
2	FAD	5-A	701	53/53	0.97	0.11	6,10,14,15	53
2	FAD	16-A	701	53/53	0.97	0.11	0,8,15,17	53
2	FAD	1-A	701	53/53	0.97	0.11	0,10,14,14	53
2	FAD	12-A	701	53/53	0.97	0.11	0,9,14,15	53
2	FAD	7-A	701	53/53	0.97	0.11	0,10,14,14	53
2	FAD	8-A	701	53/53	0.97	0.11	0,10,14,17	53
2	FAD	9-A	701	53/53	0.97	0.11	3,10,14,15	53
2	FAD	3-A	701	53/53	0.97	0.11	0,9,14,19	53
2	FAD	14-A	701	53/53	0.97	0.11	0,8,14,16	53
2	FAD	15-A	701	53/53	0.97	0.11	0,8,15,17	53
2	FAD	6-A	701	53/53	0.97	0.11	0,10,16,17	53

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.