



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 12:18 am BST

PDB ID : 3Q6I
Title : Crystal structure of FabG4 and coenzyme binary complex
Authors : Dutta, D.; Bhattacharyya, S.; Das, A.K.
Deposited on : 2011-01-01
Resolution : 2.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 76%, yellow 76%, yellow 93%, grey 93%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> 76% 17% </div> </div>
1	B	446	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 2%, green 2%, green 77%, yellow 77%, yellow 95%, grey 95%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> 77% 18% 5% </div> </div>
1	C	446	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 3%, green 3%, green 74%, yellow 74%, yellow 91%, grey 91%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> 74% 17% 6% </div> </div>
1	D	446	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 7%, green 7%, green 69%, yellow 69%, yellow 90%, grey 90%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> 69% 21% 8% </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12462 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-oxoacyl-(Acyl-carrier-protein) reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	427	Total	C	N	O	S	0	0	0
			3080	1925	553	595	7			
1	B	425	Total	C	N	O	S	0	0	0
			3059	1916	543	593	7			
1	C	418	Total	C	N	O	S	0	0	0
			2991	1871	530	584	6			
1	D	409	Total	C	N	O	S	0	0	0
			2927	1836	514	571	6			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	HIS	-	EXPRESSION TAG	UNP O53665
A	10	HIS	-	EXPRESSION TAG	UNP O53665
A	11	HIS	-	EXPRESSION TAG	UNP O53665
A	12	HIS	-	EXPRESSION TAG	UNP O53665
A	13	HIS	-	EXPRESSION TAG	UNP O53665
A	14	HIS	-	EXPRESSION TAG	UNP O53665
A	15	GLY	-	EXPRESSION TAG	UNP O53665
A	16	SER	-	EXPRESSION TAG	UNP O53665
B	9	HIS	-	EXPRESSION TAG	UNP O53665
B	10	HIS	-	EXPRESSION TAG	UNP O53665
B	11	HIS	-	EXPRESSION TAG	UNP O53665
B	12	HIS	-	EXPRESSION TAG	UNP O53665
B	13	HIS	-	EXPRESSION TAG	UNP O53665
B	14	HIS	-	EXPRESSION TAG	UNP O53665
B	15	GLY	-	EXPRESSION TAG	UNP O53665
B	16	SER	-	EXPRESSION TAG	UNP O53665
C	9	HIS	-	EXPRESSION TAG	UNP O53665
C	10	HIS	-	EXPRESSION TAG	UNP O53665
C	11	HIS	-	EXPRESSION TAG	UNP O53665
C	12	HIS	-	EXPRESSION TAG	UNP O53665
C	13	HIS	-	EXPRESSION TAG	UNP O53665

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	14	HIS	-	EXPRESSION TAG	UNP O53665
C	15	GLY	-	EXPRESSION TAG	UNP O53665
C	16	SER	-	EXPRESSION TAG	UNP O53665
D	9	HIS	-	EXPRESSION TAG	UNP O53665
D	10	HIS	-	EXPRESSION TAG	UNP O53665
D	11	HIS	-	EXPRESSION TAG	UNP O53665
D	12	HIS	-	EXPRESSION TAG	UNP O53665
D	13	HIS	-	EXPRESSION TAG	UNP O53665
D	14	HIS	-	EXPRESSION TAG	UNP O53665
D	15	GLY	-	EXPRESSION TAG	UNP O53665
D	16	SER	-	EXPRESSION TAG	UNP O53665

- # NAD
-
- The image displays the chemical structure of Nicotinamide Adenine Dinucleotide (NAD). It consists of two nucleotides linked by a pyrophosphate bridge. The first nucleotide is composed of a nicotinamide ring (labeled with N1A, C6A, N7A, C5A, C2A, N3A, C4A, N4A) and a ribose sugar (labeled with C5B, C4B, C3B, C2B, C1B, O2B, O3B, O4B, O5B). The second nucleotide is composed of an adenine ring (labeled with N6A, C6A, N7A, C5A, C2A, N3A, C4A, N4A) and a ribose sugar (labeled with C5B, C4B, C3B, C2B, C1B, O2B, O3B, O4B, O5B). The pyrophosphate bridge connects the 5' carbon of the first ribose to the 5' carbon of the second ribose. The structure is shown in a 3D representation with various atoms and bonds labeled.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
2	B	1	Total 44	C 21	N 7	O 14	P 2	0	0
2	C	1	Total 44	C 21	N 7	O 14	P 2	0	0
2	D	1	Total 44	C 21	N 7	O 14	P 2	0	0

- 
- WORLD WIDE
PDB
PROTEIN DATA BANK




- Molecule 4 is water.





• Molecule 1: 3-oxoacyl-(Acyl-carrier-protein) reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	63.07Å 71.03Å 92.92Å 105.02° 97.06° 93.66°	Depositor
Resolution (Å)	19.78 – 2.59 19.78 – 2.59	Depositor EDS
% Data completeness (in resolution range)	95.5 (19.78-2.59) 95.7 (19.78-2.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.59Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.168 , 0.242 0.194 , 0.255	Depositor DCC
R_{free} test set	2322 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.129	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12462	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, ZPG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/3125	0.83	1/4248 (0.0%)
1	B	0.70	0/3102	0.77	1/4212 (0.0%)
1	C	0.65	0/3036	0.76	0/4132
1	D	0.61	0/2968	0.73	0/4032
All	All	0.67	0/12231	0.77	2/16624 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	345	LEU	CA-CB-CG	-6.02	101.45	115.30
1	B	345	LEU	CA-CB-CG	5.00	126.80	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3080	0	3100	63	0
1	B	3059	0	3073	55	0
1	C	2991	0	2963	51	0
1	D	2927	0	2916	56	0
2	A	44	0	26	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	44	0	26	3	0
2	C	44	0	26	1	0
2	D	44	0	26	1	0
3	A	13	0	19	0	0
4	A	78	0	0	1	0
4	B	69	0	0	4	0
4	C	39	0	0	1	0
4	D	30	0	0	0	0
All	All	12462	0	12175	212	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 212 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:ARG:HH11	1:A:38:ARG:HG2	0.96	1.10
1:D:337:GLY:HA2	1:D:338:GLU:HB2	1.51	0.92
1:A:38:ARG:CG	1:A:38:ARG:HH11	1.82	0.92
1:B:299:THR:O	1:B:300:ARG:HB2	1.72	0.89
1:C:173:GLY:HA2	1:C:201:SER:OG	1.72	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/446 (95%)	398 (94%)	22 (5%)	3 (1%)	22	43
1	B	417/446 (94%)	393 (94%)	21 (5%)	3 (1%)	22	43
1	C	412/446 (92%)	384 (93%)	22 (5%)	6 (2%)	10	21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	399/446 (90%)	364 (91%)	30 (8%)	5 (1%)	12	24
All	All	1651/1784 (92%)	1539 (93%)	95 (6%)	17 (1%)	15	32

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	250	GLU
1	B	300	ARG
1	C	347	SER
1	A	397	MET
1	C	39	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/323 (95%)	287 (94%)	19 (6%)	18	37
1	B	303/323 (94%)	289 (95%)	14 (5%)	27	51
1	C	292/323 (90%)	279 (96%)	13 (4%)	27	52
1	D	287/323 (89%)	267 (93%)	20 (7%)	15	30
All	All	1188/1292 (92%)	1122 (94%)	66 (6%)	21	42

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	327	THR
1	C	195	SER
1	D	394	GLU
1	B	348	ILE
1	C	69	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	239	HIS
1	B	284	HIS
1	C	284	HIS
1	B	112	ASN
1	C	420	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAD	B	3	-	42,48,48	1.87	4 (9%)	50,73,73	1.39	3 (6%)
2	NAD	C	2	-	42,48,48	1.90	4 (9%)	50,73,73	1.35	3 (6%)
2	NAD	A	1	-	42,48,48	1.87	6 (14%)	50,73,73	1.52	5 (10%)
2	NAD	D	4	-	42,48,48	1.85	4 (9%)	50,73,73	1.37	5 (10%)
3	ZPG	A	455	-	9,12,28	0.82	0	10,14,34	1.37	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	B	3	-	-	10/26/62/62	0/5/5/5
2	NAD	C	2	-	-	7/26/62/62	0/5/5/5
2	NAD	A	1	-	-	2/26/62/62	0/5/5/5
2	NAD	D	4	-	-	5/26/62/62	0/5/5/5
3	ZPG	A	455	-	-	2/12/12/31	-

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	NAD	O7N-C7N	9.85	1.43	1.24
2	C	2	NAD	O7N-C7N	9.65	1.42	1.24
2	D	4	NAD	O7N-C7N	9.45	1.42	1.24
2	A	1	NAD	O7N-C7N	9.20	1.41	1.24
2	D	4	NAD	C2A-N3A	4.16	1.38	1.32

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	NAD	N3A-C2A-N1A	-6.97	117.78	128.68
2	B	3	NAD	N3A-C2A-N1A	-6.14	119.08	128.68
2	C	2	NAD	N3A-C2A-N1A	-6.00	119.30	128.68
2	D	4	NAD	N3A-C2A-N1A	-5.10	120.70	128.68
2	D	4	NAD	C6N-N1N-C2N	-4.09	118.24	121.97

There are no chirality outliers.

5 of 26 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	3	NAD	C5B-O5B-PA-O1A
2	B	3	NAD	C5B-O5B-PA-O2A
2	B	3	NAD	C5D-O5D-PN-O3
2	C	2	NAD	C5D-O5D-PN-O3
2	C	2	NAD	C5D-O5D-PN-O2N


There are no ring outliers.


4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	NAD	3	0
2	C	2	NAD	1	0
2	A	1	NAD	4	0
2	D	4	NAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	427/446 (95%)	-0.16	4 (0%) 84 82	17, 33, 54, 67	0
1	B	425/446 (95%)	-0.01	11 (2%) 56 50	14, 36, 71, 83	0
1	C	418/446 (93%)	-0.01	15 (3%) 42 35	19, 40, 66, 78	0
1	D	409/446 (91%)	0.41	32 (7%) 13 9	27, 52, 88, 103	0
All	All	1679/1784 (94%)	0.06	62 (3%) 41 34	14, 40, 75, 103	0

The worst 5 of 62 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	400	ALA	6.1
1	D	72	GLY	5.2
1	D	239	HIS	4.3
1	D	81	ASP	4.2
1	D	272	ASP	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ZPG	A	455	13/29	0.85	0.17	60,63,66,67	0
2	NAD	D	4	44/44	0.93	0.15	54,66,71,74	0
2	NAD	B	3	44/44	0.93	0.17	55,62,70,71	0
2	NAD	C	2	44/44	0.95	0.13	39,46,53,55	0
2	NAD	A	1	44/44	0.95	0.13	30,36,46,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NAD D 4:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD B 3:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD C 2:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 1:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.