



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 6, 2020 – 11:56 pm BST

PDB ID : 3Q6M  
Title : Crystal Structure of Human MC-HSP90 in C2221 Space Group  
Authors : Lee, C.C.; Lin, T.W.; Ko, T.P.; Wang, A.H.-J.  
Deposited on : 2011-01-03  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	<div> <div>2%</div> <div> <div></div> <div>59%</div> <div>19%</div> <div>5%</div> <div>17%</div> </div> </div>
1	B	448	<div> <div>2%</div> <div> <div></div> <div>63%</div> <div>16%</div> <div>5%</div> <div>17%</div> </div> </div>
1	C	448	<div> <div>2%</div> <div> <div></div> <div>59%</div> <div>19%</div> <div>•</div> <div>18%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9345 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heat shock protein HSP 90-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	372	Total	C	N	O	S	0	0	0
			3069	1954	518	582	15			
1	B	374	Total	C	N	O	S	0	0	0
			3085	1964	520	586	15			
1	C	369	Total	C	N	O	S	0	0	0
			3043	1939	512	577	15			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	291	ALA	-	EXPRESSION TAG	UNP P07900
A	292	ALA	-	EXPRESSION TAG	UNP P07900
A	733	HIS	-	EXPRESSION TAG	UNP P07900
A	734	HIS	-	EXPRESSION TAG	UNP P07900
A	735	HIS	-	EXPRESSION TAG	UNP P07900
A	736	HIS	-	EXPRESSION TAG	UNP P07900
A	737	HIS	-	EXPRESSION TAG	UNP P07900
A	738	HIS	-	EXPRESSION TAG	UNP P07900
B	291	ALA	-	EXPRESSION TAG	UNP P07900
B	292	ALA	-	EXPRESSION TAG	UNP P07900
B	733	HIS	-	EXPRESSION TAG	UNP P07900
B	734	HIS	-	EXPRESSION TAG	UNP P07900
B	735	HIS	-	EXPRESSION TAG	UNP P07900
B	736	HIS	-	EXPRESSION TAG	UNP P07900
B	737	HIS	-	EXPRESSION TAG	UNP P07900
B	738	HIS	-	EXPRESSION TAG	UNP P07900
C	291	ALA	-	EXPRESSION TAG	UNP P07900
C	292	ALA	-	EXPRESSION TAG	UNP P07900
C	733	HIS	-	EXPRESSION TAG	UNP P07900
C	734	HIS	-	EXPRESSION TAG	UNP P07900
C	735	HIS	-	EXPRESSION TAG	UNP P07900
C	736	HIS	-	EXPRESSION TAG	UNP P07900
C	737	HIS	-	EXPRESSION TAG	UNP P07900

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	738	HIS	-	EXPRESSION TAG	UNP P07900

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	44	Total	O	0	0
			44	44		
3	B	53	Total	O	0	0
			53	53		
3	C	36	Total	O	0	0
			36	36		





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	162.70Å 304.55Å 87.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.00 29.84 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (30.00-3.00) 95.0 (29.84-3.00)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.49 (at 3.00Å)	Xtriage
Refinement program	CNS 1.2, REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.209 , 0.252 0.210 , 0.208	Depositor DCC
$R_{free}$ test set	2184 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	88.1	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 76.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9345	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	104.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	0/3120	0.69	0/4188
1	B	0.66	0/3136	0.71	1/4210 (0.0%)
1	C	0.57	0/3094	0.65	1/4154 (0.0%)
All	All	0.61	0/9350	0.68	2/12552 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	646	LEU	CA-CB-CG	7.56	132.69	115.30
1	C	696	LEU	CA-CB-CG	5.52	128.00	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3069	0	3108	72	0
1	B	3085	0	3123	70	0
1	C	3043	0	3081	64	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	44	0	0	1	0
3	B	53	0	0	0	0
3	C	36	0	0	1	0
All	All	9345	0	9312	197	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 197 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:367:ARG:HG3	1:B:367:ARG:HH11	1.15	1.03
1:C:607:THR:HG22	1:C:610:MET:HG3	1.43	0.99
1:B:410:LYS:HE3	1:B:414:LYS:HE3	1.44	0.97
1:C:525:ILE:H	1:C:525:ILE:HD12	1.32	0.94
1:A:656:ASP:HB3	1:A:659:VAL:HG23	1.48	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/448 (81%)	344 (94%)	15 (4%)	5 (1%)	11	43
1	B	366/448 (82%)	335 (92%)	26 (7%)	5 (1%)	11	43
1	C	361/448 (81%)	323 (90%)	27 (8%)	11 (3%)	4	24
All	All	1091/1344 (81%)	1002 (92%)	68 (6%)	21 (2%)	8	36

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	580	GLU
1	B	376	GLU
1	C	432	GLU
1	A	429	GLU
1	A	696	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/412 (84%)	305 (88%)	41 (12%)	5	22
1	B	348/412 (84%)	309 (89%)	39 (11%)	6	24
1	C	343/412 (83%)	305 (89%)	38 (11%)	6	25
All	All	1037/1236 (84%)	919 (89%)	118 (11%)	5	24

5 of 118 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	435	LYS
1	B	578	ILE
1	C	614	MET
1	B	443	LYS
1	B	535	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	487	ASN
1	B	501	GLN
1	C	487	ASN
1	B	450	HIS
1	B	454	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	1	-	4,4,4	0.14	0	6,6,6	0.26	0
2	SO4	B	2	-	4,4,4	0.13	0	6,6,6	0.46	0
2	SO4	C	3	-	4,4,4	0.12	0	6,6,6	0.19	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	372/448 (83%)	-0.14	7 (1%) 66 37	67, 94, 136, 185	0
1	B	374/448 (83%)	-0.20	7 (1%) 66 37	61, 89, 143, 174	0
1	C	369/448 (82%)	-0.01	10 (2%) 54 26	69, 114, 187, 222	0
All	All	1115/1344 (82%)	-0.12	24 (2%) 62 33	61, 98, 161, 222	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	630	ALA	5.5
1	A	395	PRO	4.3
1	C	606	TRP	4.3
1	A	559	LYS	3.6
1	C	604	TYR	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	C	3	5/5	0.69	0.35	164,165,165,165	0
2	SO4	B	2	5/5	0.88	0.19	120,121,122,123	0
2	SO4	A	1	5/5	0.91	0.16	127,128,128,129	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.