



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 10:10 pm BST

PDB ID : 6Q71
Title : Crystal structure of the alanine racemase Bsu17640 from *Bacillus subtilis* in the presence of Bis-Tris propane
Authors : Bernardo-Garcia, N.; Gago, F.; Hermoso, J.A.
Deposited on : 2018-12-12
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	<div> <div>6%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>..</div> </div> </div>
1	B	394	<div> <div>7%</div> <div> <div></div> <div>87%</div> <div>10%</div> <div>..</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6630 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alanine racemase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	0	0
			3009	1924	527	544	14			
1	B	386	Total	C	N	O	S	0	0	0
			3009	1924	527	544	14			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Mg	0	0
			3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	291	Total	O	0	0
			291	291		
5	B	286	Total	O	0	0
			286	286		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alanine racemase 2



• Molecule 1: Alanine racemase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	73.24Å 73.24Å 333.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.32 – 1.92 49.32 – 1.92	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.32-1.92) 100.0 (49.32-1.92)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 1.92Å)	Xtriage
Refinement program	PHENIX (1.10 _2155: ???)	Depositor
R, R_{free}	0.204 , 0.232 0.212 , 0.239	Depositor DCC
R_{free} test set	3642 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6630	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.82	10/3063 (0.3%)	0.78	5/4135 (0.1%)
1	B	0.81	6/3063 (0.2%)	0.84	7/4135 (0.2%)
All	All	0.82	16/6126 (0.3%)	0.81	12/8270 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	105	ALA	C-N	-13.12	1.03	1.34
1	B	38	VAL	C-N	13.07	1.64	1.34
1	B	44	GLY	C-N	-10.78	1.09	1.34
1	B	105	ALA	C-N	-9.52	1.12	1.34
1	A	372	ARG	CZ-NH2	7.71	1.43	1.33
1	A	106	PHE	C-N	-7.54	1.16	1.34
1	B	106	PHE	C-N	-7.36	1.17	1.34
1	A	317	ARG	CZ-NH1	6.86	1.42	1.33
1	A	162	ARG	CZ-NH1	6.86	1.42	1.33
1	A	162	ARG	CZ-NH2	6.85	1.42	1.33
1	B	372	ARG	CZ-NH1	6.40	1.41	1.33
1	A	268	ARG	CZ-NH1	6.12	1.41	1.33
1	A	317	ARG	CZ-NH2	5.98	1.40	1.33
1	B	372	ARG	CZ-NH2	5.87	1.40	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	268	ARG	CZ-NH2	5.78	1.40	1.33
1	A	372	ARG	CZ-NH1	5.39	1.40	1.33

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	44	GLY	C-N-CA	13.82	156.25	121.70
1	B	44	GLY	O-C-N	-12.67	102.43	122.70
1	A	106	PHE	O-C-N	-10.36	106.12	122.70
1	B	44	GLY	CA-C-N	9.81	138.78	117.20
1	B	38	VAL	C-N-CA	-7.58	102.74	121.70
1	A	106	PHE	CA-C-N	7.34	133.36	117.20
1	B	106	PHE	O-C-N	-5.87	113.31	122.70
1	B	372	ARG	NE-CZ-NH1	-5.78	117.41	120.30
1	B	353	ASP	CB-CG-OD1	5.70	123.43	118.30
1	A	317	ARG	NE-CZ-NH1	-5.25	117.68	120.30
1	A	373	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	A	268	ARG	NE-CZ-NH1	-5.17	117.71	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	383	GLU	Mainchain
1	B	45	HIS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3009	0	3109	37	0
1	B	3009	0	3108	34	0
2	A	15	0	6	3	0
2	B	15	0	6	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	B	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	291	0	0	9	2
5	B	286	0	0	16	1
All	All	6630	0	6229	70	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:ILE:HD11	5:A:688:HOH:O	1.39	1.20
1:A:283:VAL:HG21	1:A:330:ASN:O	1.43	1.16
1:B:180:LEU:CD2	5:B:711:HOH:O	1.99	1.09
1:A:283:VAL:CG2	1:A:330:ASN:O	2.17	0.93
1:A:327:LEU:O	1:A:331:GLY:CA	2.15	0.93
1:B:142:VAL:HG23	1:B:147:GLU:HG2	1.52	0.91
1:A:1:MET:N	5:A:503:HOH:O	2.02	0.91
1:B:107:GLN:CG	5:B:524:HOH:O	2.22	0.86
1:B:180:LEU:HD22	5:B:711:HOH:O	1.67	0.85
1:A:327:LEU:O	1:A:331:GLY:HA3	1.77	0.84
1:B:180:LEU:HD23	5:B:711:HOH:O	1.65	0.83
1:A:372:ARG:NH2	5:A:502:HOH:O	1.78	0.83
1:B:107:GLN:HG2	5:B:524:HOH:O	1.79	0.82
1:A:169:HIS:CG	5:A:508:HOH:O	2.32	0.81
1:A:169:HIS:ND1	2:A:401:PLP:H2A2	1.99	0.76
1:B:346:LYS:O	5:B:502:HOH:O	2.02	0.76
1:A:327:LEU:O	1:A:331:GLY:HA2	1.86	0.75
1:B:301:ASN:ND2	5:B:505:HOH:O	2.19	0.73
1:B:33:LYS:NZ	5:B:503:HOH:O	2.08	0.73
1:A:169:HIS:CE1	2:A:401:PLP:C2	2.73	0.72
1:B:182:LYS:O	1:B:186:GLU:HG3	1.87	0.72
1:B:301:ASN:O	5:B:504:HOH:O	2.08	0.71
1:A:327:LEU:CB	1:A:331:GLY:HA3	2.22	0.70
1:A:354:GLU:OE1	5:A:504:HOH:O	2.10	0.69
1:B:142:VAL:HG23	1:B:147:GLU:CG	2.20	0.69
1:A:282:GLU:HA	1:A:329:GLU:HA	1.74	0.68
1:B:146:GLU:HB2	5:B:540:HOH:O	1.93	0.67
1:A:1:MET:HE3	1:A:2:ILE:HD12	1.77	0.66
1:A:169:HIS:ND1	5:A:508:HOH:O	2.27	0.66
1:B:169:HIS:CG	5:B:590:HOH:O	2.47	0.66
1:A:384:ILE:CD1	5:A:688:HOH:O	2.15	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:264:ARG:CZ	5:B:514:HOH:O	2.46	0.61
1:A:16:ALA:CB	1:A:248:GLU:HG2	2.32	0.59
1:A:169:HIS:CE1	2:A:401:PLP:N1	2.72	0.58
1:A:384:ILE:HD12	1:A:384:ILE:N	2.18	0.57
5:A:636:HOH:O	1:B:143:ARG:HD3	2.05	0.57
1:A:300:SER:HB3	1:A:317:ARG:HA	1.87	0.56
1:B:271:SER:OG	1:B:272:TYR:N	2.39	0.56
1:A:327:LEU:C	1:A:331:GLY:HA3	2.26	0.56
1:B:169:HIS:HB2	5:B:590:HOH:O	2.06	0.55
1:A:16:ALA:HB1	1:A:248:GLU:HG2	1.89	0.54
1:A:383:GLU:C	1:A:384:ILE:HD12	2.27	0.54
1:A:327:LEU:HB3	1:A:331:GLY:HA3	1.88	0.54
1:B:376:ARG:O	1:B:386:LYS:HD3	2.07	0.54
1:A:384:ILE:CD1	1:A:384:ILE:N	2.71	0.53
1:B:264:ARG:NE	5:B:514:HOH:O	2.42	0.53
1:A:282:GLU:HA	1:A:329:GLU:CA	2.40	0.52
1:A:385:PHE:HD1	1:A:386:LYS:HB2	1.75	0.51
1:B:142:VAL:CG2	1:B:147:GLU:CG	2.87	0.51
1:B:92:CYS:HA	5:B:509:HOH:O	2.12	0.49
1:B:42:GLY:O	1:B:45:HIS:HB2	2.13	0.48
1:A:1:MET:CE	1:A:2:ILE:HD12	2.42	0.48
1:B:22:ARG:HG2	1:B:26:ARG:HD3	1.95	0.48
1:A:327:LEU:HB2	1:A:331:GLY:HA3	1.96	0.47
1:B:169:HIS:CE1	2:B:401:PLP:N1	2.83	0.47
1:A:327:LEU:HB3	1:A:331:GLY:CA	2.45	0.46
1:B:293:ASP:HB3	1:B:374:ILE:HD11	1.98	0.46
1:A:353:ASP:CG	5:A:523:HOH:O	2.54	0.44
1:B:291:TYR:HA	1:B:295:TYR:O	2.18	0.44
1:A:331:GLY:O	1:A:332:GLU:C	2.56	0.44
1:B:169:HIS:CG	2:B:401:PLP:H2A2	2.52	0.43
1:B:351:SER:HB2	5:B:633:HOH:O	2.19	0.42
1:A:22:ARG:HG2	1:A:26:ARG:HD3	2.00	0.42
1:B:384:ILE:CG2	1:B:385:PHE:N	2.83	0.42
1:B:45:HIS:NE2	1:B:251:LEU:HB2	2.34	0.42
1:A:266:GLU:O	1:B:187:LYS:NZ	2.52	0.42
1:A:228:ILE:HG13	1:A:233:LEU:HB2	2.02	0.41
1:A:379:ILE:HG12	1:A:384:ILE:HG13	2.03	0.41
1:B:228:ILE:HG13	1:B:233:LEU:HB2	2.03	0.40
1:B:169:HIS:ND1	2:B:401:PLP:H2A2	2.36	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:727:HOH:O	5:A:727:HOH:O[5_657]	1.70	0.50
5:A:768:HOH:O	5:B:774:HOH:O[1_565]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/394 (98%)	372 (97%)	12 (3%)	0	100	100
1	B	384/394 (98%)	371 (97%)	12 (3%)	1 (0%)	41	31
All	All	768/788 (98%)	743 (97%)	24 (3%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	106	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/328 (98%)	314 (98%)	6 (2%)	57	51
1	B	320/328 (98%)	316 (99%)	4 (1%)	69	66
All	All	640/656 (98%)	630 (98%)	10 (2%)	62	58

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	123	SER
1	A	248	GLU
1	A	346	LYS
1	A	372	ARG
1	A	384	ILE
1	A	386	LYS
1	B	1	MET
1	B	39	LYS
1	B	107	GLN
1	B	241	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	A	401	1	15,15,16	3.29	6 (40%)	20,22,23	2.22	9 (45%)
2	PLP	B	401	1	15,15,16	2.98	6 (40%)	20,22,23	2.03	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	401	1	-	0/6/6/8	0/1/1/1
2	PLP	B	401	1	-	1/6/6/8	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	PLP	C5-C4	8.54	1.49	1.40
2	A	401	PLP	C3-C2	7.38	1.48	1.40
2	B	401	PLP	C3-C2	6.96	1.47	1.40
2	B	401	PLP	C5-C4	6.45	1.47	1.40
2	B	401	PLP	P-O3P	-3.08	1.43	1.54
2	B	401	PLP	P-O2P	-3.01	1.43	1.54
2	A	401	PLP	P-O2P	-2.92	1.43	1.54
2	A	401	PLP	P-O3P	-2.80	1.44	1.54
2	B	401	PLP	P-O1P	-2.73	1.41	1.50
2	A	401	PLP	P-O1P	-2.66	1.41	1.50
2	B	401	PLP	C3-C4	2.47	1.45	1.40
2	A	401	PLP	C3-C4	2.30	1.44	1.40

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	PLP	C4A-C4-C5	5.86	126.97	120.94
2	B	401	PLP	O4P-C5A-C5	-4.67	100.44	109.35
2	B	401	PLP	C6-N1-C2	3.76	126.14	119.17
2	A	401	PLP	O4P-P-O1P	-3.22	97.44	106.47
2	B	401	PLP	C6-C5-C4	3.01	120.53	118.16
2	A	401	PLP	C6-N1-C2	2.96	124.64	119.17
2	A	401	PLP	O3-C3-C2	2.85	123.71	117.49
2	B	401	PLP	C3-C4-C5	-2.82	115.70	118.74
2	B	401	PLP	C4A-C4-C5	2.65	123.66	120.94
2	A	401	PLP	C6-C5-C4	2.45	120.08	118.16
2	A	401	PLP	C3-C4-C5	-2.39	116.16	118.74
2	B	401	PLP	O3P-P-O1P	2.33	119.80	110.68
2	A	401	PLP	C4A-C4-C3	-2.22	116.74	120.50
2	A	401	PLP	O3P-P-O1P	2.08	118.81	110.68
2	A	401	PLP	O4P-C5A-C5	-2.03	105.48	109.35
2	B	401	PLP	C3-C2-N1	-2.01	118.18	120.77

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	PLP	C4-C5-C5A-O4P

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	PLP	3	0
2	B	401	PLP	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	4
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	38:VAL	C	39:LYS	N	1.64
1	B	106:PHE	C	107:GLN	N	1.17
1	A	106:PHE	C	107:GLN	N	1.16
1	B	105:ALA	C	106:PHE	N	1.12
1	B	44:GLY	C	45:HIS	N	1.09
1	A	105:ALA	C	106:PHE	N	1.03

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	386/394 (97%)	0.63	22 (5%)	23 26	25, 39, 85, 125	0
1	B	386/394 (97%)	0.67	29 (7%)	14 16	24, 39, 86, 121	0
All	All	772/788 (97%)	0.65	51 (6%)	18 20	24, 39, 85, 125	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	331	GLY	12.4
1	A	330	ASN	7.8
1	A	329	GLU	7.7
1	B	385	PHE	5.1
1	B	384	ILE	5.1
1	B	149	LEU	4.5
1	A	124	ALA	4.2
1	A	122	ALA	4.2
1	B	2	ILE	4.1
1	A	302	ARG	4.1
1	A	384	ILE	4.0
1	A	194	PHE	4.0
1	B	194	PHE	3.9
1	B	1	MET	3.8
1	A	2	ILE	3.8
1	A	382	GLY	3.8
1	B	124	ALA	3.7
1	B	382	GLY	3.4
1	B	153	LYS	3.1
1	B	383	GLU	3.1
1	B	302	ARG	3.0
1	A	385	PHE	2.9
1	B	201	GLU	2.8
1	A	317	ARG	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	318	VAL	2.8
1	B	195	LEU	2.6
1	B	122	ALA	2.6
1	A	161	LEU	2.5
1	A	360	ASN	2.5
1	B	200	ILE	2.5
1	B	120	LYS	2.5
1	B	191	PHE	2.5
1	A	383	GLU	2.4
1	B	357	GLU	2.4
1	B	360	ASN	2.4
1	A	125	ASN	2.4
1	B	159	LYS	2.3
1	B	346	LYS	2.3
1	A	328	GLY	2.3
1	B	93	VAL	2.3
1	A	298	ALA	2.3
1	A	1	MET	2.2
1	B	271	SER	2.2
1	B	31	LYS	2.2
1	B	237	ALA	2.2
1	A	346	LYS	2.1
1	B	193	SER	2.1
1	B	359	LEU	2.1
1	B	318	VAL	2.1
1	B	198	GLN	2.0
1	A	358	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MG	B	404	1/1	0.76	0.08	64,64,64,64	0
4	MG	B	405	1/1	0.86	0.10	64,64,64,64	0
3	CL	B	402	1/1	0.87	0.22	47,47,47,47	0
4	MG	B	403	1/1	0.88	0.22	33,33,33,33	0
3	CL	A	402	1/1	0.91	0.19	41,41,41,41	0
2	PLP	B	401	15/16	0.93	0.16	21,64,94,107	0
2	PLP	A	401	15/16	0.95	0.14	27,63,95,111	0

6.5 Other polymers [i](#)

There are no such residues in this entry.