



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:48 am BST

PDB ID : 4Q84
Title : Apo YcaO
Authors : Chekan, J.R.; Nair, S.K.
Deposited on : 2014-04-25
Resolution : 2.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1426 (2.66-2.62)
Clashscore	141614	1472 (2.66-2.62)
Ramachandran outliers	138981	1446 (2.66-2.62)
Sidechain outliers	138945	1446 (2.66-2.62)
RSRZ outliers	127900	1408 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	586	
1	B	586	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9543 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribosomal protein S12 methylthiotransferase accessory factor YcaO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	568	Total	C	N	O	S	0	0	0
			4489	2864	731	878	16			
1	B	573	Total	C	N	O	S	0	0	0
			4538	2898	738	885	17			

- Molecule 2 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Hg	0	0
			2	2		
2	A	2	Total	Hg	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	266	Total	O	0	0
			266	266		
3	B	246	Total	O	0	0
			246	246		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribosomal protein S12 methylthiotransferase accessory factor YcaO



- Molecule 1: Ribosomal protein S12 methylthiotransferase accessory factor YcaO



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.95Å 140.39Å 163.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.64 49.32 – 2.64	Depositor EDS
% Data completeness (in resolution range)	96.0 (50.00-2.64) 96.1 (49.32-2.64)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.44 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.163 , 0.218 0.168 , 0.221	Depositor DCC
R_{free} test set	2181 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	42.3	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9543	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/4599	0.82	8/6256 (0.1%)
1	B	0.73	0/4652	0.81	6/6329 (0.1%)
All	All	0.74	0/9251	0.82	14/12585 (0.1%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	56	CYS	CA-CB-SG	7.54	127.56	114.00
1	B	360	MET	CG-SD-CE	-7.38	88.39	100.20
1	B	55	LEU	CA-CB-CG	7.36	132.22	115.30
1	A	267	ASN	N-CA-CB	-7.35	97.37	110.60
1	A	505	ARG	NE-CZ-NH2	-6.71	116.94	120.30
1	B	105	ASN	CB-CA-C	-6.53	97.34	110.40
1	B	505	ARG	NE-CZ-NH1	6.23	123.42	120.30
1	A	105	ASN	CB-CA-C	-6.21	97.98	110.40
1	B	505	ARG	NE-CZ-NH2	-5.87	117.36	120.30
1	A	49	ARG	NE-CZ-NH1	5.84	123.22	120.30
1	A	286	ARG	NE-CZ-NH1	5.39	122.99	120.30
1	A	286	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	505	ARG	NE-CZ-NH1	5.35	122.97	120.30
1	A	525	LEU	CA-CB-CG	-5.14	103.49	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4489	0	4263	41	0
1	B	4538	0	4306	43	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	266	0	0	11	0
3	B	246	0	0	23	0
All	All	9543	0	8569	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:ALA:N	3:A:723:HOH:O	1.73	1.21
1:A:581:ALA:C	3:A:914:HOH:O	2.01	0.97
1:B:494:SER:OG	3:B:762:HOH:O	1.92	0.86
1:B:55:LEU:HB2	3:B:759:HOH:O	1.83	0.79
1:B:55:LEU:HD22	3:B:892:HOH:O	1.82	0.79
1:B:164:ASN:N	3:B:842:HOH:O	1.94	0.74
1:B:55:LEU:CD2	3:B:892:HOH:O	2.36	0.70
1:B:56:CYS:SG	3:B:704:HOH:O	2.50	0.70
1:A:108:TRP:CE3	1:A:151:ARG:HG2	2.29	0.68
1:B:55:LEU:CB	3:B:759:HOH:O	2.41	0.67
1:A:286:ARG:HD2	3:A:720:HOH:O	1.95	0.67
1:B:428:LEU:CD1	1:B:475:GLY:HA3	2.25	0.67
1:A:209:ILE:HD11	1:A:251:LEU:HD11	1.76	0.66
1:B:12:ALA:N	1:B:14:GLU:OE2	2.30	0.64
1:A:61:LYS:HB2	1:A:289:THR:HG23	1.79	0.64
3:A:907:HOH:O	1:B:526:ASN:HB2	1.98	0.64
1:B:519:ARG:HD3	1:B:524:TYR:OH	1.97	0.63
1:B:127:ARG:NE	3:B:937:HOH:O	2.26	0.63
1:A:519:ARG:HD3	1:A:524:TYR:OH	1.98	0.63
1:B:555:PRO:O	3:B:852:HOH:O	2.15	0.60
1:B:428:LEU:HD12	1:B:475:GLY:HA3	1.84	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:572:ALA:HB2	3:B:704:HOH:O	2.03	0.58
1:B:584:TRP:O	3:B:726:HOH:O	2.18	0.56
1:B:174:ILE:HD12	1:B:386:ILE:CD1	2.36	0.55
1:B:114:ASN:OD1	1:B:114:ASN:N	2.39	0.55
1:B:397:GLU:HG2	3:B:755:HOH:O	2.06	0.55
1:B:191:GLU:O	1:B:195:GLN:HG3	2.06	0.55
1:A:191:GLU:O	1:A:195:GLN:HG3	2.07	0.54
1:A:209:ILE:CD1	1:A:251:LEU:HD11	2.37	0.54
1:A:114:ASN:N	1:A:114:ASN:OD1	2.40	0.52
1:A:560:LEU:HD13	1:A:566:HIS:NE2	2.25	0.52
1:B:93:GLU:HG3	3:B:770:HOH:O	2.10	0.52
1:A:289:THR:HG21	3:A:904:HOH:O	2.10	0.51
1:B:502:ASN:ND2	3:B:932:HOH:O	2.43	0.51
1:A:31:ILE:HG23	1:A:48:ILE:HG23	1.92	0.51
1:A:99:PRO:HG2	3:A:959:HOH:O	2.10	0.51
1:A:98:GLY:O	1:A:151:ARG:NH2	2.44	0.51
1:A:31:ILE:CG2	1:A:48:ILE:HG23	2.40	0.50
1:A:61:LYS:HB2	1:A:289:THR:CG2	2.40	0.49
1:B:286:ARG:HD2	3:B:810:HOH:O	2.14	0.48
1:A:61:LYS:HD2	1:A:289:THR:HG22	1.94	0.48
1:B:197:LEU:HD11	1:B:360:MET:HE2	1.95	0.48
1:A:420:SER:HB2	1:A:422:TRP:CE2	2.49	0.47
1:A:555:PRO:HB3	3:A:953:HOH:O	2.14	0.47
1:B:169:ILE:HG21	1:B:174:ILE:HD11	1.95	0.47
1:A:422:TRP:N	1:A:422:TRP:CD1	2.81	0.47
1:A:190:ASN:HB3	1:A:352:THR:HG22	1.96	0.47
1:B:113:GLU:OE1	1:B:113:GLU:HA	2.15	0.46
1:A:108:TRP:CE3	1:A:151:ARG:CG	2.97	0.46
1:A:151:ARG:HG3	1:A:151:ARG:O	2.13	0.46
1:B:93:GLU:HA	3:B:770:HOH:O	2.15	0.46
1:A:108:TRP:CZ3	1:A:151:ARG:HG2	2.50	0.46
1:B:102:HIS:HD2	3:B:746:HOH:O	1.99	0.46
1:A:209:ILE:HA	1:A:214:ILE:HD12	1.99	0.45
1:B:209:ILE:HA	1:B:214:ILE:HD12	1.97	0.45
1:A:197:LEU:HD11	1:A:360:MET:HE2	1.98	0.45
1:B:169:ILE:CG2	1:B:174:ILE:HD11	2.47	0.45
1:B:560:LEU:HD13	1:B:566:HIS:NE2	2.30	0.45
1:A:333:ASP:OD1	1:B:519:ARG:NH1	2.51	0.44
1:B:171:MET:HE1	3:B:925:HOH:O	2.17	0.44
1:A:519:ARG:NH1	1:B:333:ASP:OD1	2.50	0.44
1:B:113:GLU:CG	3:B:878:HOH:O	2.65	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:108:TRP:CE3	1:B:151:ARG:HG2	2.53	0.43
1:A:560:LEU:HD13	1:A:566:HIS:CE1	2.53	0.43
1:A:197:LEU:CD1	1:A:360:MET:HE2	2.49	0.43
1:B:560:LEU:HD13	1:B:566:HIS:CE1	2.54	0.43
1:A:50:ASP:HB2	1:A:56:CYS:HB2	2.00	0.43
1:A:23:LYS:HE2	3:A:855:HOH:O	2.18	0.42
1:A:61:LYS:HB2	3:A:949:HOH:O	2.18	0.42
1:B:479:GLU:OE1	1:B:479:GLU:N	2.51	0.42
1:A:178:TYR:O	1:A:179:VAL:C	2.58	0.42
1:B:102:HIS:CD2	3:B:746:HOH:O	2.73	0.42
1:A:479:GLU:N	1:A:479:GLU:OE1	2.52	0.42
1:A:136:LEU:HD12	1:A:140:MET:SD	2.60	0.41
1:A:98:GLY:HA3	3:A:962:HOH:O	2.19	0.41
1:A:105:ASN:ND2	3:A:727:HOH:O	2.53	0.41
1:A:219:ILE:HA	1:A:220:PRO:HD3	1.94	0.41
1:B:113:GLU:HG3	3:B:878:HOH:O	2.19	0.41
1:A:58:THR:HG21	1:A:76:TYR:HA	2.02	0.41
1:B:55:LEU:CD1	3:B:892:HOH:O	2.68	0.41
1:B:58:THR:HG21	1:B:76:TYR:HA	2.02	0.41
1:B:80:LEU:HA	3:B:931:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	564/586 (96%)	548 (97%)	15 (3%)	1 (0%)	47	64
1	B	571/586 (97%)	551 (96%)	19 (3%)	1 (0%)	47	64
All	All	1135/1172 (97%)	1099 (97%)	34 (3%)	2 (0%)	47	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	179	VAL
1	B	179	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	472/487 (97%)	456 (97%)	16 (3%)	37	53
1	B	477/487 (98%)	462 (97%)	15 (3%)	40	58
All	All	949/974 (97%)	918 (97%)	31 (3%)	38	55

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	SER
1	A	39	PRO
1	A	48	ILE
1	A	49	ARG
1	A	55	LEU
1	A	119	GLU
1	A	164	ASN
1	A	251	LEU
1	A	289	THR
1	A	293	GLN
1	A	406	MET
1	A	494	SER
1	A	499	GLU
1	A	525	LEU
1	A	558	SER
1	A	560	LEU
1	B	13	LEU
1	B	25	SER
1	B	39	PRO
1	B	55	LEU
1	B	113	GLU
1	B	125	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	189	ARG
1	B	293	GLN
1	B	401	LEU
1	B	405	SER
1	B	406	MET
1	B	449	LEU
1	B	494	SER
1	B	558	SER
1	B	584	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	102	HIS
1	A	105	ASN
1	A	145	GLN
1	A	429	ASN
1	B	42	ASN
1	B	102	HIS
1	B	347	ASN
1	B	429	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	568/586 (96%)	-0.29	1 (0%) 95 96	24, 41, 69, 94	0
1	B	573/586 (97%)	-0.22	1 (0%) 95 96	24, 40, 69, 95	0
All	All	1141/1172 (97%)	-0.26	2 (0%) 95 96	24, 41, 69, 95	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	113	GLU	2.5
1	A	298	LYS	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HG	B	602	1/1	0.98	0.10	95,95,95,95	0
2	HG	A	601	1/1	0.99	0.08	86,86,86,86	0
2	HG	A	602	1/1	1.00	0.09	88,88,88,88	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	HG	B	601	1/1	1.00	0.10	86,86,86,86	0

6.5 Other polymers [i](#)

There are no such residues in this entry.