



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 12:47 am BST

PDB ID : 3Q9L
Title : The structure of the dimeric E.coli MinD-ATP complex
Authors : Wu, W.; Park, K.-T.; Lutkenhaus, J.; Holyoak, T.
Deposited on : 2011-01-08
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	<div> <div>10%</div> <div> <div></div> <div>77%</div> <div>19%</div> <div>..</div> </div> </div>
1	B	260	<div> <div>10%</div> <div> <div></div> <div>79%</div> <div>17%</div> <div>..</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4241 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Septum site-determining protein minD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	1	0
			1968	1232	345	384	7			
1	B	257	Total	C	N	O	S	0	0	0
			1969	1232	345	385	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	ALA	ASP	ENGINEERED MUTATION	UNP P0AEZ3
B	40	ALA	ASP	ENGINEERED MUTATION	UNP P0AEZ3

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	130	Total	O	0	0
			130	130		
4	B	110	Total	O	0	0
			110	110		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Septum site-determining protein minD



- Molecule 1: Septum site-determining protein minD



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.66Å 86.59Å 110.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.03 – 2.34 29.03 – 2.34	Depositor EDS
% Data completeness (in resolution range)	96.0 (29.03-2.34) 96.2 (29.03-2.34)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.34Å)	Xtriage
Refinement program	PHENIX 1.6.4 _486	Depositor
R, R_{free}	0.266 , 0.307 0.258 , 0.303	Depositor DCC
R_{free} test set	1672 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	36.0	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4241	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 62.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1172e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	1/1991 (0.1%)	0.71	6/2688 (0.2%)
1	B	0.50	2/1989 (0.1%)	0.71	5/2685 (0.2%)
All	All	0.50	3/3980 (0.1%)	0.71	11/5373 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	92	ARG	CG-CD	-6.26	1.36	1.51
1	B	92	ARG	CG-CD	-6.04	1.36	1.51
1	B	54	ARG	CB-CG	-5.08	1.38	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	92	ARG	NE-CZ-NH2	11.48	126.04	120.30
1	A	92	ARG	NE-CZ-NH1	-10.73	114.94	120.30
1	A	92	ARG	NE-CZ-NH2	10.41	125.51	120.30
1	B	92	ARG	NE-CZ-NH1	-10.22	115.19	120.30
1	B	54	ARG	CB-CA-C	-5.97	98.46	110.40
1	A	54	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	A	92	ARG	CA-CB-CG	-5.62	101.04	113.40
1	B	54	ARG	CA-CB-CG	5.25	124.94	113.40
1	B	92	ARG	CA-CB-CG	-5.20	101.97	113.40
1	A	54	ARG	CG-CD-NE	-5.13	101.03	111.80
1	A	92	ARG	CD-NE-CZ	-5.12	116.43	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1968	0	2021	70	0
1	B	1969	0	2020	62	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	0	12	0	0
3	B	31	0	12	0	0
4	A	130	0	0	4	0
4	B	110	0	0	1	0
All	All	4241	0	4065	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:ARG:HH11	1:B:204:ARG:HG3	0.98	1.14
1:A:92:ARG:NH2	1:B:159:ILE:HG12	1.68	1.08
1:A:204:ARG:HH11	1:A:204:ARG:HG3	0.95	1.04
1:A:159:ILE:HG12	1:B:92:ARG:NH2	1.75	1.01
1:A:204:ARG:HH11	1:A:204:ARG:CG	1.73	1.01
1:B:204:ARG:HH11	1:B:204:ARG:CG	1.75	0.99
1:A:92:ARG:HH12	1:B:163:LYS:HD2	1.29	0.96
1:A:92:ARG:NH1	1:B:163:LYS:HD2	1.80	0.95
1:B:204:ARG:NH1	1:B:204:ARG:HG3	1.72	0.94
1:A:92:ARG:HH12	1:B:163:LYS:CD	1.81	0.91
1:B:92:ARG:HG3	1:B:92:ARG:O	1.67	0.90
1:A:204:ARG:HG3	1:A:204:ARG:NH1	1.68	0.87
1:A:163:LYS:HD2	1:B:92:ARG:NH1	1.94	0.82
1:A:92:ARG:CG	1:A:92:ARG:O	2.29	0.81
1:A:153:SER:O	1:A:157:LEU:HD12	1.81	0.80
1:B:153:SER:O	1:B:157:LEU:HD12	1.83	0.78
1:A:92:ARG:NH2	1:B:159:ILE:CG1	2.44	0.77
1:A:92:ARG:HH21	1:B:159:ILE:CD1	1.98	0.77
1:A:163:LYS:HD2	1:B:92:ARG:HH12	1.47	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:ARG:HG2	1:A:92:ARG:O	1.86	0.75
1:B:115:GLU:HG3	4:B:332:HOH:O	1.86	0.73
1:A:92:ARG:NH1	1:B:163:LYS:CD	2.48	0.72
1:A:203:LEU:O	1:A:204:ARG:HB2	1.86	0.72
1:A:159:ILE:CD1	1:B:92:ARG:HH21	2.02	0.72
1:A:159:ILE:CG1	1:B:92:ARG:NH2	2.50	0.71
1:A:163:LYS:CD	1:B:92:ARG:HH12	2.03	0.71
1:B:203:LEU:O	1:B:204:ARG:HB2	1.91	0.70
1:A:93:ASP:HB3	1:A:96:ALA:HB2	1.73	0.69
1:A:127:THR:HA	1:A:130:LEU:HD12	1.75	0.69
1:B:93:ASP:HB3	1:B:96:ALA:HB2	1.74	0.68
1:A:94:LYS:HD3	1:B:127:THR:HG23	1.75	0.68
1:A:41:ILE:HD13	1:A:97:LEU:HD11	1.76	0.67
1:A:92:ARG:HH21	1:B:159:ILE:HG12	1.58	0.67
1:A:115:GLU:HG3	4:A:332:HOH:O	1.94	0.66
1:B:127:THR:HA	1:B:130:LEU:HD12	1.79	0.65
1:A:127:THR:HG23	1:B:94:LYS:HD3	1.77	0.64
1:A:95:ASP:N	4:A:356:HOH:O	2.29	0.64
1:B:41:ILE:HD13	1:B:97:LEU:HD11	1.78	0.64
1:B:131:MET:HE1	1:B:135:PHE:HE1	1.64	0.61
1:A:92:ARG:O	1:A:92:ARG:HG3	1.99	0.61
1:A:92:ARG:HH21	1:B:159:ILE:CG1	2.09	0.59
1:A:131:MET:HE1	1:A:135:PHE:HE1	1.68	0.57
1:A:220:ALA:HB1	1:A:225:GLU:O	2.06	0.55
1:A:36:VAL:HG22	1:A:46:LEU:HD21	1.89	0.55
1:A:133:LEU:HD12	1:A:133:LEU:C	2.26	0.55
1:A:26:GLY:HA3	1:A:243:VAL:HG21	1.88	0.55
1:A:92:ARG:HH21	1:B:159:ILE:HD13	1.71	0.55
1:B:133:LEU:C	1:B:133:LEU:HD12	2.27	0.54
1:A:159:ILE:HG12	1:B:92:ARG:HH22	1.66	0.54
1:B:172:GLU:HG3	1:B:173:PRO:HD2	1.90	0.54
1:B:145:PRO:HG3	1:B:180:LEU:HD22	1.90	0.53
1:B:26:GLY:HA3	1:B:243:VAL:HG21	1.91	0.53
1:A:126:GLU:O	1:A:130:LEU:HG	2.07	0.53
1:B:89:SER:OG	1:B:91:THR:HG22	2.08	0.53
1:A:94:LYS:NZ	1:B:94:LYS:HE2	2.24	0.53
1:A:92:ARG:HH22	1:B:159:ILE:HG23	1.75	0.52
1:A:163:LYS:CD	1:B:92:ARG:NH1	2.64	0.52
1:A:53:GLU:OE2	1:B:151:ARG:NH1	2.42	0.51
1:A:92:ARG:O	1:A:93:ASP:HB2	2.11	0.51
1:B:56:VAL:HA	1:B:75:ILE:HD11	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:LYS:HE2	1:B:94:LYS:NZ	2.26	0.51
1:A:93:ASP:CG	4:A:356:HOH:O	2.48	0.51
1:B:36:VAL:HG22	1:B:46:LEU:HD21	1.93	0.50
1:A:89:SER:OG	1:A:91:THR:HG22	2.12	0.50
1:A:172:GLU:HG3	1:A:173:PRO:HD2	1.94	0.49
1:A:42:GLY:HA3	1:B:125:ILE:HD11	1.94	0.49
1:A:56:VAL:HA	1:A:75:ILE:HD11	1.94	0.49
1:B:92:ARG:O	1:B:92:ARG:CG	2.51	0.48
1:A:159:ILE:CD1	1:B:92:ARG:NH2	2.72	0.48
1:A:25:THR:O	1:A:29:GLN:HG3	2.14	0.48
1:A:94:LYS:NZ	1:A:127:THR:HG21	2.30	0.47
1:B:25:THR:O	1:B:29:GLN:HG3	2.15	0.47
1:B:94:LYS:NZ	1:B:127:THR:HG21	2.29	0.46
1:A:92:ARG:HD3	1:A:92:ARG:HH11	1.38	0.46
1:A:159:ILE:HD13	1:B:92:ARG:HH21	1.79	0.45
1:B:203:LEU:HD23	1:B:203:LEU:HA	1.77	0.44
1:B:94:LYS:HZ3	1:B:127:THR:HG21	1.83	0.44
1:A:151:ARG:NH1	1:B:53:GLU:OE2	2.51	0.44
1:B:220:ALA:HB1	1:B:225:GLU:O	2.18	0.43
1:A:53:GLU:OE2	1:B:151:ARG:NH2	2.51	0.43
1:B:81:GLU:O	1:B:82:ASN:OD1	2.37	0.43
1:A:157:LEU:HD21	1:A:205:ILE:HG21	1.99	0.43
1:A:92:ARG:HG2	1:A:92:ARG:H	1.67	0.43
1:A:33:LYS:HA	1:A:82:ASN:HB2	1.99	0.43
1:B:157:LEU:HD21	1:B:205:ILE:HG21	2.00	0.43
1:A:145:PRO:HG3	1:A:180:LEU:HD22	2.00	0.42
1:B:204:ARG:NH1	1:B:204:ARG:CG	2.45	0.42
1:A:41:ILE:N	1:A:41:ILE:HD12	2.35	0.42
1:B:33:LYS:HA	1:B:82:ASN:HB2	2.02	0.42
1:A:94:LYS:HZ3	1:A:127:THR:HG21	1.84	0.41
1:B:95:ASP:O	1:B:96:ALA:C	2.59	0.41
1:A:245:ARG:HA	1:A:249:GLU:O	2.20	0.41
1:A:95:ASP:O	1:A:96:ALA:C	2.58	0.41
1:B:126:GLU:O	1:B:130:LEU:HG	2.21	0.41
1:B:94:LYS:HB3	1:B:94:LYS:HE3	1.80	0.41
1:A:151:ARG:NH2	1:B:53:GLU:OE2	2.54	0.40
1:A:208:VAL:O	1:A:245:ARG:HD2	2.22	0.40
1:A:11:LYS:HB3	4:A:383:HOH:O	2.20	0.40
1:A:178:LEU:HB2	1:A:205:ILE:HG13	2.04	0.40
1:A:79:ARG:HD3	1:A:229:LEU:CD1	2.50	0.40
1:B:245:ARG:HA	1:B:249:GLU:O	2.22	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:LEU:HD21	1:A:205:ILE:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/260 (98%)	241 (94%)	13 (5%)	2 (1%)	19	20
1	B	255/260 (98%)	241 (94%)	12 (5%)	2 (1%)	19	20
All	All	511/520 (98%)	482 (94%)	25 (5%)	4 (1%)	19	20

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	SER
1	A	93	ASP
1	B	9	SER
1	B	93	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/216 (99%)	202 (95%)	11 (5%)	23	28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	213/216 (99%)	204 (96%)	9 (4%)	30	37
All	All	426/432 (99%)	406 (95%)	20 (5%)	26	33

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	PHE
1	A	82	ASN
1	A	92	ARG
1	A	94	LYS
1	A	95	ASP
1	A	99	ARG
1	A	150	VAL
1	A	157	LEU
1	A	204	ARG
1	A	216	SER
1	A	254	ARG
1	B	60	PHE
1	B	82	ASN
1	B	92	ARG
1	B	94	LYS
1	B	150	VAL
1	B	157	LEU
1	B	204	ARG
1	B	216	SER
1	B	254	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ATP	A	800	2	26,33,33	0.93	2 (7%)	31,52,52	1.43	6 (19%)
3	ATP	B	800	2	26,33,33	1.06	1 (3%)	31,52,52	1.36	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	800	2	-	0/18/38/38	0/3/3/3
3	ATP	B	800	2	-	0/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	800	ATP	C5-C4	2.80	1.48	1.40
3	A	800	ATP	C2'-C1'	-2.34	1.50	1.53
3	A	800	ATP	C5-C4	2.05	1.46	1.40

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	800	ATP	N3-C2-N1	-3.83	122.69	128.68
3	B	800	ATP	PB-O3B-PG	-2.76	123.35	132.83
3	B	800	ATP	C4-C5-N7	-2.76	106.52	109.40
3	A	800	ATP	PA-O3A-PB	-2.65	123.74	132.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	800	ATP	PA-O3A-PB	-2.58	123.97	132.83
3	B	800	ATP	N3-C2-N1	-2.55	124.69	128.68
3	A	800	ATP	O3'-C3'-C4'	-2.43	104.03	111.05
3	A	800	ATP	N6-C6-N1	2.13	122.99	118.57
3	B	800	ATP	O3G-PG-O2G	2.06	115.52	107.64
3	A	800	ATP	PB-O3B-PG	-2.01	125.92	132.83
3	A	800	ATP	C2-N1-C6	2.01	122.19	118.75


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	257/260 (98%)	0.62	25 (9%)	7 12	20, 34, 60, 86	0
1	B	257/260 (98%)	0.56	25 (9%)	7 12	20, 34, 60, 86	0
All	All	514/520 (98%)	0.59	50 (9%)	7 12	20, 34, 60, 86	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	170	GLY	6.5
1	A	95	ASP	5.9
1	B	200	LEU	5.4
1	B	198	ASP	5.3
1	A	186	GLY	4.9
1	B	95	ASP	4.9
1	B	111	ALA	4.8
1	B	204	ARG	4.6
1	A	92	ARG	4.3
1	A	167	ALA	4.2
1	B	69	THR	3.6
1	B	197	GLU	3.5
1	A	94	LYS	3.5
1	A	57	VAL	3.4
1	B	170	GLY	3.2
1	B	23	ILE	3.1
1	B	24	ALA	3.1
1	A	161	ALA	3.1
1	B	250	GLU	3.0
1	A	140	ILE	3.0
1	B	201	GLU	3.0
1	A	172	GLU	3.0
1	A	171	GLU	2.9
1	A	158	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	7	VAL	2.8
1	B	36	VAL	2.8
1	B	107	ASP	2.8
1	A	118	VAL	2.8
1	B	94	LYS	2.8
1	B	199	VAL	2.8
1	A	162	SER	2.7
1	B	140	ILE	2.6
1	A	248	GLY	2.6
1	A	37	ILE	2.5
1	A	189	SER	2.5
1	A	197	GLU	2.5
1	A	90	GLN	2.5
1	B	67	ASP	2.4
1	B	14	VAL	2.4
1	A	165	ARG	2.3
1	B	228	ILE	2.2
1	B	73	ALA	2.2
1	B	104	LYS	2.2
1	B	7	VAL	2.2
1	B	101	GLY	2.2
1	B	91	THR	2.2
1	A	179	LEU	2.2
1	A	132	ALA	2.1
1	A	204	ARG	2.1
1	A	54	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	B	700	1/1	0.96	0.20	27,27,27,27	0
3	ATP	A	800	31/31	0.97	0.13	18,27,37,50	0
3	ATP	B	800	31/31	0.97	0.12	16,25,37,43	0
2	MG	A	700	1/1	0.97	0.24	24,24,24,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.