



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 03:24 pm BST

PDB ID : 4Q9J  
Title : P-glycoprotein cocrystallised with QZ-Val  
Authors : McGrath, A.P.; Szewczyk, P.; Chang, G.  
Deposited on : 2014-05-01  
Resolution : 3.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1257 (3.70-3.50)
Clashscore	141614	1353 (3.70-3.50)
Ramachandran outliers	138981	1307 (3.70-3.50)
Sidechain outliers	138945	1307 (3.70-3.50)
RSRZ outliers	127900	1161 (3.70-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1284	 87% 6% 8%
2	B	6	 50% 33% 17%
2	C	6	 50% 33% 17%
2	D	6	 50% 33% 17%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 9315 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multidrug resistance protein 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1187	9207	5919	1562	1688	38	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	83	GLN	ASN	CONFLICT	UNP P21447
A	87	GLN	ASN	CONFLICT	UNP P21447
A	90	GLN	ASN	CONFLICT	UNP P21447
A	1277	LEU	-	EXPRESSION TAG	UNP P21447
A	1278	GLU	-	EXPRESSION TAG	UNP P21447
A	1279	HIS	-	EXPRESSION TAG	UNP P21447
A	1280	HIS	-	EXPRESSION TAG	UNP P21447
A	1281	HIS	-	EXPRESSION TAG	UNP P21447
A	1282	HIS	-	EXPRESSION TAG	UNP P21447
A	1283	HIS	-	EXPRESSION TAG	UNP P21447
A	1284	HIS	-	EXPRESSION TAG	UNP P21447

- Molecule 2 is a protein called N- $\{[5\text{-AMINO-1-(5-O-PHOSPHONO-BETA-D-ARABINOFURANOSYL)-1H-IMIDAZOL-4-YL]CARBONYL}\}$ -L-ASPARTIC ACID.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
2	C	6	36	24	6	3	3	0	0	0
2	D	6	36	24	6	3	3	0	0	0
2	B	6	36	24	6	3	3	0	0	0



Chain B:  50% 33% 17%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.74Å 138.66Å 189.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.37 – 3.60 78.37 – 3.60	Depositor EDS
% Data completeness (in resolution range)	95.0 (78.37-3.60) 96.6 (78.37-3.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.01 (at 3.58Å)	Xtrriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, $R_{free}$	0.261 , 0.282 0.265 , 0.285	Depositor DCC
$R_{free}$ test set	1361 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	154.0	Xtrriage
Anisotropy	0.331	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.23 , 71.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9315	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	160.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 30F

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.25	0/9376	0.42	0/12671
2	B	1.11	0/15	3.29	2/18 (11.1%)
2	C	1.40	0/15	4.69	5/18 (27.8%)
2	D	1.16	0/15	3.34	3/18 (16.7%)
All	All	0.27	0/9421	0.49	10/12725 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	VAL	N-CA-CB	-8.69	92.39	111.50
2	C	4	VAL	N-CA-C	-8.20	88.85	111.00
2	B	4	VAL	CA-CB-CG2	7.99	122.88	110.90
2	C	4	VAL	CA-CB-CG1	7.95	122.82	110.90
2	C	2	VAL	CA-CB-CG1	7.39	121.99	110.90
2	D	4	VAL	CA-CB-CG2	7.15	121.63	110.90
2	B	2	VAL	CA-CB-CG2	5.62	119.33	110.90
2	C	4	VAL	CB-CA-C	5.43	121.71	111.40
2	D	4	VAL	N-CA-C	-5.38	96.46	111.00
2	D	6	VAL	N-CA-C	-5.30	96.70	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9207	0	9398	42	0
2	B	36	0	27	1	0
2	C	36	0	27	4	0
2	D	36	0	27	3	0
All	All	9315	0	9479	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:ASP:OD2	1:A:901:ARG:NH2	2.31	0.64
1:A:801:ASP:OD2	1:A:1083:TYR:OH	2.15	0.64
1:A:1207:GLU:OE1	1:A:1229:ARG:NH2	2.36	0.58
1:A:1173:SER:OG	1:A:1176:GLN:OE1	2.24	0.56
1:A:604:GLU:OE1	1:A:617:ILE:N	2.37	0.55
1:A:978:VAL:HB	2:C:2:VAL:HG22	1.89	0.55
1:A:997:ALA:O	1:A:1001:ALA:N	2.38	0.55
1:A:396:SER:OG	1:A:403:VAL:O	2.23	0.54
1:A:303:TYR:CE1	2:D:2:VAL:HG21	2.44	0.53
1:A:998:THR:O	1:A:1002:SER:N	2.40	0.53
1:A:1204:THR:OG1	1:A:1205:GLU:N	2.45	0.49
1:A:727:ILE:HD12	1:A:758:LEU:HD23	1.94	0.49
1:A:922:ILE:HB	1:A:923:PRO:HD3	1.94	0.49
1:A:979:PHE:HZ	2:D:2:VAL:HA	1.78	0.49
1:A:1036:VAL:O	1:A:1051:GLY:N	2.47	0.48
1:A:544:ASN:ND2	1:A:544:ASN:O	2.48	0.47
1:A:78:PHE:O	1:A:82:GLY:N	2.47	0.47
1:A:68:MET:SD	2:C:6:VAL:HG12	2.55	0.46
1:A:495:GLU:N	1:A:495:GLU:OE1	2.49	0.46
1:A:734:VAL:O	1:A:738:GLY:N	2.50	0.45
1:A:762:SER:HA	1:A:765:THR:HG22	1.99	0.45
1:A:147:PHE:CD1	1:A:361:VAL:HG13	2.53	0.44
1:A:303:TYR:HE1	2:D:2:VAL:HG21	1.83	0.44
1:A:88:SER:O	1:A:89:THR:OG1	2.30	0.43
1:A:392:ASN:N	1:A:409:LEU:O	2.52	0.43
1:A:893:ALA:HB2	1:A:916:TYR:HE1	1.83	0.43
1:A:721:GLN:HB3	1:A:722:PRO:HD3	2.00	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1032:GLN:NE2	1:A:1055:GLU:OE2	2.52	0.42
1:A:88:SER:OG	1:A:89:THR:N	2.53	0.42
1:A:970:VAL:HG12	1:A:974:PHE:CE2	2.54	0.42
1:A:758:LEU:HA	1:A:761:ILE:HG22	2.01	0.42
1:A:704:TRP:N	1:A:705:PRO:HD2	2.34	0.42
1:A:68:MET:SD	2:C:6:VAL:HA	2.59	0.42
1:A:387:ASN:O	1:A:450:ASP:N	2.52	0.42
1:A:832:ILE:HG13	2:B:6:VAL:HB	2.02	0.41
1:A:978:VAL:HG11	2:C:2:VAL:HG13	2.01	0.41
1:A:1154:ILE:HA	1:A:1157:LEU:HD12	2.01	0.41
1:A:509:ILE:HG13	1:A:510:MET:N	2.35	0.41
1:A:156:ILE:HD11	1:A:904:VAL:HG11	2.02	0.41
1:A:689:PRO:N	1:A:690:PRO:HD2	2.36	0.41
1:A:733:GLY:O	1:A:737:ASN:N	2.49	0.40
1:A:991:ALA:N	1:A:992:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1183/1284 (92%)	1127 (95%)	56 (5%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	978/1065 (92%)	973 (100%)	5 (0%)	88	95
2	B	3/3 (100%)	2 (67%)	1 (33%)	0	2
2	C	3/3 (100%)	2 (67%)	1 (33%)	0	2
2	D	3/3 (100%)	2 (67%)	1 (33%)	0	2
All	All	987/1074 (92%)	979 (99%)	8 (1%)	81	91

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	310	PHE
1	A	339	PHE
1	A	429	LYS
1	A	700	ASN
1	A	979	PHE
2	C	2	VAL
2	D	2	VAL
2	B	2	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1003	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

9 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	30F	C	5	2	3,5,6	2.00	2 (66%)	1,5,7	1.63	0
2	30F	D	5	2	3,5,6	2.29	2 (66%)	1,5,7	1.82	0
2	30F	B	5	2	3,5,6	1.95	1 (33%)	1,5,7	0.06	0
2	30F	D	1	2	3,5,6	1.14	0	1,5,7	0.07	0
2	30F	C	3	2	3,5,6	1.11	0	1,5,7	1.93	0
2	30F	B	3	2	3,5,6	1.33	0	1,5,7	2.05	1 (100%)
2	30F	D	3	2	3,5,6	1.27	0	1,5,7	0.66	0
2	30F	C	1	2	3,5,6	0.82	0	1,5,7	0.99	0
2	30F	B	1	2	3,5,6	1.25	0	1,5,7	2.04	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	30F	C	5	2	-	0/0/4/6	-
2	30F	D	5	2	-	0/0/4/6	-
2	30F	B	5	2	-	0/0/4/6	-
2	30F	D	1	2	-	0/0/4/6	-
2	30F	C	3	2	-	0/0/4/6	-
2	30F	B	3	2	-	0/0/4/6	-
2	30F	D	3	2	-	0/0/4/6	-
2	30F	C	1	2	-	0/0/4/6	-
2	30F	B	1	2	-	0/0/4/6	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	5	30F	C08-N04	2.73	1.42	1.35
2	B	5	30F	C08-N04	2.61	1.41	1.35
2	D	5	30F	C10-C08	2.50	1.49	1.45
2	C	5	30F	C08-N04	2.48	1.41	1.35
2	C	5	30F	C10-C08	2.06	1.48	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	30F	O2-C10-C08	-2.05	122.79	125.39
2	B	1	30F	O2-C10-C08	-2.04	122.80	125.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1187/1284 (92%)	0.03	18 (1%) 73 60	70, 150, 250, 381	0
2	B	3/6 (50%)	0.79	0 100 100	127, 127, 127, 127	0
2	C	3/6 (50%)	-0.38	0 100 100	127, 127, 127, 127	0
2	D	3/6 (50%)	0.14	0 100 100	127, 127, 127, 127	0
All	All	1196/1302 (91%)	0.03	18 (1%) 73 60	70, 150, 250, 381	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	556	ALA	4.0
1	A	99	MET	4.0
1	A	98	ALA	3.9
1	A	486	TYR	3.9
1	A	732	VAL	3.6
1	A	734	VAL	2.8
1	A	324	ILE	2.8
1	A	1033	PHE	2.6
1	A	605	GLN	2.5
1	A	405	ILE	2.4
1	A	1079	LEU	2.2
1	A	778	GLU	2.2
1	A	224	SER	2.2
1	A	706	TYR	2.1
1	A	784	LEU	2.1
1	A	306	TYR	2.0
1	A	351	PHE	2.0
1	A	886	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	30F	C	1	6/7	0.48	0.19	127,127,127,127	0
2	30F	B	3	6/7	0.57	0.28	127,127,127,127	0
2	30F	C	5	6/7	0.77	0.39	127,127,127,127	0
2	30F	C	3	6/7	0.77	0.09	127,127,127,127	0
2	30F	D	3	6/7	0.84	0.14	127,127,127,127	0
2	30F	B	5	6/7	0.89	0.26	127,127,127,127	0
2	30F	D	1	6/7	0.92	0.34	127,127,127,127	0
2	30F	D	5	6/7	0.94	0.11	127,127,127,127	0
2	30F	B	1	6/7	0.94	0.42	127,127,127,127	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.