



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 10, 2022 – 07:24 PM EST

PDB ID : 2QCJ  
Title : Native Structure of Lyp  
Authors : Sun, J.P.; Yu, X.; Zhang, Z.Y.  
Deposited on : 2007-06-19  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.31.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	 2% 22% 53% 14% 11%
1	B	313	 3% 21% 54% 13% 11%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4631 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 22.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	0	0
			2304	1488	374	427	15			
1	B	280	Total	C	N	O	S	0	0	0
			2304	1488	374	427	15			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	HIS	-	expression tag	UNP Q9Y2R2
A	-17	HIS	-	expression tag	UNP Q9Y2R2
A	-16	HIS	-	expression tag	UNP Q9Y2R2
A	-15	HIS	-	expression tag	UNP Q9Y2R2
A	-14	HIS	-	expression tag	UNP Q9Y2R2
A	-13	HIS	-	expression tag	UNP Q9Y2R2
A	-12	SER	-	expression tag	UNP Q9Y2R2
A	-11	SER	-	expression tag	UNP Q9Y2R2
A	-10	GLY	-	expression tag	UNP Q9Y2R2
A	-9	LEU	-	expression tag	UNP Q9Y2R2
A	-8	VAL	-	expression tag	UNP Q9Y2R2
A	-7	PRO	-	expression tag	UNP Q9Y2R2
A	-6	ARG	-	expression tag	UNP Q9Y2R2
A	-5	GLY	-	expression tag	UNP Q9Y2R2
A	-4	SER	-	expression tag	UNP Q9Y2R2
A	-3	HIS	-	expression tag	UNP Q9Y2R2
A	-2	MET	-	expression tag	UNP Q9Y2R2
A	-1	ALA	-	expression tag	UNP Q9Y2R2
A	0	SER	-	expression tag	UNP Q9Y2R2
B	-18	HIS	-	expression tag	UNP Q9Y2R2
B	-17	HIS	-	expression tag	UNP Q9Y2R2
B	-16	HIS	-	expression tag	UNP Q9Y2R2
B	-15	HIS	-	expression tag	UNP Q9Y2R2
B	-14	HIS	-	expression tag	UNP Q9Y2R2
B	-13	HIS	-	expression tag	UNP Q9Y2R2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	SER	-	expression tag	UNP Q9Y2R2
B	-11	SER	-	expression tag	UNP Q9Y2R2
B	-10	GLY	-	expression tag	UNP Q9Y2R2
B	-9	LEU	-	expression tag	UNP Q9Y2R2
B	-8	VAL	-	expression tag	UNP Q9Y2R2
B	-7	PRO	-	expression tag	UNP Q9Y2R2
B	-6	ARG	-	expression tag	UNP Q9Y2R2
B	-5	GLY	-	expression tag	UNP Q9Y2R2
B	-4	SER	-	expression tag	UNP Q9Y2R2
B	-3	HIS	-	expression tag	UNP Q9Y2R2
B	-2	MET	-	expression tag	UNP Q9Y2R2
B	-1	ALA	-	expression tag	UNP Q9Y2R2
B	0	SER	-	expression tag	UNP Q9Y2R2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	12	Total O 12 12	0	0
2	B	11	Total O 11 11	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 22



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 22



I239	I240	D241	Y242	T243	W244	M245	L246	L247	K248	D249	G250	I251	I252	P253	E254	N255	F256	S257	V258	F259	S260	L261	I262	R263	E264	M265	R266	T267	Q268	R269	P270	S271	L272	V273	Q274	T275	Q276	E277	Q278	Y279	E280	L281	V282	Y283	N284	L287	E288	L289	F290	LYS	ARG	GLN	MET
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.31Å 93.95Å 148.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.00 28.85 – 2.79	Depositor EDS
% Data completeness (in resolution range)	92.2 (50.00-3.00) 90.7 (28.85-2.79)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.14	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.18 (at 2.80Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.284 , 0.307 0.284 , 0.306	Depositor DCC
$R_{free}$ test set	990 reflections (6.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.2	Xtriage
Anisotropy	0.227	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 103.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4631	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 58.92 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.9045e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/2360	0.76	2/3191 (0.1%)
1	B	0.49	0/2360	0.74	1/3191 (0.0%)
All	All	0.48	0/4720	0.75	3/6382 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	ILE	N-CA-C	7.74	131.90	111.00
1	A	11	LEU	N-CA-C	-5.66	95.73	111.00
1	B	37	LYS	N-CA-C	-5.18	97.02	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2304	0	2291	301	0
1	B	2304	0	2291	292	0
2	A	12	0	0	2	0
2	B	11	0	0	5	0
All	All	4631	0	4582	588	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 64.



The worst 5 of 588 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:PHE:HD2	1:A:13:GLU:HB2	1.09	1.13
1:B:46:THR:HG21	1:B:64:LEU:HB3	1.28	1.12
1:A:91:LYS:HG3	1:A:96:PRO:HA	1.30	1.10
1:A:46:THR:HG21	1:A:64:LEU:HB3	1.15	1.09
1:B:51:LYS:HG3	1:B:52:PRO:HD2	1.35	1.07

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/313 (88%)	191 (69%)	59 (21%)	26 (9%)	0	3
1	B	276/313 (88%)	205 (74%)	50 (18%)	21 (8%)	1	5
All	All	552/626 (88%)	396 (72%)	109 (20%)	47 (8%)	1	4

5 of 47 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	21	LYS
1	A	42	LYS
1	A	80	GLU
1	A	148	GLU
1	A	193	TRP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/287 (89%)	221 (86%)	35 (14%)	3	17
1	B	256/287 (89%)	221 (86%)	35 (14%)	3	17
All	All	512/574 (89%)	442 (86%)	70 (14%)	3	17

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	197	ASP
1	B	211	ASP
1	B	256	PHE
1	A	211	ASP
1	A	206	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	150	GLN
1	B	276	GLN
1	B	284	ASN
1	B	278	GLN
1	A	276	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	280/313 (89%)	-0.13	6 (2%) 63 34	9, 52, 93, 104	0
1	B	280/313 (89%)	-0.15	8 (2%) 51 23	5, 52, 95, 104	0
All	All	560/626 (89%)	-0.14	14 (2%) 57 29	5, 52, 95, 104	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	7	LEU	3.6
1	A	197	ASP	3.4
1	B	17	LYS	3.4
1	A	37	LYS	3.3
1	B	15	GLN	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.