



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2022 – 10:24 pm BST

PDB ID : 7QHN
Title : CRYSTAL STRUCTURE OF LYSYL-TRNA SYNTHETASE FROM *Mycobacterium tuberculosis* COMPLEXED WITH L-LYSINE and an inhibitor
Authors : Dawson, A.; Robinson, D.A.; Tamjar, J.; Wyatt, P.; Green, S.
Deposited on : 2021-12-13
Resolution : 2.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3676 (2.60-2.56)
Clashscore	141614	4049 (2.60-2.56)
Ramachandran outliers	138981	3979 (2.60-2.56)
Sidechain outliers	138945	3979 (2.60-2.56)
RSRZ outliers	127900	3614 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3604 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lysine-tRNA ligase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	450	Total	C	N	O	S	0	0	0
			3504	2213	622	659	10			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP P9WUFU8
A	-19	GLY	-	expression tag	UNP P9WUFU8
A	-18	SER	-	expression tag	UNP P9WUFU8
A	-17	SER	-	expression tag	UNP P9WUFU8
A	-16	HIS	-	expression tag	UNP P9WUFU8
A	-15	HIS	-	expression tag	UNP P9WUFU8
A	-14	HIS	-	expression tag	UNP P9WUFU8
A	-13	HIS	-	expression tag	UNP P9WUFU8
A	-12	HIS	-	expression tag	UNP P9WUFU8
A	-11	HIS	-	expression tag	UNP P9WUFU8
A	-10	SER	-	expression tag	UNP P9WUFU8
A	-9	SER	-	expression tag	UNP P9WUFU8
A	-8	GLY	-	expression tag	UNP P9WUFU8
A	-7	GLU	-	expression tag	UNP P9WUFU8
A	-6	ASN	-	expression tag	UNP P9WUFU8
A	-5	LEU	-	expression tag	UNP P9WUFU8
A	-4	TYR	-	expression tag	UNP P9WUFU8
A	-3	PHE	-	expression tag	UNP P9WUFU8
A	-2	GLN	-	expression tag	UNP P9WUFU8
A	-1	GLY	-	expression tag	UNP P9WUFU8
A	0	HIS	-	expression tag	UNP P9WUFU8
A	455	ALA	MET	engineered mutation	UNP P9WUFU8

- Molecule 2 is LYSINE (three-letter code: LYS) (formula: $C_6H_{15}N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			10	6	2	2		

- Molecule 3 is 6-azanyl-2-cyclohexyl-4-fluoranyl-1 {H}-pyrrolo[3,4-c]pyridin-3-one (three-letter code: C6I) (formula: C₁₃H₁₆FN₃O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			18	13	1	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	72	Total	O	0	0
			72	72		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lysine-tRNA ligase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	82.89Å 82.89Å 146.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.26 – 2.58 72.16 – 2.58	Depositor EDS
% Data completeness (in resolution range)	99.3 (72.26-2.58) 99.3 (72.16-2.58)	Depositor EDS
R_{merge}	0.29	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.293 , 0.339 0.293 , 0.342	Depositor DCC
R_{free} test set	787 reflections (4.73%)	wwPDB-VP
Wilson B-factor (Å ²)	42.5	Xtriage
Anisotropy	0.687	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	3604	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: C6I

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/3563	0.82	0/4830

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	427	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3504	0	3504	69	1
2	A	10	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	18	0	0	0	0
4	A	72	0	0	4	0
All	All	3604	0	3516	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:LEU:HD23	1:A:95:SER:HA	1.43	0.96
1:A:76:LYS:HE3	1:A:77:LEU:CD1	1.98	0.93
1:A:76:LYS:HE3	1:A:77:LEU:HD12	1.54	0.89
1:A:161:VAL:CG1	1:A:498:PRO:HG2	2.10	0.81
1:A:161:VAL:HG13	1:A:498:PRO:HG2	1.63	0.80
1:A:418:LEU:HD22	1:A:485:SER:HB2	1.69	0.74
1:A:76:LYS:HG3	1:A:77:LEU:HG	1.69	0.73
1:A:17:ARG:NH2	1:A:113:LEU:O	2.26	0.68
1:A:274:THR:HG21	4:A:767:HOH:O	1.94	0.68
1:A:54:ILE:HG23	1:A:126:SER:O	1.95	0.66
1:A:325:TYR:HB2	1:A:326:PRO:HD3	1.76	0.66
1:A:77:LEU:CD2	1:A:95:SER:CB	2.74	0.65
1:A:107:TRP:CE3	1:A:111:VAL:HG21	2.34	0.63
1:A:77:LEU:HD23	1:A:95:SER:CA	2.22	0.62
1:A:54:ILE:CG2	1:A:126:SER:O	2.48	0.62
1:A:462:LEU:O	1:A:466:GLU:HG2	2.00	0.61
1:A:68:VAL:HG12	1:A:115:ASP:O	2.02	0.60
1:A:77:LEU:CD2	1:A:95:SER:HA	2.27	0.59
1:A:441:ALA:O	1:A:445:ARG:HG2	2.03	0.58
1:A:491:ILE:HG22	4:A:744:HOH:O	2.04	0.58
1:A:58:THR:CG2	1:A:123:VAL:HG13	2.36	0.56
1:A:114:GLY:O	1:A:144:SER:HB3	2.05	0.56
1:A:246:PHE:HB3	1:A:249:VAL:HG13	1.88	0.56
1:A:74:SER:HB2	1:A:77:LEU:O	2.07	0.54
1:A:157:GLU:O	1:A:161:VAL:HG23	2.08	0.54
1:A:262:ASP:OD1	1:A:262:ASP:C	2.47	0.53
1:A:206:GLN:NE2	4:A:711:HOH:O	2.42	0.52
1:A:14:PHE:O	1:A:18:ARG:NH1	2.41	0.51
1:A:404:HIS:HB2	1:A:410:VAL:HG23	1.93	0.51
1:A:116:ILE:HD12	1:A:142:ALA:HB3	1.93	0.50
1:A:401:THR:HG23	1:A:427:TYR:HB3	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:THR:C	1:A:275:TYR:CD1	2.86	0.49
1:A:156:SER:OG	1:A:159:SER:OG	2.21	0.49
1:A:68:VAL:HG23	1:A:80:ALA:HB1	1.95	0.49
1:A:13:GLN:HA	1:A:16:ILE:HG22	1.95	0.49
1:A:161:VAL:HG11	1:A:498:PRO:HG2	1.95	0.49
1:A:412:GLU:HB2	1:A:428:SER:HB2	1.94	0.49
1:A:77:LEU:CD2	1:A:95:SER:HB2	2.43	0.48
1:A:13:GLN:HA	1:A:16:ILE:CG2	2.44	0.47
1:A:161:VAL:HG13	1:A:498:PRO:CG	2.38	0.47
1:A:69:ILE:HD11	1:A:81:THR:CG2	2.45	0.47
1:A:77:LEU:HD21	1:A:95:SER:HB2	1.95	0.47
1:A:57:ALA:HA	1:A:124:ILE:HG22	1.96	0.46
1:A:76:LYS:CE	1:A:77:LEU:CD1	2.85	0.46
1:A:58:THR:HG22	1:A:123:VAL:HG13	1.98	0.45
1:A:325:TYR:CB	1:A:326:PRO:HD3	2.44	0.45
1:A:369:GLY:HA2	1:A:394:PRO:CB	2.47	0.45
1:A:279:GLY:HA2	1:A:283:ASP:OD2	2.16	0.45
1:A:418:LEU:CD2	1:A:485:SER:HB2	2.42	0.45
1:A:40:HIS:CE1	1:A:62:VAL:HG12	2.52	0.45
1:A:414:TRP:CE2	1:A:426:GLY:HA3	2.52	0.44
1:A:496:LEU:HB2	4:A:710:HOH:O	2.17	0.44
1:A:167:ASP:OD1	1:A:171:ARG:NE	2.49	0.44
1:A:112:ASP:OD2	1:A:112:ASP:N	2.51	0.43
1:A:210:GLY:HA3	1:A:236:LEU:CD1	2.48	0.43
1:A:76:LYS:HE3	1:A:77:LEU:HD11	1.91	0.43
1:A:77:LEU:CD2	1:A:95:SER:CA	2.92	0.43
1:A:271:MET:HE2	1:A:273:GLU:CG	2.48	0.43
1:A:246:PHE:HB3	1:A:249:VAL:CG1	2.49	0.42
1:A:342:VAL:O	1:A:346:ARG:HB2	2.19	0.42
1:A:321:THR:HA	1:A:389:PHE:O	2.19	0.42
1:A:41:THR:O	1:A:44:GLU:HB2	2.20	0.42
1:A:291:LEU:HD13	1:A:291:LEU:C	2.41	0.41
1:A:369:GLY:O	1:A:373:GLU:HG3	2.20	0.41
1:A:289:ARG:HA	1:A:319:TRP:CZ3	2.56	0.41
1:A:17:ARG:HG2	1:A:112:ASP:OD1	2.19	0.41
1:A:33:PRO:HD2	1:A:141:ALA:O	2.20	0.41
1:A:66:GLY:HA3	1:A:82:LEU:HB3	2.04	0.40
1:A:414:TRP:CZ2	1:A:426:GLY:HA3	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:TYR:OH	1:A:460:ASP:OD1[8_555]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	431/526 (82%)	397 (92%)	32 (7%)	2 (0%)	29 50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	53	PRO
1	A	297	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	372/432 (86%)	356 (96%)	16 (4%)	29 52

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	GLU
1	A	16	ILE
1	A	67	ARG
1	A	110	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	112	ASP
1	A	132	SER
1	A	170	VAL
1	A	254	ARG
1	A	257	ARG
1	A	263	SER
1	A	274	THR
1	A	281	TYR
1	A	309	ASP
1	A	438	GLU
1	A	443	GLN
1	A	480	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	ASN
1	A	206	GLN
1	A	276	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LYS	A	601	-	8,9,9	0.83	1 (12%)	9,10,10	0.66	0
3	C6I	A	602	-	20,20,20	1.67	5 (25%)	25,29,29	3.68	11 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LYS	A	601	-	-	2/9/9/9	-
3	C6I	A	602	-	-	4/4/24/24	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	C6I	C6-C7	-3.85	1.45	1.50
3	A	602	C6I	C6-N	-3.53	1.44	1.46
3	A	602	C6I	C10-N2	3.25	1.35	1.31
3	A	602	C6I	O-C12	2.95	1.28	1.22
3	A	602	C6I	C12-N	-2.40	1.33	1.36
2	A	601	LYS	OXT-C	-2.30	1.23	1.30

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	C6I	C3-N-C12	12.98	136.89	122.96
3	A	602	C6I	C7-C6-N	-5.71	100.28	102.18
3	A	602	C6I	C11-C10-N2	-5.40	118.59	126.64
3	A	602	C6I	C10-C11-C12	4.62	138.50	131.32
3	A	602	C6I	O-C12-C11	-4.14	123.20	129.09
3	A	602	C6I	C2-C3-N	4.00	117.56	111.85
3	A	602	C6I	C10-N2-C9	3.63	122.25	118.06
3	A	602	C6I	F-C10-N2	2.87	119.58	116.15
3	A	602	C6I	N1-C9-N2	2.61	120.61	116.49
3	A	602	C6I	C4-C3-C2	-2.37	105.74	111.19
3	A	602	C6I	C4-C3-N	-2.36	108.47	111.85

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	C6I	C2-C3-N-C6
3	A	602	C6I	C2-C3-N-C12
3	A	602	C6I	C4-C3-N-C6
3	A	602	C6I	C4-C3-N-C12
2	A	601	LYS	O-C-CA-CB
2	A	601	LYS	OXT-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	450/526 (85%)	1.12	67 (14%) 2 1	29, 57, 110, 132	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	127	ARG	5.3
1	A	309	ASP	4.9
1	A	28	GLY	4.9
1	A	57	ALA	4.7
1	A	71	ALA	4.6
1	A	82	LEU	4.3
1	A	109	ALA	4.1
1	A	113	LEU	4.0
1	A	455	ALA	4.0
1	A	133	VAL	4.0
1	A	110	ASP	3.9
1	A	444	ALA	3.8
1	A	32	TYR	3.7
1	A	25	LEU	3.6
1	A	499	ILE	3.6
1	A	58	THR	3.6
1	A	117	VAL	3.5
1	A	95	SER	3.5
1	A	445	ARG	3.5
1	A	22	ALA	3.4
1	A	155	MET	3.4
1	A	145	LEU	3.3
1	A	76	LYS	3.3
1	A	14	PHE	3.1
1	A	119	VAL	3.1
1	A	112	ASP	3.0
1	A	131	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	456	VAL	3.0
1	A	134	LEU	2.8
1	A	443	GLN	2.7
1	A	118	TYR	2.7
1	A	105	ASP	2.7
1	A	70	PHE	2.7
1	A	102	ALA	2.7
1	A	29	ARG	2.7
1	A	400	LEU	2.6
1	A	78	CYS	2.6
1	A	77	LEU	2.5
1	A	104	LEU	2.5
1	A	333	GLY	2.5
1	A	308	PRO	2.5
1	A	74	SER	2.5
1	A	90	LEU	2.5
1	A	162	ARG	2.5
1	A	69	ILE	2.4
1	A	226	ILE	2.4
1	A	73	ASN	2.3
1	A	140	ILE	2.3
1	A	180	LEU	2.3
1	A	462	LEU	2.3
1	A	204	VAL	2.3
1	A	126	SER	2.3
1	A	123	VAL	2.2
1	A	72	ARG	2.2
1	A	111	VAL	2.2
1	A	142	ALA	2.2
1	A	150	VAL	2.1
1	A	42	LEU	2.1
1	A	107	TRP	2.1
1	A	208	LEU	2.1
1	A	138	TRP	2.1
1	A	425	THR	2.1
1	A	125	SER	2.1
1	A	176	ALA	2.0
1	A	68	VAL	2.0
1	A	366	PHE	2.0
1	A	31	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	LYS	A	601	10/10	0.88	0.27	43,52,57,57	0
3	C6I	A	602	18/18	0.93	0.22	43,46,51,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.