



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 05:25 pm BST

PDB ID : 2QJW
Title : CRYSTAL STRUCTURE OF A PUTATIVE HYDROLASE OF THE ALPHA/BETA SUPERFAMILY (XCC1541) FROM XANTHOMONAS CAMPESTRIS PV. CAMPESTRIS AT 1.35 Å RESOLUTION
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2007-07-09
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	<div> <div>0%</div> <div> <div>89%</div> <div>11%</div> </div> </div>
1	B	176	<div> <div>0%</div> <div> <div>92%</div> <div>7%</div> </div> </div>
1	C	176	<div> <div>0%</div> <div> <div>93%</div> <div>5%</div> </div> </div>
1	D	176	<div> <div>2%</div> <div> <div>88%</div> <div>9%</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6211 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein XCC1541.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	176	Total	C	N	O	S	Se	0	10	0
			1380	875	247	252	1	5			
1	B	176	Total	C	N	O	S	Se	0	7	0
			1362	861	250	245	1	5			
1	C	173	Total	C	N	O	S	Se	0	5	0
			1316	839	237	236	1	3			
1	D	172	Total	C	N	O	S	Se	0	6	0
			1321	837	238	241	1	4			


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP Q8PAE4
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
A	102	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
A	108	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
B	0	GLY	-	LEADER SEQUENCE	UNP Q8PAE4
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
B	102	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
B	108	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
C	0	GLY	-	LEADER SEQUENCE	UNP Q8PAE4
C	1	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
C	102	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
C	108	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
D	0	GLY	-	LEADER SEQUENCE	UNP Q8PAE4
D	1	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
D	102	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4
D	108	MSE	MET	MODIFIED RESIDUE	UNP Q8PAE4

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Cl	0	0
			3	3		
2	A	3	Total	Cl	0	0
			3	3		
2	C	2	Total	Cl	0	0
			2	2		

- Molecule 3 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: $C_4H_6O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			10	4	6		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			7	4	3		
4	C	1	Total	C	O	0	0
			7	4	3		
4	C	1	Total	C	O	0	0
			7	4	3		
4	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: C₁₂H₂₆O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			19	12	7		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	189	Total	O	0	0
			189	189		
6	B	199	Total	O	0	0
			199	199		
6	C	178	Total	O	0	0
			178	178		
6	D	201	Total	O	0	0
			201	201		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uncharacterized protein XCC1541




- Molecule 1: Uncharacterized protein XCC1541



- Molecule 1: Uncharacterized protein XCC1541



- Molecule 1: Uncharacterized protein XCC1541



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.80Å 60.64Å 74.72Å 90.00° 94.57° 90.00°	Depositor
Resolution (Å)	29.77 – 1.35 29.77 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.77-1.35) 99.7 (29.77-1.35)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.2.0019, PHENIX	Depositor
R, R_{free}	0.182 , 0.216 0.188 , 0.219	Depositor DCC
R_{free} test set	7030 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.024 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6211	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TLA, PEG, P6G, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/1404	0.95	3/1906 (0.2%)
1	B	0.88	2/1386 (0.1%)	0.98	7/1881 (0.4%)
1	C	0.80	0/1341	0.93	6/1823 (0.3%)
1	D	0.88	1/1344 (0.1%)	0.92	1/1824 (0.1%)
All	All	0.85	3/5475 (0.1%)	0.95	17/7434 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	108[A]	MSE	CG-SE	-6.54	1.73	1.95
1	B	108[B]	MSE	CG-SE	-6.54	1.73	1.95
1	D	85	TYR	CE2-CZ	-5.57	1.31	1.38

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	147	ARG	NE-CZ-NH2	-8.12	116.24	120.30
1	C	144[A]	ARG	NE-CZ-NH1	6.91	123.75	120.30
1	C	144[B]	ARG	NE-CZ-NH1	6.91	123.75	120.30
1	A	54	ASP	CB-CG-OD2	-6.90	112.09	118.30
1	C	144[A]	ARG	NE-CZ-NH2	-6.07	117.27	120.30
1	C	144[B]	ARG	NE-CZ-NH2	-6.07	117.27	120.30
1	A	54	ASP	CB-CG-OD1	6.00	123.70	118.30
1	B	144[A]	ARG	NE-CZ-NH1	5.89	123.25	120.30
1	B	144[B]	ARG	NE-CZ-NH1	5.89	123.25	120.30
1	B	144[A]	ARG	NE-CZ-NH2	-5.83	117.38	120.30
1	B	144[B]	ARG	NE-CZ-NH2	-5.83	117.38	120.30
1	C	58	ARG	NE-CZ-NH1	5.74	123.17	120.30
1	A	58	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	B	58	ARG	NE-CZ-NH2	-5.57	117.52	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	37	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	B	152	ASP	CB-CG-OD2	-5.33	113.50	118.30
1	B	39	ASP	CB-CG-OD1	5.29	123.06	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1380	0	1371	12	0
1	B	1362	0	1365	7	0
1	C	1316	0	1307	13	0
1	D	1321	0	1318	11	0
2	A	3	0	0	0	0
2	B	3	0	0	1	0
2	C	2	0	0	1	0
3	B	10	0	4	0	0
4	B	7	0	10	0	0
4	C	14	0	20	3	0
4	D	7	0	10	0	0
5	C	19	0	26	0	0
6	A	189	0	0	1	0
6	B	199	0	0	0	0
6	C	178	0	0	2	0
6	D	201	0	0	2	0
All	All	6211	0	5431	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:52:LEU:HD12	1:B:108[B]:MSE:HE1	1.52	0.89

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:121:ILE:HD12	1:C:144[A]:ARG:HD3	1.73	0.68
1:C:108[B]:MSE:HE3	1:C:108[B]:MSE:HA	1.76	0.67
1:C:147[B]:ARG:HE	1:D:149:LEU:HD11	1.70	0.57
1:A:94[A]:VAL:HG22	1:A:95:PRO:HD2	1.87	0.56
1:D:52:LEU:HD12	1:D:108[B]:MSE:HE1	1.88	0.56
1:C:108[B]:MSE:HA	1:C:108[B]:MSE:CE	2.37	0.55
1:C:120:PRO:HG3	4:C:179:PEG:H42	1.89	0.55
1:D:27:VAL:HG22	1:D:30[B]:ARG:NH1	2.22	0.55
1:B:80:SER:HB2	1:B:102:MSE:HE3	1.92	0.52
1:C:108[B]:MSE:HG3	1:C:111:LEU:HD12	1.92	0.51
1:A:94[B]:VAL:HG13	1:A:95:PRO:HD2	1.92	0.51
1:A:70:THR:HG21	1:A:94[B]:VAL:CG1	2.40	0.51
1:D:80[A]:SER:HB2	1:D:102:MSE:HE3	1.93	0.50
1:B:47:ARG:O	2:B:178:CL:CL	2.68	0.49
1:B:52:LEU:CD1	1:B:108[B]:MSE:HE1	2.33	0.49
1:C:121:ILE:HD12	1:C:144[A]:ARG:CD	2.42	0.48
1:A:170[A]:GLU:HG3	6:A:260:HOH:O	2.14	0.47
1:D:142:GLN:HG2	6:D:356:HOH:O	2.14	0.47
1:A:80:SER:HB2	1:A:102:MSE:HE3	1.97	0.47
1:A:103:VAL:HA	1:A:125:HIS:ND1	2.30	0.47
1:D:52:LEU:CD1	1:D:108[B]:MSE:HE1	2.45	0.46
1:C:85:TYR:CE1	1:C:108[A]:MSE:HG3	2.50	0.46
1:A:70:THR:HG21	1:A:94[B]:VAL:HG12	1.96	0.46
1:A:52:LEU:HD12	1:A:108[A]:MSE:SE	2.67	0.45
1:B:103:VAL:HA	1:B:125:HIS:ND1	2.31	0.44
1:D:147:ARG:NH2	1:D:170[B]:GLU:OE1	2.51	0.43
4:C:179:PEG:C3	6:D:249:HOH:O	2.66	0.43
1:D:14:SER:HB2	1:D:18:ALA:HB2	2.01	0.43
1:D:111:LEU:HD12	1:D:111:LEU:N	2.34	0.42
1:C:170[B]:GLU:HG3	6:C:229:HOH:O	2.18	0.42
1:B:121:ILE:HD12	1:B:144[A]:ARG:CD	2.49	0.42
1:C:85:TYR:CD1	1:C:108[A]:MSE:HG3	2.55	0.42
1:B:49:LEU:HD12	1:B:54:ASP:HB2	2.01	0.42
4:C:179:PEG:H11	6:C:287:HOH:O	2.20	0.42
1:C:147[B]:ARG:HB2	2:C:176:CL:CL	2.57	0.41
1:A:78:ALA:HA	1:A:100:PHE:O	2.21	0.41
1:A:126:ALA:HB1	1:A:153:ASP:O	2.21	0.41
1:A:85:TYR:CE1	1:A:108[B]:MSE:HG3	2.56	0.41
1:A:91:SER:HB3	1:A:99[B]:LEU:HD21	2.03	0.41
1:D:49:LEU:HD12	1:D:54:ASP:HB2	2.03	0.40
1:C:108[B]:MSE:HG3	1:C:108[B]:MSE:O	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:147[B]:ARG:HD2	1:D:151:VAL:HG12	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 8 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PEG	C	180	-	6,6,6	0.47	0	5,5,5	0.39	0
4	PEG	C	179	-	6,6,6	0.50	0	5,5,5	0.94	0
4	PEG	D	176	-	6,6,6	0.59	0	5,5,5	0.46	0
5	P6G	C	178	-	18,18,18	0.62	0	17,17,17	0.82	0
3	TLA	B	179	-	3,9,9	0.59	0	6,12,12	1.23	1 (16%)
4	PEG	B	180	-	6,6,6	0.46	0	5,5,5	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	C	180	-	-	2/4/4/4	-
4	PEG	C	179	-	-	2/4/4/4	-
4	PEG	D	176	-	-	3/4/4/4	-
5	P6G	C	178	-	-	11/16/16/16	-
3	TLA	B	179	-	-	0/4/12/12	-
4	PEG	B	180	-	-	1/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	179	TLA	C1-C2-C3	-2.07	108.64	113.11

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	178	P6G	C6-C5-O4-C3
5	C	178	P6G	C9-C8-O7-C6
5	C	178	P6G	C18-C17-O16-C15
4	D	176	PEG	O2-C3-C4-O4
4	C	179	PEG	O2-C3-C4-O4
4	C	180	PEG	O1-C1-C2-O2
5	C	178	P6G	O4-C5-C6-O7
5	C	178	P6G	C15-C14-O13-C12
5	C	178	P6G	O7-C8-C9-O10
4	D	176	PEG	C1-C2-O2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	C	178	P6G	O1-C2-C3-O4
5	C	178	P6G	C2-C3-O4-C5
4	B	180	PEG	O1-C1-C2-O2
4	D	176	PEG	O1-C1-C2-O2
4	C	179	PEG	C1-C2-O2-C3
4	C	180	PEG	C1-C2-O2-C3
5	C	178	P6G	O16-C17-C18-O19
5	C	178	P6G	O13-C14-C15-O16
5	C	178	P6G	O10-C11-C12-O13

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	179	PEG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	173/176 (98%)	-0.21	2 (1%) 79 83	9, 14, 24, 35	0
1	B	173/176 (98%)	-0.13	1 (0%) 89 91	9, 15, 25, 34	0
1	C	171/176 (97%)	-0.17	0 100 100	9, 15, 25, 31	0
1	D	169/176 (96%)	0.09	4 (2%) 59 65	9, 15, 26, 31	0
All	All	686/704 (97%)	-0.11	7 (1%) 82 85	9, 15, 26, 35	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	70	THR	5.2
1	A	49	LEU	3.3
1	D	71	GLU	3.0
1	D	110	PRO	3.0
1	D	95	PRO	2.7
1	B	50	GLY	2.5
1	A	50	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PEG	D	176	7/7	0.63	0.16	44,46,55,56	0
5	P6G	C	178	19/19	0.70	0.29	25,39,54,57	0
4	PEG	C	180	7/7	0.74	0.13	47,52,54,57	0
4	PEG	C	179	7/7	0.79	0.21	27,39,42,42	0
4	PEG	B	180	7/7	0.89	0.12	28,35,41,43	0
2	CL	B	178	1/1	0.92	0.28	37,37,37,37	0
3	TLA	B	179	10/10	0.95	0.06	16,20,23,25	0
2	CL	B	177	1/1	0.97	0.10	31,31,31,31	0
2	CL	C	177	1/1	0.98	0.12	28,28,28,28	0
2	CL	A	178	1/1	0.98	0.07	23,23,23,23	0
2	CL	C	176	1/1	0.99	0.06	18,18,18,18	0
2	CL	A	177	1/1	0.99	0.04	17,17,17,17	0
2	CL	A	176	1/1	0.99	0.09	29,29,29,29	0
2	CL	B	176	1/1	0.99	0.04	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around P6G C 178:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.