



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 19, 2020 – 07:41 pm BST

PDB ID : 3QJX
Title : Crystal Structure of E. coli Aminopeptidase N in complex with L-Serine
Authors : Addlagatta, A.; Gumpena, R.; Kishor, C.; Ganji, R.J.
Deposited on : 2011-01-31
Resolution : 1.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

i

X-RAY DIFFRACTION

A.

Ramachandran outliers16

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	A	977	-	-	X	-
6	GOL	A	978	-	X	-	-
6	GOL	A	979	-	X	-	-
6	GOL	A	982	-	-	X	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8314 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminopeptidase N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	866	Total	C	N	O	S	0	62	0
			7296	4638	1253	1377	28			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP P04825
A	-19	GLY	-	EXPRESSION TAG	UNP P04825
A	-18	SER	-	EXPRESSION TAG	UNP P04825
A	-17	SER	-	EXPRESSION TAG	UNP P04825
A	-16	HIS	-	EXPRESSION TAG	UNP P04825
A	-15	HIS	-	EXPRESSION TAG	UNP P04825
A	-14	HIS	-	EXPRESSION TAG	UNP P04825
A	-13	HIS	-	EXPRESSION TAG	UNP P04825
A	-12	HIS	-	EXPRESSION TAG	UNP P04825
A	-11	HIS	-	EXPRESSION TAG	UNP P04825
A	-10	SER	-	EXPRESSION TAG	UNP P04825
A	-9	SER	-	EXPRESSION TAG	UNP P04825
A	-8	GLY	-	EXPRESSION TAG	UNP P04825
A	-7	GLU	-	EXPRESSION TAG	UNP P04825
A	-6	ASN	-	EXPRESSION TAG	UNP P04825
A	-5	LEU	-	EXPRESSION TAG	UNP P04825
A	-4	TYR	-	EXPRESSION TAG	UNP P04825
A	-3	PHE	-	EXPRESSION TAG	UNP P04825
A	-2	GLN	-	EXPRESSION TAG	UNP P04825
A	-1	GLY	-	EXPRESSION TAG	UNP P04825
A	0	HIS	-	EXPRESSION TAG	UNP P04825


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Na	0	0
			3	3		

- Molecule 4 is SERINE (three-letter code: SER) (formula: C₃H₇NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			7	3	1	3		

- Molecule 5 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	922	Total	O	0	0
			922	922		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aminopeptidase N



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	120.23Å 120.23Å 170.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.74 – 1.45 35.74 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.9 (35.74-1.45) 99.9 (35.74-1.45)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.120 , 0.147 0.119 , 0.146	Depositor DCC
R_{free} test set	7641 reflections (3.04%)	wwPDB-VP
Wilson B-factor (Å ²)	12.1	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for -h,-k,l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	8314	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, MLI, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.54	73/7629 (1.0%)	1.42	99/10345 (1.0%)

The worst 5 of 73 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	231	LYS	CE-NZ	9.03	1.71	1.49
1	A	606	GLU	CG-CD	9.01	1.65	1.51
1	A	207	ARG	CD-NE	8.47	1.60	1.46
1	A	585	GLN	CG-CD	8.47	1.70	1.51
1	A	763	GLU	CG-CD	8.22	1.64	1.51

The worst 5 of 99 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	349[A]	ARG	NE-CZ-NH2	-12.18	114.21	120.30
1	A	349[C]	ARG	NE-CZ-NH2	-12.18	114.21	120.30
1	A	681	ARG	NE-CZ-NH2	11.82	126.21	120.30
1	A	279	ARG	NE-CZ-NH1	-10.73	114.94	120.30
1	A	832	ARG	NE-CZ-NH2	-10.45	115.07	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7296	0	7277	143	1
2	A	1	0	0	0	0
3	A	3	0	0	0	0
4	A	7	0	4	0	0
5	A	7	0	2	0	0
6	A	78	0	76	31	0
7	A	922	0	0	44	1
All	All	8314	0	7359	151	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 151 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:LYS:NZ	1:A:231:LYS:CE	1.71	1.48
1:A:825[B]:ARG:NH2	7:A:1776:HOH:O	1.58	1.31
1:A:627[A]:GLU:HG3	7:A:1384:HOH:O	1.23	1.27
1:A:783[B]:ARG:NH2	1:A:825[B]:ARG:HD2	1.49	1.26
1:A:783[B]:ARG:HH21	1:A:825[B]:ARG:CD	1.52	1.23

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204[B]:ARG:NE	7:A:963:HOH:O[4_565]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	926/891 (104%)	912 (98%)	14 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	802/763 (105%)	791 (99%)	11 (1%)	67 37

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	356	ASP
1	A	507[A]	ASN
1	A	741[A]	GLN
1	A	327	ASP
1	A	606	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	569	GLN
1	A	623	ASN
1	A	682	ASN
1	A	514	GLN
1	A	665	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 4 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	A	978	-	5,5,5	1.24	0	5,5,5	2.28	3 (60%)
6	GOL	A	977	-	5,5,5	0.75	0	5,5,5	2.23	2 (40%)
6	GOL	A	974	-	5,5,5	1.68	2 (40%)	5,5,5	1.32	1 (20%)
5	MLI	A	950	-	0,6,6	0.00	-	0,7,7	0.00	-
6	GOL	A	973	-	5,5,5	1.00	0	5,5,5	1.56	1 (20%)
6	GOL	A	970	-	5,5,5	2.18	2 (40%)	5,5,5	2.24	2 (40%)
6	GOL	A	979	-	5,5,5	0.95	0	5,5,5	1.79	2 (40%)
6	GOL	A	983	-	5,5,5	0.94	0	5,5,5	1.14	0
6	GOL	A	975	-	5,5,5	1.24	0	5,5,5	1.37	1 (20%)
6	GOL	A	971	-	5,5,5	0.81	0	5,5,5	0.86	0
6	GOL	A	976	-	5,5,5	1.24	1 (20%)	5,5,5	1.75	1 (20%)
6	GOL	A	982	-	5,5,5	0.87	0	5,5,5	1.70	1 (20%)
6	GOL	A	972	-	5,5,5	3.08	3 (60%)	5,5,5	2.34	2 (40%)
6	GOL	A	981	-	5,5,5	0.55	0	5,5,5	1.51	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	978	-	-	3/4/4/4	-
6	GOL	A	977	-	-	3/4/4/4	-
6	GOL	A	974	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MLI	A	950	-	-	0/0/4/4	-
6	GOL	A	973	-	-	4/4/4/4	-
6	GOL	A	970	-	-	2/4/4/4	-
6	GOL	A	979	-	-	4/4/4/4	-
6	GOL	A	983	-	-	2/4/4/4	-
6	GOL	A	975	-	-	1/4/4/4	-
6	GOL	A	971	-	-	0/4/4/4	-
6	GOL	A	976	-	-	2/4/4/4	-
6	GOL	A	982	-	-	2/4/4/4	-
6	GOL	A	972	-	-	3/4/4/4	-
6	GOL	A	981	-	-	3/4/4/4	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	972	GOL	O2-C2	-5.21	1.27	1.43
6	A	972	GOL	C3-C2	3.47	1.66	1.51
6	A	970	GOL	O2-C2	3.18	1.52	1.43
6	A	972	GOL	O3-C3	2.87	1.54	1.42
6	A	970	GOL	C1-C2	2.71	1.62	1.51

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	972	GOL	O1-C1-C2	-4.06	90.73	110.20
6	A	970	GOL	C3-C2-C1	-3.76	97.08	111.70
6	A	978	GOL	O2-C2-C1	-3.63	93.14	109.12
6	A	977	GOL	O2-C2-C1	3.60	124.96	109.12
6	A	982	GOL	O2-C2-C3	3.29	123.60	109.12

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	978	GOL	C1-C2-C3-O3
6	A	978	GOL	O2-C2-C3-O3
6	A	977	GOL	O1-C1-C2-C3
6	A	973	GOL	O1-C1-C2-C3
6	A	973	GOL	C1-C2-C3-O3

There are no ring outliers.

9 monomers are involved in 31 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	978	GOL	3	0
6	A	977	GOL	16	0
6	A	974	GOL	1	0
6	A	973	GOL	1	0
6	A	979	GOL	3	0
6	A	983	GOL	3	0
6	A	976	GOL	1	0
6	A	982	GOL	6	0
6	A	972	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	866/891 (97%)	-0.26	13 (1%) 73 74	6, 12, 25, 46	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	504	HIS	5.6
1	A	503	GLY	5.1
1	A	169	ASN	4.2
1	A	168	GLU	3.7
1	A	502	GLY	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	A	982	6/6	0.76	0.48	31,34,35,35	6
6	GOL	A	983	6/6	0.90	0.17	27,33,35,35	6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	GOL	A	979	6/6	0.91	0.17	21,26,27,27	6
6	GOL	A	977	6/6	0.92	0.29	27,29,30,35	6
5	MLI	A	950	7/7	0.93	0.22	20,26,31,32	0
6	GOL	A	981	6/6	0.93	0.15	24,27,32,34	6
6	GOL	A	972	6/6	0.94	0.23	21,25,30,30	0
6	GOL	A	978	6/6	0.94	0.14	11,21,25,33	6
6	GOL	A	976	6/6	0.94	0.55	12,23,30,31	6
6	GOL	A	970	6/6	0.95	0.12	17,25,26,28	0
6	GOL	A	973	6/6	0.96	0.12	17,36,49,58	0
6	GOL	A	974	6/6	0.97	0.14	16,17,22,26	6
6	GOL	A	971	6/6	0.97	0.09	19,27,31,37	0
6	GOL	A	975	6/6	0.98	0.15	9,13,15,18	6
4	SER	A	900	7/7	0.99	0.08	7,7,9,10	0
3	NA	A	892	1/1	0.99	0.15	21,21,21,21	1
3	NA	A	891	1/1	0.99	0.15	28,28,28,28	0
2	ZN	A	880	1/1	1.00	0.05	6,6,6,6	0
3	NA	A	890	1/1	1.00	0.07	15,15,15,15	0

6.5 Other polymers ⓘ

There are no such residues in this entry.