



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 11:48 am BST

PDB ID : 3QKA  
Title : Crystal structure of enoyl-CoA hydratase EchA5 from *Mycobacterium marinum*  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2011-01-31  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	<div> <div>84%</div> <div>5%</div> <div>11%</div> </div>
1	B	261	<div> <div>88%</div> <div>10%</div> </div>
1	C	261	<div> <div>3%</div> <div>88%</div> <div>7%</div> </div>
1	D	261	<div> <div>5%</div> <div>89%</div> <div>7%</div> <div>5%</div> </div>
1	E	261	<div> <div>3%</div> <div>87%</div> <div>5%</div> <div>7%</div> </div>
1	F	261	<div> <div>2%</div> <div>88%</div> <div>8%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11036 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enoyl-CoA hydratase, EchA5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	233	Total	C	N	O	S	0	0	0
			1697	1058	314	315	10			
1	B	236	Total	C	N	O	S	0	3	0
			1708	1067	309	321	11			
1	C	243	Total	C	N	O	S	0	0	0
			1732	1076	319	327	10			
1	D	248	Total	C	N	O	S	0	2	0
			1770	1103	326	331	10			
1	E	243	Total	C	N	O	S	0	0	0
			1734	1077	317	329	11			
1	F	239	Total	C	N	O	S	0	0	0
			1723	1074	315	324	10			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP B2HSK3
A	-3	PRO	-	EXPRESSION TAG	UNP B2HSK3
A	-2	GLY	-	EXPRESSION TAG	UNP B2HSK3
A	-1	SER	-	EXPRESSION TAG	UNP B2HSK3
A	0	MET	-	EXPRESSION TAG	UNP B2HSK3
A	1	VAL	-	EXPRESSION TAG	UNP B2HSK3
B	-4	GLY	-	EXPRESSION TAG	UNP B2HSK3
B	-3	PRO	-	EXPRESSION TAG	UNP B2HSK3
B	-2	GLY	-	EXPRESSION TAG	UNP B2HSK3
B	-1	SER	-	EXPRESSION TAG	UNP B2HSK3
B	0	MET	-	EXPRESSION TAG	UNP B2HSK3
B	1	VAL	-	EXPRESSION TAG	UNP B2HSK3
C	-4	GLY	-	EXPRESSION TAG	UNP B2HSK3
C	-3	PRO	-	EXPRESSION TAG	UNP B2HSK3
C	-2	GLY	-	EXPRESSION TAG	UNP B2HSK3
C	-1	SER	-	EXPRESSION TAG	UNP B2HSK3
C	0	MET	-	EXPRESSION TAG	UNP B2HSK3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	VAL	-	EXPRESSION TAG	UNP B2HSK3
D	-4	GLY	-	EXPRESSION TAG	UNP B2HSK3
D	-3	PRO	-	EXPRESSION TAG	UNP B2HSK3
D	-2	GLY	-	EXPRESSION TAG	UNP B2HSK3
D	-1	SER	-	EXPRESSION TAG	UNP B2HSK3
D	0	MET	-	EXPRESSION TAG	UNP B2HSK3
D	1	VAL	-	EXPRESSION TAG	UNP B2HSK3
E	-4	GLY	-	EXPRESSION TAG	UNP B2HSK3
E	-3	PRO	-	EXPRESSION TAG	UNP B2HSK3
E	-2	GLY	-	EXPRESSION TAG	UNP B2HSK3
E	-1	SER	-	EXPRESSION TAG	UNP B2HSK3
E	0	MET	-	EXPRESSION TAG	UNP B2HSK3
E	1	VAL	-	EXPRESSION TAG	UNP B2HSK3
F	-4	GLY	-	EXPRESSION TAG	UNP B2HSK3
F	-3	PRO	-	EXPRESSION TAG	UNP B2HSK3
F	-2	GLY	-	EXPRESSION TAG	UNP B2HSK3
F	-1	SER	-	EXPRESSION TAG	UNP B2HSK3
F	0	MET	-	EXPRESSION TAG	UNP B2HSK3
F	1	VAL	-	EXPRESSION TAG	UNP B2HSK3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	104	Total O 104 104	0	0
2	B	94	Total O 94 94	0	0
2	C	130	Total O 130 130	0	0
2	D	125	Total O 125 125	0	0
2	E	107	Total O 107 107	0	0
2	F	112	Total O 112 112	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Enoyl-CoA hydratase, EchA5

Chain A: 




- Molecule 1: Enoyl-CoA hydratase, EchA5

Chain B: 





- Molecule 1: Enoyl-CoA hydratase, EchA5

Chain C: 




- Molecule 1: Enoyl-CoA hydratase, EchA5

Chain D: 




- Molecule 1: Enoyl-CoA hydratase, EchA5

Chain E: 



● Molecule 1: Enoyl-CoA hydratase, EchA5



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.04Å 118.92Å 132.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.15 41.35 – 2.15	Depositor EDS
% Data completeness (in resolution range)	97.1 (50.00-2.15) 97.1 (41.35-2.15)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.50 (at 2.16Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.168 , 0.213 0.175 , 0.222	Depositor DCC
$R_{free}$ test set	3698 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.1	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11036	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	1/1725 (0.1%)	0.69	1/2347 (0.0%)
1	B	0.68	0/1745	0.71	0/2378
1	C	0.75	1/1759 (0.1%)	0.75	2/2396 (0.1%)
1	D	0.73	0/1801	0.67	0/2453
1	E	0.68	0/1761	0.71	0/2399
1	F	0.68	0/1751	0.71	0/2384
All	All	0.70	2/10542 (0.0%)	0.71	3/14357 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	202	CYS	CB-SG	-7.06	1.70	1.82
1	A	202	CYS	CB-SG	-6.45	1.71	1.82

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	20	ARG	NE-CZ-NH2	-8.16	116.22	120.30
1	C	20	ARG	NE-CZ-NH1	6.93	123.77	120.30
1	A	20	ARG	NE-CZ-NH2	-5.25	117.68	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1697	0	1695	10	0
1	B	1708	0	1688	4	0
1	C	1732	0	1699	7	0
1	D	1770	0	1744	11	0
1	E	1734	0	1710	10	0
1	F	1723	0	1706	6	0
2	A	104	0	0	0	0
2	B	94	0	0	0	0
2	C	130	0	0	1	0
2	D	125	0	0	0	0
2	E	107	0	0	1	0
2	F	112	0	0	0	0
All	All	11036	0	10242	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:121:GLU:HG3	1:F:184:ALA:HB2	1.76	0.67
1:C:121:GLU:HG3	1:C:184:ALA:HB2	1.79	0.64
1:F:180:PRO:HB2	1:F:183:GLN:HG3	1.82	0.61
1:D:121:GLU:HG3	1:D:184:ALA:HB2	1.82	0.61
1:B:121:GLU:HG3	1:B:184:ALA:HB2	1.84	0.60
1:C:165:GLN:HB3	2:C:297:HOH:O	2.01	0.60
1:A:121:GLU:HG3	1:A:184:ALA:HB2	1.85	0.59
1:C:4:LEU:HD22	1:C:20:ARG:HD3	1.87	0.57
1:A:4:LEU:HD22	1:A:20:ARG:HD3	1.89	0.55
1:A:136:PRO:HA	1:B:202[B]:CYS:SG	2.49	0.53
1:A:51:ALA:HB3	1:A:96:VAL:HG22	1.92	0.51
1:F:130:CYS:HB2	1:F:135:VAL:O	2.10	0.51
1:D:121:GLU:CG	1:D:184:ALA:HB2	2.40	0.50
1:E:153:ARG:HD3	1:E:172:ILE:HB	1.92	0.50
1:A:121:GLU:CG	1:A:184:ALA:HB2	2.41	0.50
1:A:230:SER:HB3	1:C:231:ARG:HA	1.94	0.49
1:E:232:VAL:HG12	1:E:232:VAL:O	2.13	0.48
1:A:131:ARG:HA	1:A:131:ARG:NH1	2.28	0.47
1:D:123:ASP:OD2	1:D:181:ASN:ND2	2.36	0.47
1:C:151:HIS:HB3	1:D:148[A]:LEU:HD23	1.96	0.46
1:B:120:ALA:O	1:B:178:VAL:HA	2.16	0.46
1:F:131:ARG:NH1	1:F:131:ARG:HA	2.31	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:198:LEU:HD13	1:E:131:ARG:CG	2.47	0.45
1:D:109:LEU:O	1:D:113:LEU:HG	2.16	0.45
1:D:120:ALA:O	1:D:178:VAL:HA	2.16	0.45
1:F:203:LEU:C	1:F:203:LEU:HD23	2.36	0.45
1:E:14:THR:O	1:E:51:ALA:HA	2.18	0.44
1:E:202:CYS:HB3	2:E:281:HOH:O	2.17	0.44
1:E:121:GLU:CG	1:E:184:ALA:HB2	2.47	0.44
1:A:128:VAL:O	1:A:128:VAL:HG12	2.18	0.43
1:D:133:TRP:O	1:E:239:GLY:HA3	2.18	0.43
1:C:66:LEU:HD11	1:C:133:TRP:CE3	2.53	0.43
1:A:217:PRO:HB3	1:D:217:PRO:HG3	2.00	0.43
1:D:14:THR:O	1:D:51:ALA:HA	2.19	0.43
1:D:151:HIS:HB3	1:E:148:LEU:HD23	2.00	0.42
1:A:203:LEU:C	1:A:203:LEU:HD23	2.40	0.42
1:B:131:ARG:HD2	1:F:203:LEU:HD12	2.02	0.41
1:E:145:LEU:HB3	1:E:146:PRO:HD3	2.02	0.41
1:D:28:ASN:HB2	1:D:73:GLU:O	2.21	0.41
1:E:205:SER:HB3	1:E:228:SER:OG	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/261 (88%)	222 (96%)	9 (4%)	0	100	100
1	B	237/261 (91%)	227 (96%)	10 (4%)	0	100	100
1	C	241/261 (92%)	232 (96%)	9 (4%)	0	100	100
1	D	248/261 (95%)	242 (98%)	6 (2%)	0	100	100
1	E	241/261 (92%)	228 (95%)	13 (5%)	0	100	100
1	F	237/261 (91%)	231 (98%)	6 (2%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1435/1566 (92%)	1382 (96%)	53 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/184 (91%)	165 (99%)	2 (1%)	71	76
1	B	167/184 (91%)	167 (100%)	0	100	100
1	C	165/184 (90%)	162 (98%)	3 (2%)	59	63
1	D	167/184 (91%)	166 (99%)	1 (1%)	86	90
1	E	168/184 (91%)	167 (99%)	1 (1%)	86	90
1	F	167/184 (91%)	165 (99%)	2 (1%)	71	76
All	All	1001/1104 (91%)	992 (99%)	9 (1%)	78	83

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	ARG
1	A	205	SER
1	C	59	THR
1	C	133	TRP
1	C	205	SER
1	D	132	ARG
1	E	131	ARG
1	F	46	GLU
1	F	82	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	233/261 (89%)	-0.25	1 (0%) 92 94	15, 23, 37, 47	0
1	B	236/261 (90%)	-0.12	1 (0%) 92 94	14, 27, 43, 53	0
1	C	243/261 (93%)	-0.16	7 (2%) 51 61	12, 20, 35, 57	0
1	D	248/261 (95%)	-0.07	14 (5%) 24 33	12, 20, 43, 51	0
1	E	243/261 (93%)	0.01	9 (3%) 41 49	13, 25, 48, 58	0
1	F	239/261 (91%)	-0.24	5 (2%) 63 71	13, 23, 37, 54	0
All	All	1442/1566 (92%)	-0.14	37 (2%) 56 64	12, 23, 41, 58	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	241	GLY	4.2
1	D	242	ARG	4.2
1	F	72	ALA	3.9
1	E	236	ALA	3.8
1	E	237	LEU	3.7
1	D	244	ALA	3.7
1	F	69	PHE	3.4
1	E	1	VAL	3.4
1	D	246	GLY	3.4
1	D	237	LEU	3.3
1	D	215[A]	GLY	3.2
1	E	-1	SER	3.1
1	D	243	PHE	3.1
1	C	242	ARG	3.0
1	E	0	MET	2.9
1	E	233	ALA	2.9
1	E	70	GLY	2.8
1	F	70	GLY	2.8
1	E	240	ALA	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	245	ALA	2.6
1	E	2	SER	2.5
1	D	69	PHE	2.5
1	C	243	PHE	2.4
1	B	232	VAL	2.3
1	A	216	LEU	2.3
1	C	234	ALA	2.3
1	F	240	ALA	2.3
1	C	238	GLU	2.3
1	D	236	ALA	2.3
1	C	1	VAL	2.2
1	D	133	TRP	2.2
1	C	233	ALA	2.2
1	D	249	ARG	2.2
1	F	66	LEU	2.1
1	D	240	ALA	2.1
1	D	47	SER	2.1
1	D	9	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.