



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2020 – 10:29 AM BST

PDB ID : 6QMV  
Title : E87D sulfur oxygenase reductase  
Authors : Frazao, C.; Klezin, A.; Veith, A.  
Deposited on : 2019-02-08  
Resolution : 1.87 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	<div> <div>87%</div> <div>9%</div> <div>.</div> </div>
1	B	318	<div> <div>%</div> <div>91%</div> <div>5%</div> <div>.</div> </div>
1	C	318	<div> <div>%</div> <div>90%</div> <div>7%</div> <div>.</div> </div>
1	D	318	<div> <div>%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
1	E	318	<div> <div>%</div> <div>92%</div> <div>5%</div> <div>.</div> </div>
1	F	318	<div> <div>%</div> <div>90%</div> <div>7%</div> <div>.</div> </div>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 16476 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sulfur oxygenase/reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	307	Total	C	N	O	S	0	10	0
			2502	1619	412	446	25			
1	B	307	Total	C	N	O	S	0	10	0
			2498	1615	412	448	23			
1	C	307	Total	C	N	O	S	0	9	0
			2503	1619	415	446	23			
1	D	307	Total	C	N	O	S	0	11	0
			2508	1623	413	448	24			
1	E	307	Total	C	N	O	S	0	7	0
			2488	1609	409	446	24			
1	F	307	Total	C	N	O	S	0	11	0
			2504	1621	412	447	24			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	87	ASP	GLU	engineered mutation	UNP P29082
A	309	ASN	-	expression tag	UNP P29082
A	310	ALA	-	expression tag	UNP P29082
A	311	TRP	-	expression tag	UNP P29082
A	312	ARG	-	expression tag	UNP P29082
A	313	HIS	-	expression tag	UNP P29082
A	314	PRO	-	expression tag	UNP P29082
A	315	GLN	-	expression tag	UNP P29082
A	316	PHE	-	expression tag	UNP P29082
A	317	GLY	-	expression tag	UNP P29082
A	318	GLY	-	expression tag	UNP P29082
B	87	ASP	GLU	engineered mutation	UNP P29082
B	309	ASN	-	expression tag	UNP P29082
B	310	ALA	-	expression tag	UNP P29082
B	311	TRP	-	expression tag	UNP P29082
B	312	ARG	-	expression tag	UNP P29082
B	313	HIS	-	expression tag	UNP P29082

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	314	PRO	-	expression tag	UNP P29082
B	315	GLN	-	expression tag	UNP P29082
B	316	PHE	-	expression tag	UNP P29082
B	317	GLY	-	expression tag	UNP P29082
B	318	GLY	-	expression tag	UNP P29082
C	87	ASP	GLU	engineered mutation	UNP P29082
C	309	ASN	-	expression tag	UNP P29082
C	310	ALA	-	expression tag	UNP P29082
C	311	TRP	-	expression tag	UNP P29082
C	312	ARG	-	expression tag	UNP P29082
C	313	HIS	-	expression tag	UNP P29082
C	314	PRO	-	expression tag	UNP P29082
C	315	GLN	-	expression tag	UNP P29082
C	316	PHE	-	expression tag	UNP P29082
C	317	GLY	-	expression tag	UNP P29082
C	318	GLY	-	expression tag	UNP P29082
D	87	ASP	GLU	engineered mutation	UNP P29082
D	309	ASN	-	expression tag	UNP P29082
D	310	ALA	-	expression tag	UNP P29082
D	311	TRP	-	expression tag	UNP P29082
D	312	ARG	-	expression tag	UNP P29082
D	313	HIS	-	expression tag	UNP P29082
D	314	PRO	-	expression tag	UNP P29082
D	315	GLN	-	expression tag	UNP P29082
D	316	PHE	-	expression tag	UNP P29082
D	317	GLY	-	expression tag	UNP P29082
D	318	GLY	-	expression tag	UNP P29082
E	87	ASP	GLU	engineered mutation	UNP P29082
E	309	ASN	-	expression tag	UNP P29082
E	310	ALA	-	expression tag	UNP P29082
E	311	TRP	-	expression tag	UNP P29082
E	312	ARG	-	expression tag	UNP P29082
E	313	HIS	-	expression tag	UNP P29082
E	314	PRO	-	expression tag	UNP P29082
E	315	GLN	-	expression tag	UNP P29082
E	316	PHE	-	expression tag	UNP P29082
E	317	GLY	-	expression tag	UNP P29082
E	318	GLY	-	expression tag	UNP P29082
F	87	ASP	GLU	engineered mutation	UNP P29082
F	309	ASN	-	expression tag	UNP P29082
F	310	ALA	-	expression tag	UNP P29082
F	311	TRP	-	expression tag	UNP P29082

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	312	ARG	-	expression tag	UNP P29082
F	313	HIS	-	expression tag	UNP P29082
F	314	PRO	-	expression tag	UNP P29082
F	315	GLN	-	expression tag	UNP P29082
F	316	PHE	-	expression tag	UNP P29082
F	317	GLY	-	expression tag	UNP P29082
F	318	GLY	-	expression tag	UNP P29082


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	255	Total 255	O 255	0	0
2	B	245	Total 245	O 245	0	0
2	C	234	Total 234	O 234	0	0
2	D	240	Total 240	O 240	0	0
2	E	244	Total 244	O 244	0	0
2	F	255	Total 255	O 255	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sulfur oxygenase/reductase




- Molecule 1: Sulfur oxygenase/reductase



- Molecule 1: Sulfur oxygenase/reductase




- Molecule 1: Sulfur oxygenase/reductase



- Molecule 1: Sulfur oxygenase/reductase



• Molecule 1: Sulfur oxygenase/reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	161.92Å 161.92Å 154.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.12 – 1.87 43.12 – 1.87	Depositor EDS
% Data completeness (in resolution range)	99.1 (43.12-1.87) 99.1 (43.12-1.87)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 1.87Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, $R_{free}$	0.159 , 0.193 0.159 , 0.192	Depositor DCC
$R_{free}$ test set	1986 reflections (1.22%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.1	Xtriage
Anisotropy	0.236	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 46.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h 0.007 for -l,-k,-h 0.008 for -h,-l,-k 0.000 for -h,l,k 0.019 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	16476	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSS, TSY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2602	0.58	1/3523 (0.0%)
1	B	0.38	0/2598	0.56	0/3519
1	C	0.40	0/2597	0.56	0/3517
1	D	0.38	0/2610	0.56	0/3534
1	E	0.37	0/2575	0.54	0/3488
1	F	0.38	0/2607	0.55	0/3531
All	All	0.38	0/15589	0.56	1/21112 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	LEU	CA-CB-CG	7.64	132.86	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2502	0	2460	20	0
1	B	2498	0	2448	12	0
1	C	2503	0	2463	15	0
1	D	2508	0	2465	18	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2488	0	2439	10	0
1	F	2504	0	2461	15	0
2	A	255	0	0	0	2
2	B	245	0	0	0	0
2	C	234	0	0	0	0
2	D	240	0	0	1	0
2	E	244	0	0	2	2
2	F	255	0	0	0	0
All	All	16476	0	14736	79	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:ARG:NH2	1:A:232:ASN:OD1	1.98	0.96
1:A:178[B]:ASP:OD1	1:A:182[B]:ARG:NH1	2.08	0.86
1:B:178[B]:ASP:OD1	1:B:182[B]:ARG:NH1	2.09	0.86
1:C:223:ASN:ND2	1:C:227:LEU:O	2.09	0.85
1:C:91:ARG:NH2	1:C:232:ASN:OD1	2.13	0.80

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:621:HOH:O	2:E:434:HOH:O[6_564]	2.15	0.05
2:A:424:HOH:O	2:E:431:HOH:O[6_564]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/318 (98%)	307 (98%)	6 (2%)	0	100	100
1	B	313/318 (98%)	307 (98%)	6 (2%)	0	100	100
1	C	312/318 (98%)	305 (98%)	7 (2%)	0	100	100
1	D	314/318 (99%)	307 (98%)	7 (2%)	0	100	100
1	E	310/318 (98%)	303 (98%)	7 (2%)	0	100	100
1	F	314/318 (99%)	308 (98%)	6 (2%)	0	100	100
All	All	1876/1908 (98%)	1837 (98%)	39 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/271 (101%)	271 (99%)	2 (1%)	84	83
1	B	273/271 (101%)	273 (100%)	0	100	100
1	C	272/271 (100%)	272 (100%)	0	100	100
1	D	274/271 (101%)	274 (100%)	0	100	100
1	E	270/271 (100%)	269 (100%)	1 (0%)	91	90
1	F	274/271 (101%)	274 (100%)	0	100	100
All	All	1636/1626 (101%)	1633 (100%)	3 (0%)	93	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	VAL
1	A	260	LEU
1	E	167	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSS	F	31	1	4,6,7	1.16	0	1,6,8	1.37	0
1	CSS	A	31	1	4,6,7	1.23	0	1,6,8	0.80	0
1	CSS	B	31	1	4,6,7	1.25	0	1,6,8	1.11	0
1	TSY	C	104	1	5,7,8	1.19	0	2,7,9	0.90	0
1	TSY	A	104	1	5,7,8	1.13	0	2,7,9	1.28	0
1	CSS	D	31	1	4,6,7	1.16	0	1,6,8	1.10	0
1	TSY	E	104	1	5,7,8	1.24	0	2,7,9	1.24	0
1	CSS	C	31	1	4,6,7	1.17	0	1,6,8	1.31	0
1	TSY	F	104	1	5,7,8	1.14	0	2,7,9	1.22	0
1	TSY	D	104	1	5,7,8	1.21	0	2,7,9	2.38	1 (50%)
1	TSY	B	104	1	5,7,8	1.20	0	2,7,9	1.34	0
1	CSS	E	31	1	4,6,7	1.21	0	1,6,8	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	F	31	1	-	0/1/5/7	-
1	CSS	A	31	1	-	0/1/5/7	-
1	CSS	B	31	1	-	0/1/5/7	-
1	TSY	C	104	1	-	1/2/6/8	-
1	TSY	A	104	1	-	1/2/6/8	-
1	CSS	D	31	1	-	0/1/5/7	-
1	TSY	E	104	1	-	1/2/6/8	-
1	CSS	C	31	1	-	0/1/5/7	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TSY	F	104	1	-	1/2/6/8	-
1	TSY	D	104	1	-	1/2/6/8	-
1	TSY	B	104	1	-	1/2/6/8	-
1	CSS	E	31	1	-	0/1/5/7	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	104	TSY	CB-SG-S1	3.31	112.54	103.82

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	104	TSY	CA-CB-SG-S1
1	A	104	TSY	CA-CB-SG-S1
1	E	104	TSY	CA-CB-SG-S1
1	F	104	TSY	CA-CB-SG-S1
1	D	104	TSY	CA-CB-SG-S1

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	104	TSY	1	0
1	A	104	TSY	1	0
1	D	31	CSS	1	0
1	B	104	TSY	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	305/318 (95%)	-0.50	0 <b>100</b> <b>100</b>	11, 16, 37, 55	0
1	B	305/318 (95%)	-0.46	4 (1%) <b>77</b> <b>79</b>	12, 18, 38, 59	0
1	C	305/318 (95%)	-0.53	3 (0%) <b>82</b> <b>83</b>	12, 18, 39, 58	0
1	D	305/318 (95%)	-0.48	2 (0%) <b>87</b> <b>88</b>	12, 18, 37, 57	0
1	E	305/318 (95%)	-0.44	3 (0%) <b>82</b> <b>83</b>	13, 18, 37, 59	0
1	F	305/318 (95%)	-0.49	2 (0%) <b>87</b> <b>88</b>	12, 18, 35, 61	0
All	All	1830/1908 (95%)	-0.48	14 (0%) <b>86</b> <b>87</b>	11, 18, 38, 61	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	102	TYR	3.7
1	E	144	GLY	2.9
1	E	147	LEU	2.7
1	B	148	ASP	2.6
1	C	17	PRO	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TSY	E	104	8/9	0.75	0.13	35,46,79,81	0
1	TSY	F	104	8/9	0.80	0.22	34,53,99,107	0
1	TSY	D	104	8/9	0.82	0.14	33,48,84,92	0
1	TSY	C	104	8/9	0.85	0.15	41,48,77,81	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	TSY	B	104	8/9	0.85	0.18	35,44,75,80	0
1	TSY	A	104	8/9	0.87	0.13	41,48,73,76	0
1	CSS	D	31	7/8	0.94	0.09	15,17,30,56	0
1	CSS	F	31	7/8	0.94	0.09	16,17,28,56	0
1	CSS	C	31	7/8	0.94	0.09	13,16,31,54	0
1	CSS	E	31	7/8	0.94	0.10	15,16,33,58	0
1	CSS	B	31	7/8	0.95	0.10	15,18,32,57	0
1	CSS	A	31	7/8	0.96	0.09	13,16,31,58	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.